



US010947548B2

(12) **United States Patent**
Deng et al.

(10) **Patent No.:** **US 10,947,548 B2**
(45) **Date of Patent:** **Mar. 16, 2021**

(54) **PRODUCTION OF ORGANIC ACIDS FROM *ASPERGILLUS CIS-ACONITIC ACID* DECARBOXYLASE (CADA) DELETION STRAINS**

(71) Applicant: **Battelle Memorial Institute**, Richland, WA (US)

(72) Inventors: **Shuang Deng**, Richland, WA (US); **Ziyu Dai**, Richland, WA (US); **Jon K. Magnuson**, Richland, WA (US)

(73) Assignee: **Battelle Memorial Institute**, Richland, WA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/393,149**

(22) Filed: **Apr. 24, 2019**

(65) **Prior Publication Data**

US 2019/0323020 A1 Oct. 24, 2019

Related U.S. Application Data

(60) Provisional application No. 62/661,804, filed on Apr. 24, 2018.

(51) **Int. Cl.**

C12N 1/15 (2006.01)
C12N 9/88 (2006.01)
C12N 9/10 (2006.01)
C12N 9/04 (2006.01)
C12P 7/42 (2006.01)
C12P 7/52 (2006.01)
C12N 15/80 (2006.01)
C12P 7/48 (2006.01)

(52) **U.S. Cl.**

CPC **C12N 15/80** (2013.01); **C12N 9/0006** (2013.01); **C12N 9/1096** (2013.01); **C12N 9/88** (2013.01); **C12P 7/42** (2013.01); **C12P 7/48** (2013.01); **C12Y 101/01059** (2013.01); **C12Y 206/01018** (2013.01); **C12Y 401/01011** (2013.01)

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,740,464 A * 4/1988 Holdom C12P 7/48 435/135
2008/0199926 A1* 8/2008 Burgard C12N 15/52 435/141
2015/0267228 A1* 9/2015 Borodina C12N 9/88 435/146

FOREIGN PATENT DOCUMENTS

CN 110527637 A * 12/2019
WO WO 2018/213349 A1 11/2018

OTHER PUBLICATIONS

Samson et al., *Studies in Mycology* 78:141-173, 2014 (Year: 2014).*

Van der Straat et al., *Microb. Cell Fac.* 13:11, 2014, 9 pages (Year: 2014).*

Rodrigues et al., *Fung. Genet. Biol.* 138 (2020) 103367, 11 pages (Year: 2020).*

Deng et al., *Appl. Microbiol. Biotechnol.* 104:3981-3992, 2020 (Year: 2020).*

UniProt Database Accession No. B3IUN8, Oct. 2017, 2 pages (Year: 2017).*

GenBank Database Accession No. AB326105, Aug. 2008, 2 pages (Year: 2008).*

Nielsen, J., *Appl. Microbiol. Biotechnol.* 55:263-283, 2001 (Year: 2001).*

Steiger et al., *Frontiers Microbiol.* 4:23, 2013, 5 pages (Year: 2013).*

Samson et al., *Studies in Mycology* 69:39-55, 2011 (Year: 2011).*

Deng et al., "Deletion Analysis of the Itaconic Acid Production Gene Cluster Components in *Aspergillus pseudoterreus* ATCC32359," Poster presented at 40th Symposium on Biotechnology for Fuels and Chemicals, Apr. 29-May 2, 2018, Clearwater, Florida.

* cited by examiner

Primary Examiner — David Steadman
(74) *Attorney, Agent, or Firm* — Klarquist Sparkman, LLP

(57) **ABSTRACT**

This application provides recombinant *Aspergillus* fungi having an endogenous cis-aconitic acid decarboxylase (cadA) gene genetically inactivated, which allows aconitic acid production by the recombinant fungi. Such recombinant fungi can further include an exogenous nucleic acid molecule encoding aspartate decarboxylase (panD), an exogenous nucleic acid molecule encoding β -alanine-pyruvate aminotransferase (BAPAT), and an exogenous nucleic acid molecule encoding 3-hydroxypropionate dehydrogenase (HPDH). Kits including these fungi, and methods of using these fungi to produce aconitic acid and 3-hydroxypropionic acid (3-HP) are also provided.

15 Claims, 10 Drawing Sheets

Specification includes a Sequence Listing.

FIG. 1




FIG. 2




FIG. 3B




FIG. 3A




FIG. 4




FIG. 5




FIG. 6B




FIG. 6A




FIG. 6C




FIG. 7




FIG. 8




FIG. 9A





FIG. 9B



1

**PRODUCTION OF ORGANIC ACIDS FROM
ASPERGILLUS CIS-ACONITIC ACID
DECARBOXYLASE (CADA) DELETION
STRAINS**

CROSS REFERENCE TO RELATED
APPLICATION

This application claims priority to U.S. Provisional Application No. 62/661,804 filed Apr. 24, 2018, herein incorporated by reference in its entirety.

ACKNOWLEDGMENT OF GOVERNMENT
SUPPORT

This disclosure was made with Government support under Contract DE-AC05-76RL0 1830 awarded by the U.S. Department of Energy. The Government has certain rights in the invention.

FIELD

Recombinant *Aspergillus* genetically inactivated for an endogenous cis-aconitic acid decarboxylase (*cadA*) gene are provided, as are methods of using such recombinant fungi to produce aconitic acid and 3-hydroxypropionic acid (3-HP).

BACKGROUND

Itaconic acid (IA) is utilized as a monomer or co-monomer to form polymers that are used as raw material for plastics, resins, synthetic fibers and elastomers, detergents and cleaners. *Aspergillus terreus* Thom, produces an appreciable amount of itaconic acid when grown in a glucose medium. Cell-free extracts of *Aspergillus terreus* contain cis-aconitic decarboxylase (*cadA*), which can decarboxylate cis-aconitic acid into equal moles of itaconic acid and carbon dioxide.

The itaconic acid gene cluster (IA cluster) includes four genes, including cis-aconitic acid decarboxylase (*cadA*), a predicted transcription factor (*tf*), mitochondrial organic acid transporter (*mttA*), and MFS (Major Facilitator Superfamily) type transporter (*mfsA*) located in plasma membranes. Expression of one or more genes of the IA gene cluster in heterologous hosts, including *E. coli*, *A. niger*, and *S. cerevisiae*, can result in the production of itaconic acid in non itaconic acid host microorganisms.

Characterization and regulation of genes in the IA biosynthesis cluster through gene deletion had not been previously investigated. The inventors used protoplast transformation to delete each gene in the IA cluster in *Aspergillus terreus/Aspergillus pseudoterreus*, which allowed for the effect on cell growth and IA production to be investigated.

SUMMARY

The role of cis-aconitic acid decarboxylase (*cadA*), a predicted transcription factor (*tf*), mitochondrial organic acid transporter (*mttA*), and MFS (Major Facilitator Superfamily) type transporter (*mfsA*) in IA biosynthesis in *A. pseudoterreus* ATCC 32359 is shown herein. Expressed Sequence Tag (EST) analysis showed a similar expression pattern among those four genes distinct from neighboring genes. Systematic gene deletion analysis demonstrated that *tf*, *cadA*, *mttA* and *mfsA* genes in the cluster are essential for IA production. Interestingly, significant amounts of aconitic

2

acid production was detected in the *cadA* deletion strain but not in the other deletion strains.

Based on these observations, a novel recombinant *AcadA Aspergillus* strain is provided, which can be used for aconitic acid and other organic acid production. Provided herein are isolated recombinant fungi (such as *Aspergillus* filamentous fungi) having a gene inactivation (also referred to herein as a gene deletion or functional deletion) of a cis-aconitic acid decarboxylase (*cadA*) gene (referred to herein as *AcadA* strains). In some examples, the *Aspergillus* fungi is *Aspergillus terreus* or *Aspergillus pseudoterreus*, or particular strains thereof (for example *A. pseudoterreus* ATCC32359 and *A. terreus* NRRL 1960). In particular examples, a *AcadA* strain exhibits one or more of the following characteristics: produces at least 2-fold, at least 3-fold, at least 3.5 fold, at least 5-fold, at least 8-fold, or at least 10-fold more total aconitic acid than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus* (for example at day 3, 4, 5, 6, 7, 8, 9 or 10 of production); produces at least 2-fold more cis-aconitic acid at day 5, 6, 7, 8, 9, or 10 of culturing in Riscaldati medium than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus*; produces at least 2-fold, at least 3-fold, at least 5-fold, or at least 10-fold more trans-aconitic acid at day 10 of culturing in Riscaldati medium than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus*; or combinations thereof. In some examples, such increases are relative to *Aspergillus terreus* strain ATCC 32359 grown under the same conditions.

In particular examples, a *AcadA* fungi further includes an exogenous nucleic acid molecule encoding aspartate 1-decarboxylase (*panD*), an exogenous nucleic acid molecule encoding β -alanine-pyruvate aminotransferase (*BAPAT*), and an exogenous nucleic acid molecule encoding 3-hydroxypropionate dehydrogenase (*HPDH*). The *AcadA* fungi expressing *panD*, *BAPAT*, and *HPDH* can be used to produce 3-HP. Such exogenous nucleic acid molecules can be part of one or more exogenous nucleic acid molecules, such as 1, 2 or 3 exogenous nucleic acid molecules. In one example, the exogenous nucleic acid molecule encoding *panD* has at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 53 or 65 and/or encodes a *panD* protein having at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 54. In one example, the exogenous nucleic acid molecule encoding *BAPAT* has at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 55, and/or encodes a *BAPAT* protein having at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 56. In one example, the exogenous nucleic acid molecule encoding *HPDH* has at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 57, and/or encodes a *HPDH* protein having at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 58. Such *panD*, *BAPAT*, and *HPDH* nucleic acid molecules can be part of a vector. In addition, expression of the *panD*, *BAPAT*, and *HPDH* can be driven by one or more promoters.

The endogenous *cadA* gene is genetically inactivated in some examples by a deletion mutation (complete or partial) or by insertional mutation (e.g., by insertion of an antibiotic resistance gene, such as hygromycin). In some examples, prior to its genetic inactivation, the *cadA* gene encodes a protein having at least 80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 50 or 52. In some examples, prior to its genetic inactivation, the *cadA* gene (or a *cadA* coding sequence) comprises at least

80%, at least 90%, at least 95%, at least 98%, or 100% sequence identity to SEQ ID NO: 49, 51, 59 or 92.

Also provided herein are compositions (such as a culture media or fermentation broth) and kits that include a *Aspergillus* Δ cadA strain. Also provided herein are compositions (such as a culture media or fermentation broth) and kits that include an *Aspergillus* Δ cadA strain that also express panD, BAPAT, and HPDH, in some examples such genes are exogenous to the fungi. In some examples, the composition or kit includes Riscaldati medium (such as modified Riscaldati medium with 20 \times trace elements).

Also provided herein are methods of making aconitic acid (such as cis-aconitic acid, trans-aconitic acid, or both) using the disclosed *Aspergillus* Δ cadA strains. For example, such a method can include culturing an isolated Δ cadA *Aspergillus* under conditions that permit the fungus to make aconitic acid, thereby producing aconitic acid. For example, the Δ cadA fungus can be cultured in Riscaldati medium. In some examples, the method further includes isolating the aconitic acid produced, for example isolating it from the culture media or from the fungus.

Also provided herein are methods of making 3-hydroxypropionic acid (3-HP using the disclosed *Aspergillus* Δ cadA strains that also expresses panD, BAPAT, and HPDH (which can be exogenous). For example, such a method can include culturing an isolated Δ cadA *Aspergillus* that also expresses panD, BAPAT, and HPDH under conditions that permit the fungus to make 3-HP, thereby producing 3-HP. For example, the Δ cadA fungus that also expresses panD, BAPAT, and HPDH can be cultured in Riscaldati medium (such as one including 20 \times trace elements). In some examples, the method further includes isolating the 3-HP produced, for example isolating it from the culture media or from the fungus.

The foregoing and other objects and features of the disclosure will become more apparent from the following detailed description, which proceeds with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Hypothesized itaconic acid (IA) production and transport pathway in *Aspergillus pseudoterreus* and *Aspergillus terreus*. Glucose is utilized by *A. terreus* and *A. pseudoterreus* to form pyruvate and is subsequently converted to citric acid for tricarboxylic acid (TCA) cycle in the mitochondria. Citric acid is dehydrated to cis-aconitic acid, which is then transported from mitochondria to cytosol through transporter mttA. In the cytosol, cis-aconitic acid is decarboxylated into itaconic acid and CO₂ by cis-aconitic decarboxylase. Finally, itaconic acid secreted outside of cell through transporters, for example mfsA.

FIG. 2. *Aspergillus pseudoterreus* ATCC 32359 fermentation data for collecting samples for EST sequencing. A 20 L volume of Riscaldati production medium (see Riscaldati et al., *J Biotechnol* 2000, 83:219-230) in a 30 L working volume Sartorius fermenter was inoculated with 10⁶ *A. pseudoterreus* spores per ml. The three samples referred to as "preproduction, production onset and production" were collected at 40, 50 and 62 hours, respectively. Itaconic acid and glucose data are shown on the left y-axis and fungal ash free dry weight (AFDW) is shown on the right y-axis.

FIGS. 3A-3B. *Aspergillus pseudoterreus* IA Cluster Analysis after five days growth in Riscaldati medium. Spores 0.5 \times 10⁸ were inoculated into 50 ml of production media for IA production as described in Riscaldati et al. (*J Biotechnol* 2000, 83:219-230). The cultivation was per-

formed at 30 $^{\circ}$ C. on a rotary shaker at 150 rpm. At the end of five days, samples were obtained for HPLC analysis and biomass measurement. (A) Dry mass measurement of wild type and mutant strains (B) Itaconic acid production of wild type and mutant strains. The average obtained from three independent experiments are shown. Error bars represent standard deviations from the means.

FIG. 4. Kinetics of itaconic acid production by wild type *A. pseudoterreus* and Δ tfr strains grown in production media at 30 $^{\circ}$ C. Spores 0.5 \times 10⁸ were inoculated into 50 ml of production media for itaconic acid production as described in Riscaldati et al. (*J Biotechnol* 2000, 83:219-230). The cultivation was performed at 30 $^{\circ}$ C. on a rotary shaker at 150 rpm. All experiments were done in three replicates. At day 2, 4, 6, and 7, HPLC analysis was performed to determine amount of IA produced. Each sample was measured in five replicates. Error bars represent standard deviation from the means.

FIG. 5. Real-time (RT)-PCR analysis of the relative levels of mttA, cadA, mfsA mRNAs in wild type and Δ tfr strains. Spores 0.5 \times 10⁸ were inoculated into 50 ml of production media for itaconic acid production as described in Riscaldati et al. (*J Biotechnol* 2000, 83:219-230). The cultivation was performed at 30 $^{\circ}$ C. on a rotary shaker at 150 rpm. All experiments were done in three biological replicates. At day 3, samples were collected and RNA was extracted for RT-PCR. The average of results obtained from five independent RNA preparations is shown. All transcript levels were measured in triplicate for each RNA preparation. Error bars represent standard deviations from the means. Compared to wild type, expression level of mttA, cadA and mfsA were decreased 57, 37 and 23 fold in the Δ tfr strain.

FIGS. 6A-6C. Aconitic acid production in Δ cadA strain. The cultivation was performed at 30 $^{\circ}$ C. on a rotary shaker at 150 rpm. All experiments were done in three biological replicates. (A) at day 5, only Δ cadA produced cis-aconitic and trans-aconitic acid, while wild type and other mutants did not. (B) Time course of cis- and trans-aconitic acid production in Δ cad strain over 10 days. (C) Comparison of total aconitic acid production between wild type and Δ cadA mutant strains.

FIG. 7. Arrangement of transgene expression cassette for 3-HP Production in *A. pseudoterreus* with a synthetic beta-alanine pathway. A description of each Fragment is described in Example 8. The relevant fragments were cloned into pBlueScript SK(-) vector linearized with restriction enzyme H3/PstI. The whole expression cassette was linearized with restriction enzyme XhoI for the protoplast transformation for homologous recombination at cadA locus.

FIG. 8. Southern blot confirmation of cadA gene interruption by 3HP transgene expression cassette (FIG. 7). The cadA gene in the transgenic strains #2 (3HP-2) and #6 (3HP-6) was disrupted by the homologous recombination, while the random integration occurred in the strains #4 (3HP-4) and #5 (3HP-5). No insertion was observed in strains #1 and #3.

FIGS. 9A-9B. 3-HP production. *A. pseudoterreus* having a genetically inactivated cadA locus alone (cad1A), or additionally expressing panD, BAPAT, and HPDH (3HP-2, 3HP-4, 3HP-5, and 3-HP6), were grown at 30 $^{\circ}$ C. on a rotary shaker at 200 rpm for (A) 7 days, or (B) over 8 days, in the Riscaldati media with 20 \times TE, and 3-HP present in the supernatant measured using HPLC.

SEQUENCE LISTING

The nucleic acid sequences listed in the accompanying sequence listing are shown using standard abbreviations for

nucleotide bases and amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. The sequence listing submitted herewith, generated on Apr. 24, 2019, 80 kb, is herein incorporated by reference. In the accompanying sequence listing:

SEQ ID NOS: 1-8 are primers used to delete the *tf* gene in *A. pseudoterreus*.

SEQ ID NOS: 9-16 are primers used to delete the *mttA* gene in *A. pseudoterreus*.

SEQ ID NOS: 17-24 are primers used to delete the *cadA* gene in *A. pseudoterreus*.

SEQ ID NOS: 25-32 are primers used to delete the *mfsA* gene in *A. pseudoterreus*.

SEQ ID NOS: 33-40 are primers used to delete the *p450* gene in *A. pseudoterreus*.

SEQ ID NOS: 41-42 are primers used to amplify *mttA* in *A. pseudoterreus*.

SEQ ID NOS: 43-44 are primers used to amplify *cadA* in *A. pseudoterreus*.

SEQ ID NOS: 45-46 are primers used to amplify *mfsA* in *A. pseudoterreus*.

SEQ ID NOS: 47-48 are primers used to amplify *benA* in *A. pseudoterreus*.

SEQ ID NOS: 49 and 50 are exemplary *cadA* nucleic acid and protein sequences, respectively, from *A. terreus* (GenBank Accession Nos. AB326105.1 and BAG49047.1).

SEQ ID NOS: 51 and 52 are exemplary *cadA* nucleic acid and protein sequences, respectively, from *A. vadensis* CBS 113365 (GenBank Accession Nos. XM_025706777.1 and XP_025563141.1).

SEQ ID NOS: 53 and 54 are exemplary aspartate 1-decarboxylase (*panD*) nucleic acid and protein sequences, respectively, from *Tribolium castaneum* (GenBank Accession Nos. NM_001102585.1 and NP_001096055.1). Coding sequence nt 41-1663.

SEQ ID NOS: 55 and 56 are exemplary β -alanine-pyruvate aminotransferase (BAPAT) nucleic acid and protein sequences, respectively, from *Bacillus cereus* AH1272 (GenBank Accession Nos. ACMS01000158.1 (complement (10606 . . . 11961)) and EEL86940.1).

SEQ ID NOS: 57 and 58 are exemplary 3-hydroxypropionate dehydrogenase (HPDH) nucleic acid and protein sequences (GenBank Accession No. WP_000636571), respectively.

SEQ ID NO: 59 is an *A. pseudoterreus* 5'-*cadA* nucleic acid sequence.

SEQ ID NOS: 60-61 are primers used to isolate an *A. pseudoterreus* 5'-*cadA* gene.

SEQ ID NO: 62 is an *A. niger* *gpdA* promoter nucleic acid sequence.

SEQ ID NOS: 63-64 are primers used to isolate an *A. niger* *gpdA* promoter.

SEQ ID NO: 65 is *panD* cDNA of *Tribolium castaneum* with codon optimization for *A. pseudoterreus*.

SEQ ID NOS: 66-67 are primers used to isolate *panD* cDNA of *Tribolium castaneum* with codon optimization for *A. pseudoterreus*.

SEQ ID NO: 68 is a bidirectional terminator from *A. niger* *elf3*/multifunctional chaperone.

SEQ ID NOS: 69-70 are primers used to isolate bidirectional terminator from *A. niger* *elf3*/multifunctional chaperone.

SEQ ID NO: 71 is codon optimized synthetic cDNA of β -alanine-pyruvate aminotransferase (BAPAT) of *Bacillus cereus*.

SEQ ID NOS: 72-73 are primers used to isolate a codon optimized synthetic cDNA of BAPAT of *Bacillus cereus*.

SEQ ID NO: 74 is an *A. niger* *enol* promoter.

SEQ ID NOS: 75-76 are primers used to isolate an *A. niger* *enol* promoter.

SEQ ID NO: 77 is an *A. nidulans* *gpdA* promoter.

SEQ ID NOS: 78-79 are primers used to isolate an *A. nidulans* *gpdA* promoter.

SEQ ID NO: 80 is the codon optimized synthetic cDNA of *E. coli* 3-hydroxypropionate dehydrogenase (HPDH).

SEQ ID NOS: 81-82 are primers used to isolate a codon optimized synthetic cDNA of *E. coli* HPDH.

SEQ ID NO: 83 is a *trpC* terminator of *A. nidulans*.

SEQ ID NOS: 84-85 are primers used to isolate the *trpC* terminator of *A. nidulans*.

SEQ ID NO: 86 is a *trpC* terminator of *A. nidulans*.

SEQ ID NOS: 87-88 are primers used to isolate a *trpC* terminator of *A. nidulans*.

SEQ ID NO: 89 is an *A. oryzae* *ptrA* selection marker gene.

SEQ ID NOS: 90-91 are primers used to isolate the *A. oryzae* *ptrA* selection marker gene.

SEQ ID NO: 92 is an *A. pseudoterreus* 3'-*cadA* gene.

SEQ ID NOS: 93-94 are primers used to isolate an *A. pseudoterreus* 3'-*cadA* gene fragment.

SEQ ID NO: 95 is a combination of Fragments 7 to 9 (SEQ ID NOS: 77, 80, and 83, respectively).

SEQ ID NO: 96 is a primer used to isolate Fragments 7 to 9 (in combination with SEQ ID NO: 88).

SEQ ID NO: 97 is a combination of Fragments 11 and 12 (SEQ ID NOS: 89 and 92, respectively).

SEQ ID NO: 98 is a primer used to isolate Fragments 11 to 12 (in combination with SEQ ID NO: 90).

DETAILED DESCRIPTION

Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew et al. (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).


The singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. Similarly, the word "or" is intended to include "and" unless the context clearly indicates otherwise. Hence "comprising A or B" means including A, or B, or A and B. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. All publications, references and Genbank® Accession numbers (the sequence available on Apr. 24, 2019) mentioned herein are incorporated by reference in their entirety. The materials, methods, and examples are illustrative only and not intended to be limiting.

In order to facilitate review of the various embodiments of the disclosure, the following explanations of specific terms are provided:

3-hydroxypropionate dehydrogenase (HPDH): EC 1.1.1.59 An enzyme that catalyzes the chemical reaction: 3-hydroxypropionate+NAD⁺↔3-oxopropionate+NADH+H⁺. The term HPDH includes any HPDH gene (such as a bacterial or fungal panD sequence), cDNA, mRNA, or protein, that is a HPDH that can convert 3-hydroxypropionate and NAD into 3-oxopropionate, NADH, and H⁺ and vice versa. Expression or increased expression of HPDH, for example in an *Aspergillus* also expressing BAPAT and panD and having a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

HPDH sequences are publicly available. For example, SEQ ID NO: 57 discloses an HPDH coding sequence and GenBank® Accession No: WP_000636571 discloses an HPDH protein sequence (SEQ ID NO: 58); GenBank® Accession Nos. FR729477.2 (nt 1005136 . . . 1005885) and CBY27203.1 disclose exemplary *Yersinia enterocolitica* subsp. *palaearctica* Y11 HPDH nucleic acid and protein sequences, respectively; and GenBank® Accession Nos: CP004083.1 (complement (1399227 . . . 1399973) and AJQ99264.1 disclose exemplary *Enterobacteriaceae bacterium* bta3-1 HPDH nucleic acid and protein sequences, respectively. However, one skilled in the art will appreciate that in some examples, a HPDH sequence can include variant sequences (such as allelic variants and homologs) that retain HPDH activity and when expressed in an *Aspergillus* also expressing BAPAT and panD and with a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

Aconitic acid: An organic acid with two isomers, cis- and trans-acconitic acid. The Δ cadA fungi provided herein can be used to produce cis- and trans-acconitic acid.



Aspartate 1-decarboxylase (panD): EC 4.1.1.11. An enzyme that catalyzes the chemical reaction: L-aspartate↔beta-alanine+CO₂. The term panD includes any panD gene (such as a bacterial or fungal panD sequence), cDNA, mRNA, or protein, that is a panD that can convert L-aspartate into beta-alanine+CO₂ and vice versa. Expression or increased expression of panD, for example in an *Aspergillus* also expressing BAPAT and HPDH and having a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60%

at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions)

panD sequences are publicly available. For example, GenBank® Accession Nos: NM_001102585.1 and NP_001096055.1 disclose *Tribolium castaneum* panD nucleic acid and protein sequences, respectively (SEQ ID NOS: 55 and 56); GenBank® Accession Nos. CP002745.1 (complement (4249351 . . . 4249824)) and AEK63458.1 disclose exemplary *Collimonas fungivorans* Ter331 panD nucleic acid and protein sequences, respectively; and GenBank® Accession Nos: CP029034.1 (nt 1201611 . . . 1201994) and AWE15802.1 disclose exemplary *Bacillus velezensis* panD nucleic acid and protein sequences, respectively. However, one skilled in the art will appreciate that in some examples, a panD sequence can include variant sequences (such as allelic variants and homologs) that retain panD activity and when expressed in an *Aspergillus* also expressing BAPAT and HPDH and with a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

β-alanine-pyruvate aminotransferase (BAPAT): EC 2.6.1.18. An enzyme that can catalyze the reaction L-alanine+3-oxopropionate↔beta-alanine+pyruvate. The term BAPAT includes any BAPAT gene (such as a bacterial or fungal panD sequence), cDNA, mRNA, or protein, that is a BAPAT that can convert beta-alanine and pyruvate to L-alanine and 3-oxopropionate [or malonic semialdehyde], and vice versa. Expression or increased expression of BAPAT, for example in an *Aspergillus* also expressing HPDH and panD and having a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

BAPAT sequences are publicly available. For example, GenBank® Accession Nos: ACMS01000158.1 (complement (10606 . . . 11961)) and EEL86940.1 disclose *Bacillus cereus* AH1272 BAPAT nucleic acid and protein sequences, respectively (SEQ ID NOS: 55 and 56); GenBank® Accession Nos. DF820429.1 (complement (241627 . . . 242967)) and GAK28710.1 disclose exemplary *Serratia liquefaciens* FK01 BAPAT nucleic acid and protein sequences, respectively; and GenBank Accession Nos: LGUJ01000001.1 complement (92812 . . . 94140) and KOY12524.1 disclose exemplary *Bradyrhizobium diazoefficiens* BAPAT nucleic acid and protein sequences, respectively. However, one skilled in the art will appreciate that in some examples, a BAPAT sequence can include variant sequences (such as allelic variants and homologs) that retain BAPAT activity and when expressed in an *Aspergillus* also expressing HPDH and panD and with a genetically inactivated cadA gene (Δ cadA), results in a fungus that has an ability to produce more 3-HP than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60% at least 70%, at least 100%, at least 200%, at least 300%, or at least 400% more than a parent strain under the same growing conditions).

cadA (cis-acconitic acid decarboxylase): The cadA gene encodes an enzyme (EC 4.1.1.6) that catalyzes the chemical reaction cis-acconitate↔itaconate+CO₂. The term cadA (or

cadA) includes any cadA gene (such as a fungal cadA sequence), cDNA, mRNA, or protein, that is a cadA that can catalyze the decarboxylation of cis-aconitate to itaconate and CO₂ and vice versa, and when genetically inactivated results in a fungus that produces more aconitic acid than the parent strain without a genetically inactivated cadA gene (such as at least 20%, at least 30%, at least 50%, at least 60%, at least 75%, at least 100%, at least 200%, at least 500%, or 1000% more than a parent strain under the same growing conditions, for example at day 5 of production). In some examples, a parental strain containing a functional native cadA sequence does not produce detectable aconitic acid. In some examples, genetic inactivation of cadA results in a fungus that produces more trans-aconitic acid than cis-aconitic acid at day 10 of production, (such as at least 2-fold, at least 3-fold, at least 4-fold, or at least 5-fold more at day 10 of production).

cadA sequences are publicly available for many species of *Aspergillus*. For example, GenBank® Accession Nos: AB326105.1 and BAG49047.1 disclose *Aspergillus terreus* cadA nucleic acid and protein sequences, respectively (SEQ ID NOS: 49 and 50); GenBank® Accession Nos: XM_025706777.1 and XP_025563141.1 disclose *Aspergillus vadensis* CBS 113365 cadA nucleic acid and protein sequences, respectively (SEQ ID NOS: 51 and 52); and GenBank® Accession Nos: XM_025663103.1 and XP_025520527.1 disclose *Aspergillus piperis* CBS 112811 cadA nucleic acid and protein sequences, respectively. However, one skilled in the art will appreciate that in some examples, a cadA sequence can include variant sequences (such as allelic variants and homologs) that retain cadA activity but when genetically inactivated in *Aspergillus* results in a fungus that has an ability to produce more aconitic acid than the parent strain without a genetically inactivated cadA gene (such as at least 20%, at least 30%, at least 50%, at least 60%, at least 75%, at least 100%, at least 200%, at least 500%, or 1000% more than a parent strain under the same growing conditions, for example at day 5 of production).

Detectable: Capable of having an existence or presence ascertained. For example, production of aconitic acid or 3-HP is detectable if the signal generated is strong enough to be measurable.

Exogenous: The term “exogenous” as used herein with reference to nucleic acid and a particular cell refers to any nucleic acid that does not originate from that particular cell as found in nature. Thus, a non-naturally-occurring nucleic acid is considered to be exogenous to a cell once introduced into the cell. A nucleic acid that is naturally-occurring also can be exogenous to a particular cell. For example, an entire chromosome isolated from cell X is an exogenous nucleic acid with respect to cell Y once that chromosome is introduced into cell Y, even if X and Y are the same cell type.

In some examples, the panD, BAPAT, and HPDH nucleic acid or protein expressed in an *Aspergillus terreus* or *Aspergillus pseudoterreus* fungi does not naturally occur in the *Aspergillus terreus* or *Aspergillus pseudoterreus* fungi and is therefore exogenous to that fungi. For example, the panD, BAPAT, and HPDH nucleic acid molecule introduced into an *Aspergillus terreus* or *Aspergillus pseudoterreus* fungi can be from another organism, such as a bacterial panD, BAPAT, and HPDH sequence.

Genetic enhancement or up-regulation: When used in reference to the expression of a nucleic acid molecule, such as a gene, refers to any process which results in an increase in production of a gene product. A gene product can be RNA (such as mRNA, rRNA, tRNA, and structural RNA) or

protein. Examples of processes that increase transcription include those that facilitate formation of a transcription initiation complex, those that increase transcription initiation rate, those that increase transcription elongation rate, those that increase processivity of transcription and those that relieve transcriptional repression (for example by blocking the binding of a transcriptional repressor). Gene up-regulation can include inhibition of repression as well as expression above an existing level. Examples of processes that increase translation include those that increase translational initiation, those that increase translational elongation and those that increase mRNA stability. In one example, additional copies of genes are introduced into a cell in order to increase expression of that gene in the resulting transgenic cell.

Gene up-regulation includes any detectable increase in the production of a gene product. In certain examples, production of a gene product increases by at least 1.5-fold, at least 2-fold, or at least 5-fold), such as aspartate decarboxylase (panD), β-alanine-pyruvate aminotransferase (BAPAT), and 3-hydroxypropionate dehydrogenase (HPDH). For example, expression of a panD, BAPAT, and HPDH genes in *Aspergillus* (e.g., *A. terreus*) results in an *Aspergillus* strain having increased levels of the panD, BAPAT, and HPDH proteins, respectively, relative to the parent strain, which can permit the recombinant fungus to produce 3-HP. Genetic enhancement is also referred to herein as “enhancing or increasing expression.”

Genetic inactivation or down-regulation: When used in reference to the expression of a nucleic acid molecule, such as a gene, refers to any process which results in a decrease in production of a gene product. A gene product can be RNA (such as mRNA, rRNA, tRNA, and structural RNA) or protein. Therefore, gene down-regulation or deactivation includes processes that decrease transcription of a gene or translation of mRNA.

For example, a mutation, such as a substitution, partial or complete deletion, insertion, or other variation, can be made to a gene sequence that significantly reduces (and in some cases eliminates) production of the gene product or renders the gene product substantially or completely non-functional. For example, a genetic inactivation of the cadA gene in *Aspergillus* (e.g., *A. pseudoterreus*) results in *Aspergillus* having a non-functional or non-existent cadA protein, which results in the recombinant fungus to produce more aconitic acid. Genetic inactivation is also referred to herein as “functional deletion”.

Isolated: To be significantly separated from other agents. An “isolated” biological component (such as a nucleic acid molecule or protein) has been substantially separated, produced apart from, or purified away from other biological components in the cell of the organism in which the component occurs, for example, other chromosomal and extra-chromosomal DNA and RNA, and proteins. Nucleic acid molecules and proteins which have been “isolated” include nucleic acid molecules and proteins purified by standard purification methods. The term also embraces nucleic acid molecules and proteins prepared by recombinant expression in a host cell as well as chemically synthesized proteins and nucleic acids. Samples of isolated biological components include samples of the biological component wherein the biological component represents greater than 90% (for example, greater than 95%, such as greater than 98%) of the sample.

An “isolated” microorganism (such as a ΔcadA strain of *Aspergillus*) has been substantially separated or purified away from microorganisms of different types, strains, or

species. Microorganisms can be isolated by a variety of techniques, including serial dilution and culturing and resistance to certain chemicals, such as antibiotics. In some examples, an isolated Δ cadA strain of *Aspergillus* is at least 90% (for example, at least 95%, as at least 98%, at least 99%, or at least 99.99%) pure.

Mutation: A change in a nucleic acid sequence (such as a gene sequence) or amino acid sequence, for example as compared to a nucleic acid or amino acid sequence present in a wild-type or native organism. In particular examples, a mutation is introduced into a cadA gene in *Aspergillus*. Mutations can occur spontaneously, or can be introduced, for example using molecular biology methods (e.g., thereby generating a recombinant or transformed cell or microorganism). In particular examples, a mutation includes one or more nucleotide substitutions, deletions, insertions, or combinations thereof. In particular examples, the presence of one or more mutations in a gene can significantly inactivate and reduce expression of that gene.

Promoter: An array of nucleic acid control sequences which direct transcription of a nucleic acid. A promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements which can be located as much as several thousand base pairs from the start site of transcription. In some examples, a promoter is bi-directional. Native and non-native promoters can be used to drive expression of a gene, such as panD, BAPAT, and HPDH. Exemplary promoters that can be used include but are not limited to: enol promoter from *A. niger*, and dth1 from *A. nidulans* or *A. niger*.

Examples of promoters include, but are not limited to the SV40 promoter, the CMV enhancer-promoter, and the CMV enhancer/ β -actin promoter. Both constitutive and inducible promoters can be used in the methods provided herein (see e.g., Bitter et al., *Methods in Enzymology* 153:516-544, 1987). Also included are those promoter elements which are sufficient to render promoter-dependent gene expression controllable for cell-type specific, tissue-specific, or inducible by external signals or agents; such elements may be located in the 5' or 3' regions of the gene. Promoters produced by recombinant DNA or synthetic techniques can also be used to provide for transcription of the nucleic acid sequences.

Recombinant: A recombinant nucleic acid molecule or protein is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. In particular examples, this artificial combination is accomplished by chemical synthesis or by the artificial manipulation of isolated segments of nucleic acids, for example, by genetic engineering techniques such as those described in Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 3d ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N Y, 2001. The term recombinant includes nucleic acid molecules that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid molecule. A recombinant or transformed organism or cell, such as a recombinant *Aspergillus*, is one that includes at least one exogenous nucleic acid molecule, such as one used to genetically inactivate an endogenous cadA gene, and one used to express a non-native protein, such as exogenous panD, BAPAT, and HPDH nucleic acid coding sequences.

Sequence identity/similarity: The identity/similarity between two or more nucleic acid sequences, or two or more

amino acid sequences, is expressed in terms of the identity or similarity between the sequences. Sequence identity can be measured in terms of percentage identity; the higher the percentage, the more identical the sequences are. Sequence similarity can be measured in terms of percentage similarity (which takes into account conservative amino acid substitutions); the higher the percentage, the more similar the sequences are.

Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith & Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman & Wunsch, *J. Mol. Biol.* 48:443, 1970; Pearson & Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444, 1988; Higgins & Sharp, *Gene*, 73:237-44, 1988; Higgins & Sharp, *CABIOS* 5:151-3, 1989; Corpet et al., *Nuc. Acids Res.* 16:10881-90, 1988; Huang et al. *Computer Appls. in the Biosciences* 8, 155-65, 1992; and Pearson et al., *Meth. Mol. Bio.* 24:307-31, 1994. Altschul et al., *J. Mol. Biol.* 215:403-10, 1990, presents a detailed consideration of sequence alignment methods and homology calculations.

The NCBI Basic Local Alignment Search Tool (BLAST) (Altschul et al., *J. Mol. Biol.* 215:403-10, 1990) is available from several sources, including the National Center for Biological Information (NCBI, National Library of Medicine, Building 38A, Room 8N805, Bethesda, Md. 20894) and on the Internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx. Additional information can be found at the NCBI web site.

BLASTN is used to compare nucleic acid sequences, while BLASTP is used to compare amino acid sequences. To compare two nucleic acid sequences, the options can be set as follows: -i is set to a file containing the first nucleic acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second nucleic acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastn; -o is set to any desired file name (e.g., C:\output.txt); -q is set to -1; -r is set to 2; and all other options are left at their default setting. For example, the following command can be used to generate an output file containing a comparison between two sequences: C:\B12seq -i c:\seq1.txt -j c:\seq2.txt -p blastn -o c:\output.txt -q -1 -r 2.

To compare two amino acid sequences, the options of B12seq can be set as follows: -i is set to a file containing the first amino acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second amino acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastp; -o is set to any desired file name (e.g., C:\output.txt); and all other options are left at their default setting. For example, the following command can be used to generate an output file containing a comparison between two amino acid sequences: C:\B12seq c:\seq1.txt -j c:\seq2.txt -p blastp -o c:\output.txt. If the two compared sequences share homology, then the designated output file will present those regions of homology as aligned sequences. If the two compared sequences do not share homology, then the designated output file will not present aligned sequences.

Once aligned, the number of matches is determined by counting the number of positions where an identical nucleotide or amino acid residue is presented in both sequences. The percent sequence identity is determined by dividing the number of matches either by the length of the sequence set forth in the identified sequence, or by an articulated length (e.g., 100 consecutive nucleotides or amino acid residues from a sequence set forth in an identified sequence), followed by multiplying the resulting value by 100. For example, a nucleic acid sequence that has 1166 matches

when aligned with a test sequence having 1554 nucleotides is 75.0 percent identical to the test sequence (i.e., $1166 \div 1554 * 100 = 75.0$). The percent sequence identity value is rounded to the nearest tenth. For example, 75.11, 75.12, 75.13, and 75.14 are rounded down to 75.1, while 75.15, 75.16, 75.17, 75.18, and 75.19 are rounded up to 75.2. The length value will always be an integer. In another example, a target sequence containing a 20-nucleotide region that aligns with 20 consecutive nucleotides from an identified sequence as follows contains a region that shares 75 percent sequence identity to that identified sequence (i.e., $15 \div 20 * 100 = 75$).

For comparisons of amino acid sequences of greater than about 30 amino acids, the Blast 2 sequences function is employed using the default BLOSUM62 matrix set to default parameters, (gap existence cost of 11, and a per residue gap cost of 1). Homologs are typically characterized by possession of at least 70% sequence identity counted over the full-length alignment with an amino acid sequence using the NCBI Basic Blast 2.0, gapped blastp with databases such as the nr or swissprot database. Queries searched with the blastn program are filtered with DUST (Hancock and Armstrong, 1994, *Comput. Appl. Biosci.* 10:67-70). Other programs use SEG. In addition, a manual alignment can be performed. Proteins with even greater similarity will show increasing percentage identities when assessed by this method, such as at least 75%, 80%, 85%, 90%, 95%, or 99% sequence identity.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode identical or similar (conserved) amino acid sequences, due to the degeneracy of the genetic code. Changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid molecules that all encode substantially the same protein. Such homologous nucleic acid sequences can, for example, possess at least 60%, 70%, 80%, 90%, 95%, 98%, or 99% sequence identity determined by this method.

One of skill in the art will appreciate that these sequence identity ranges are provided for guidance only; it is possible that strongly significant homologs could be obtained that fall outside the ranges provided. Thus, a variant *cadA*, *panD*, *BAPAT*, or *HPDH* protein or nucleic acid molecule that can be used with the organisms and methods of the present disclosure can have at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to the SEQ ID NOs; and GenBank® Accession Nos. provided herein.

Transformed: A cell, such as a fungal cell, into which a nucleic acid molecule has been introduced, for example by molecular biology methods. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including, but not limited to chemical methods (e.g., calcium-phosphate transfection), physical methods (e.g., electroporation, microinjection, particle bombardment), fusion (e.g., liposomes), receptor-mediated endocytosis (e.g., DNA-protein complexes, viral envelope/capsid-DNA complexes) and by biological infection by viruses such as recombinant viruses. In one example, the protoplast transformation provide herein, such as in Example 1, is used.

Vector: A nucleic acid molecule as introduced into a host cell, thereby producing a transformed or recombinant host cell. A vector may include nucleic acid sequences that permit it to replicate in the host cell, such as an origin of replication. A vector may also include a *panD*, *BAPAT*, or *HPDH* coding sequence, or a sequence used to genetically inactivate *cadA*

for example in combination with a promoter, and/or selectable marker genes, and other genetic elements. A vector can transduce, transform or infect a cell, thereby causing the cell to express nucleic acids and/or proteins other than those native to the cell. A vector optionally includes materials to aid in achieving entry of the nucleic acid into the cell, such as a viral particle, liposome, protein coating or the like. In one example, a vector is a plasmid.

Overview

The filamentous fungus *Aspergillus pseudoterreus* has been used for industrial production of itaconic acid. *cis*-aconitic acid decarboxylase (*cadA*) is the key enzyme in itaconic acid production. The itaconic acid biosynthesis cluster is composed of genes *tf*, *mttA*, *cadA* and *mfsA*. As shown in FIG. 1, itaconic acid (IA) is produced from glucose. Glucose is utilized in the cell mainly by the glycolytic pathway and metabolized to pyruvate, which forms citric acid. *cis*-aconitic acid is derived from citric acid as a primary precursor of IA. *cis*-aconitic acid decarboxylase (*cadA*) removes carbon dioxide from *cis*-aconitic acid and forms itaconic acid. However, *cadA* is localized in the cytosol, while *cis*-aconitic acid is formed from the TCA cycle in the mitochondria. *mttA* is localized on the mitochondrial membrane and functioned to transport *cis*-aconitic acid from mitochondria to cytosol. Another transporter, *mfsA* is also an organic acid transporter that may be involved in exporting itaconic acid out of cells.

The first demonstration of genetically inactivating the *cadA* gene in *Aspergillus pseudoterreus* is shown herein. In the *cadA* deletion strain (Δ *cadA*), no more itaconic acid is produced. At the same time significant amount of *cis*-aconitic acid and *trans*-aconitic acid are detected. Blocking the itaconic acid production pathway permits the carbon to be diverted towards other organic acid production. The Δ *cadA* *Aspergillus* can be used as a host for chemical platform, and provides a new way to produce aconitic acid and other organic acids (for example by expressing other genes needed for production of those acids, such as *panD*, *BAPAT*, and *HPDH* for 3-HP production). This strain works as biocatalyst that converts biomass into aconitic acid through bioproduction method at room temperature (such as about 20-35° C.) and ordinary pressure (such as about 1 atm). Current processes of aconitic acid production include chemical synthesis that require high temperatures and harmful reagents.

The EST data provided herein demonstrated that four genes, *tf*, *cadA*, *mttA* and *mfsA* show high transcription frequency after IA production starts, but not before IA production begins. The high expression of these genes persists through the production process. Genes upstream and downstream of the cluster did not show expression differences before and after production. One gene downstream next to *mfsA*, a *p450* enzyme, also showed high expression after IA production started, however, deletion of this gene did not effect IA yield.

Correlations between the IA gene cluster and IA production were further investigated by constructing deletion strains. In a Δ *cad* strain, no IA was detected, while trace amounts of IA were detected in an *mttA* knockout. IA production in an *mfsA* deletion strain decreased one third compared with wild type. This indicates *mfsA* can transport IA across the cell membrane. In the Δ *tf* strain, IA production decreased eight fold and slowed the production rate compared to wild type. Also in the *tf* deletion strain, expression of *cadA*, *mttA* and *mfsA* significantly decreased. RT-PCR

results indicated that the expression level of genes in the IA cluster was regulated by *tf*, which is turned on by IA production conditions.

The Δ cadA strain produced aconitic acid. During the production, cis-aconitic acid was detected first, followed by the appearance of trans-aconitic acid. cis-aconitic acid levels remained consistent from day 5 forward. The trans-aconitic acid levels continued to increase from days 4 to 10. By day 10, more than 10 g/L trans-aconitic acid was detected in the supernatant. In the Δ cadA strain, cis-aconitic acid decarboxylase is not produced, and the cis-aconitic acid cannot be converted to itaconic acid by decarboxylation and accumulates in the cell. cis-aconitic acid was transported outside the cell. cis-aconitic acid is not stable in the acid solution and is rapidly converted into trans-aconitic acid.

Aconitic acid is an unsaturated tricarboxylic acid and is noted as a top 30 potential building block by United States Department of Energy (DOE). Trans-aconitic acid can be used to make polymers. Currently, trans-aconitic acid is produced by chemical synthesis and requires high temperature and harmful solvents. Generation of trans-aconitic acid has been achieved by metabolic engineering aconitase isomerase from *Pseudomonas* sp. WU-0701 into *E. coli*. However, the substrate for the recombinant *E. coli* to produce trans-aconitic acid is citric acid, which has to be generated first from fermentation. In contrast, the disclosed Δ cadA fungi can produce trans-aconitic acid directly from renewable biomass substrates. Also since the *cadA* is not functional and precursors from TAC cycle accumulate in the cell, the carbon can be rerouted to generate other organic acid since *A. pseudoterreus* is industrial filamentous fungi and tolerant to low pH.

Based on these observations, provided herein are isolated recombinant (i.e., transformed) *Aspergillus* fungi that include a genetic inactivation (also referred to as a functional deletion) of an endogenous cis-aconitic acid decarboxylase (*cadA*) gene. Such fungi are referred to herein as Δ cadA fungi. Exemplary *Aspergillus* species that can be used include *Aspergillus pseudoterreus* and *Aspergillus terreus*. In some examples, the endogenous *cadA* gene is genetically inactivated by mutation (such as a complete or partial deletion of the *cadA* gene) or by insertional mutation (such as by insertion of another nucleic acid molecule into the *cadA* gene, such as an antibiotic resistance marker).

In some examples, the *cadA* gene prior to its genetic inactivation encodes a protein having at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 50 or 52. In some examples, the *cadA* gene (or its coding sequence) prior to its genetic inactivation comprises at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 49, 51, 59 or 92.

The disclosed Δ cadA fungi can include other exogenous genes to express proteins needed to permit the fungi to produce other organic acids. For example, the disclosed Δ cadA fungi can further include an exogenous nucleic acid molecule encoding aspartate 1-decarboxylase (*panD*), an exogenous nucleic acid molecule encoding β -alanine-pyruvate aminotransferase (*BAPAT*), and an exogenous nucleic acid molecule encoding 3-hydroxypropionate dehydrogenase (*HPDH*). *panD*, *BAPAT*, and *HPDH* coding sequences can be part of a one or more nucleic acid molecules, such as a vector. In addition, expression of the *panD*, *BAPAT*, and *HPDH* coding sequences can be driven by one or more promoters, such as a bi-directional promoter. In some examples, the promoter is native to the gene it is expressing. In some examples, the promoter is from *A. niger*. In some

examples, the *panD*, *BAPAT*, and/or *HPDH* coding sequences are inserted into the *cadA* gene, genetically inactivating *cadA*. In some examples, the exogenous nucleic acid molecule encoding *panD* has at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 53 or 65, and/or encodes a *panD* protein comprising at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 54. In some examples, the exogenous nucleic acid molecule encoding *BAPAT* has at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 55, and/or encodes a *BAPAT* protein having at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 56. In some examples, the exogenous nucleic acid molecule encoding *HPDH* has at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 57, and/or encodes a *HPDH* protein having at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 58.

The disclosure also provides compositions that include the Δ cadA fungi, and the Δ cadA fungi expressing other genes (such as *panD*, *BAPAT*, and *HPDH*). Such a composition can include a solid or liquid culture or growth media, such as complete media, minimal media, or Riscaldati medium (such as modified Riscaldati medium with 20 \times trace elements).

The disclosure also provides kits that include the Δ cadA fungi, and the Δ cadA fungi expressing other genes (such as *panD*, *BAPAT*, and *HPDH*). Such a kits can include a solid or liquid culture or growth media, such as complete media, minimal media, or Riscaldati medium (such as modified Riscaldati medium with 20 \times trace elements).

Also provided are methods of using the disclosed Δ cadA fungi to make aconitic acid. Such a method can include culturing the recombinant *Aspergillus* Δ cadA fungi under conditions that permit the fungus to make aconitic acid, such as growth in Riscaldati medium, thereby making aconitic acid. In some examples the aconitic acid generated is cis-aconitic acid, trans-aconitic acid, or both. In some examples, the fungi are cultured at room temperature (e.g., 20-35 $^{\circ}$ C.) at normal atmospheric pressure (e.g., 1 atm). In some examples, the method includes purifying or isolating the aconitic acid, for example from the culture media or from the cultured fungus. In some examples, the aconitic acid is isolated at least 2 days, at least 3 days, at least 5 days, at least 8 days or at least 10 days after the start of culturing, such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 days after the start of culturing.

Also provided are methods of using the disclosed Δ cadA fungi expressing *panD*, *BAPAT*, and *HPDH* to make 3-HP. Such a method can include culturing the recombinant *Aspergillus* Δ cadA fungi expressing *panD*, *BAPAT*, and *HPDH* under conditions that permit the fungus to make 3-HP, such as growth in Riscaldati medium (such as modified Riscaldati medium with 20 \times trace elements), thereby making 3-HP. In some examples, the fungi are cultured at room temperature (e.g., 20-35 $^{\circ}$ C.) at normal atmospheric pressure (e.g., 1 atm). In some examples, the method includes purifying or isolating the 3-HP, for example from the culture media or from the cultured fungus. In some examples, the 3-HP is isolated at least 2 days, at least 3 days, at least 5 days, at least 8 days or at least 10 days after the start of culturing, such as

2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 days after the start of culturing.

Recombinant Δ cadA Fungi

The present disclosure provides isolated recombinant *Aspergillus* fungi having its endogenous cadA gene genetically inactivated (e.g., functional deletion) of. Such fungi are referred to herein as Δ cadA fungal strains. It is shown herein that Δ cadA *Aspergillus* strains have increased aconitic acid production as compared to *Aspergillus* having native levels of cadA expression.

Any variety or strain of *Aspergillus* can be used. In particular examples, the *Aspergillus* fungus is *A. terreus* or *A. pseudoterreus*, as well as particular strains thereof (for example *A. terreus* NRRL 1960, *A. pseudoterreus* ATCC 32359).

In addition, any method for genetic inactivation can be used, as long as the expression of the cadA gene is significantly reduced or eliminated, or the function of the cadA protein is significantly reduced or eliminated. In particular examples, the cadA gene is genetically inactivated by complete or partial deletion mutation or by insertional mutation. In some examples genetic inactivation need not be 100%. In some embodiments, genetic inactivation refers to at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95% gene or protein inactivation. The term "reduced" or "decreased" as used herein with respect to a cell and a particular gene or protein activity refers to a lower level of activity than that measured in a comparable cell of the same species. For example, a particular *A. terreus* or *A. pseudoterreus* lacking cadA activity has reduced cadA activity if a comparable *A. terreus* or *A. pseudoterreus* not having an cadA genetic inactivation has detectable cadA activity.

cadA sequences are disclosed herein and others are publicly available, for example from GenBank or EMBL. In some examples, the cadA gene functionally deleted encoded a protein having at least 80%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 50 or 52 prior to its genetic inactivation. In some examples, the endogenous cadA gene functionally deleted comprises at least 80%, at least 90%, at least 95%, at least 97%, or at least 98% sequence identity to SEQ ID NO: 49, 51, 59, or 92 prior to its genetic inactivation.

The genetic inactivation of cadA results in many phenotypes in the recombinant Δ cadA *Aspergillus*, such as *A. terreus* or *A. pseudoterreus*. For example, Δ cadA mutants can have one or more of the following phenotypes: produces at least 2-fold, at least 3-fold, at least 3.5 fold, at least 5-fold, at least 8-fold, or at least 10-fold more total aconitic acid than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus* (for example at day 3, 4, 5, 6, 7, 8, 9 or 10 of production); produces at least 2-fold more cis-acetic acid at day 5, 6, 7, 8, 9, or 10 of culturing in Riscaldati medium than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus*; produces at least 2-fold, at least 3-fold, at least 5-fold, or at least 10-fold more trans-acetic acid at day 10 of culturing in Riscaldati medium than a wild-type *Aspergillus terreus* or *Aspergillus pseudoterreus*; or combinations thereof. In some examples, such increases are relative to *Aspergillus terreus* strain ATCC 32359 grown under the same conditions as the Δ cadA mutant. In some examples, an increased total aconitic acid production by Δ cadA fungi occurs at least 3 days (such as at least 4, 5, 6, 7, 8, 9, or 10 days) after inoculation in Riscaldati medium (such as at least 0.5 g/L aconitic acid or at least 1 g/L aconitic acid), as

compared to no detectable aconitic acid produced by *Aspergillus terreus* strain ATCC 32359 at the same time point.

Additional genes can also be inactivated in the Δ cadA fungi, wherein the additional genes may or may not provide additional enhancement of aconitic acid production to the fungus. In one example, the Δ cadA fungi includes overexpressed or upregulated aconitic acid transporters.

In some examples, Δ cadA fungi include one or more additional exogenous nucleic acid molecules, for example to permit production of other organic acids by the recombinant fungi. In one example, the Δ cadA fungi includes an exogenous nucleic acid molecule encoding aspartate decarboxylase (panD), an exogenous nucleic acid molecule encoding β -alanine-pyruvate aminotransferase (BAPAT), and an exogenous nucleic acid molecule encoding 3-hydroxypropionate dehydrogenase (HPDH). Such exogenous nucleic acid molecules can be part of one or more exogenous nucleic acid molecules (such as 1, 2 or 3 exogenous nucleic acid molecules). In some examples, exogenous nucleic acid molecules can be part of a vector, such as a plasmid or viral vector. In some examples, expression of the exogenous nucleic acid molecules is driven by one or more promoters, such as a constitutive or inducible promoter, or a bi-directional promoter. In some examples, the promoter used to drive expression of panD, BAPAT, and HPDH is a native promoter (e.g., native to the panD, BAPAT, and HPDH gene expressed). In other examples, the promoter used to drive expression of panD, BAPAT, and HPDH is a non-native promoter (e.g., exogenous to the panD, BAPAT, and HPDH gene expressed). In some examples, such a Δ cadA fungi expressing panD, BAPAT, and HPDH are used to produce 3-HP.

A. Methods of Functionally Deleting cadA

As used herein, an "inactivated" or "functionally deleted" cadA gene means that the cadA gene has been mutated, for example by insertion, deletion, or substitution (or combinations thereof) of one or more nucleotides such that the mutation substantially reduces (and in some cases abolishes) expression or biological activity of the encoded cadA gene product. The mutation can act through affecting transcription or translation of the cadA gene or its mRNA, or the mutation can affect the cadA polypeptide product itself in such a way as to render it substantially inactive.

In one example, a strain of *Aspergillus* is transformed with a vector which has the effect of down-regulating or otherwise inactivating a cadA gene. This can be done by mutating control elements such as promoters and the like which control gene expression, by mutating the coding region of the gene so that any protein expressed is substantially inactive, or by deleting the cadA gene entirely. For example, a cadA gene can be functionally deleted by complete or partial deletion mutation (for example by deleting a portion of the coding region of the gene) or by insertional mutation (for example by inserting a sequence of nucleotides into the coding region of the gene, such as a sequence of about 1-5000 nucleotides). In one example, the cadA gene is genetically inactivated by inserting coding sequences for panD, BAPAT, and/or HPDH. Thus, the disclosure provides transformed fungi that include at least one exogenous nucleic acid molecule which genetically inactivates a cadA gene. In one example, such a transformed cell produces more aconitic acid, for example relative to a comparable fungus with a native or wild-type cadA sequence.

In particular examples, an insertional mutation includes introduction of a sequence that is in multiples of three bases (e.g., a sequence of 3, 9, 12, or 15 nucleotides) to reduce the possibility that the insertion will be polar on downstream

genes. For example, insertion or deletion of even a single nucleotide that causes a frame shift in the open reading frame, which in turn can cause premature termination of the encoded *cadA* polypeptide or expression of a substantially inactive polypeptide. Mutations can also be generated through insertion of foreign gene sequences, for example the insertion of a gene encoding antibiotic resistance (such as hygromycin or bleomycin), or *panD*, *BAPAT*, and/or *HPDH* coding sequences.

In one example, genetic inactivation is achieved by deletion of a portion of the coding region of the *cadA* gene. For example, some, most (such as at least 50%) or virtually the entire coding region can be deleted. In particular examples, about 5% to about 100% of the gene is deleted, such as at least 20% of the gene, at least 40% of the gene, at least 75% of the gene, or at least 90% of the *cadA* gene.

Deletion mutants can be constructed using any of a number of techniques. In one example, homologous double crossover with fusion PCR products is employed to genetically inactivate one or more genes in *Aspergillus*. A specific example of such a method is described in Example 1 below.

In one example, a strategy using counterselectable markers can be employed which has been utilized to delete genes. For a review, see Reyrat et al. (*Infect. Immun.* 66:4011-4017, 1998). In this technique, a double selection strategy is employed wherein a plasmid is constructed encoding both a selectable and counterselectable marker, with flanking DNA sequences derived from both sides of the desired deletion. The selectable marker is used to select for fungi in which the plasmid has integrated into the genome in the appropriate location and manner. The counterselectable marker is used to select for the very small percentage of fungi that have spontaneously eliminated the integrated plasmid. A fraction of these fungi will then contain only the desired deletion with no other foreign DNA present.

In another technique, the *cre-lox* system is used for site specific recombination of DNA (for example see Steiger et al., *Appl. Environ. Microbiol.* 77(1):114, 2011). The system includes 34 base pair *lox* sequences that are recognized by the bacterial *cre* recombinase gene. If the *lox* sites are present in the DNA in an appropriate orientation, DNA flanked by the *lox* sites will be excised by the *cre* recombinase, resulting in the deletion of all sequences except for one remaining copy of the *lox* sequence. Using standard recombination techniques, the targeted gene of interest (e.g., *cadA*) can be deleted in the *Aspergillus* genome and to replace it with a selectable marker (for example a gene coding for kanamycin resistance) that is flanked by the *lox* sites. Transient expression (by electroporation of a suicide plasmid containing the *cre* gene under control of a promoter that functions in *Aspergillus*) of the *cre* recombinase should result in efficient elimination of the *lox* flanked marker. This process will produce a mutant containing the desired deletion mutation and one copy of the *lox* sequence.

In another method, a *cadA* gene sequence in the *Aspergillus* genome is replaced with a marker gene, such as green fluorescent protein, β -galactosidase, or luciferase. In this technique, DNA segments flanking a desired deletion are prepared by PCR and cloned into a suicide (non-replicating) vector for *Aspergillus*. An expression cassette, containing a promoter active in *Aspergillus* and the appropriate marker gene, is cloned between the flanking sequences. The plasmid is introduced into wild-type *Aspergillus*. Fungi that incorporate and express the marker gene are isolated and examined for the appropriate recombination event (replacement of the wild type *cadA* gene with the marker gene).

Thus, for example, a fungal cell can be engineered to have a disrupted *cadA* gene using common mutagenesis or knock-out technology. (Methods in Yeast Genetics (1997 edition), Adams, Gottschling, Kaiser, and Sterns, Cold Spring Harbor Press, 1998; Datsenko and Wanner, *Proc. Natl. Acad. Sci. USA* 97: 6640-5, 2000; and Dai et al., *Appl. Environ. Microbiol.* 70(4):2474-85, 2004). Alternatively, antisense technology can be used to reduce or eliminate the activity of *cadA*. For example, a fungal cell can be engineered to contain a cDNA that encodes an antisense molecule that prevents *cadA* from being translated. The term "antisense molecule" encompasses any nucleic acid molecule or nucleic acid analog (e.g., peptide nucleic acids) that contains a sequence that corresponds to the coding strand of an endogenous *cadA* gene. An antisense molecule also can have flanking sequences (e.g., regulatory sequences). Thus, antisense molecules can be ribozymes or antisense oligonucleotides. A ribozyme can have any general structure including, without limitation, hairpin, hammerhead, or axehead structures, provided the molecule cleaves RNA. Further, gene silencing can be used to reduce the activity of *cadA*.

In one example, to genetically inactivate *cadA* in *A. pseudoterreus* or *A. terreus*, protoplast transformation is used, for example as described in Example 1. For example, conidia of *A. pseudoterreus* or *A. terreus* are grown in liquid complete medium at room temperature (e.g., about 20-35° C., such as 30° C.) and grown for at least 12 hours (such as at least 16 hours, or at least 18 hours, such as 12-24 hours, or 16-18 hours), at least 100 rpm, such as at least 150 rpm, for example 100 to 200 rpm. The resulting mycelia are subsequently harvested, for example by filtration. Protoplasts are prepared, for example by treating the harvested mycelia with a lysing enzyme (for example in an osmotic wash buffer for at least 30 min, at least 60 min, at least 120 min, or at leave 240 min, such as 2 h). The resulting protoplasts are collected (e.g., by filtering). Protoplasts can be washed, for example with a Washing Solution (0.6M KCl, 0.1M Tris/HCl, pH 7.0) and Conditioning Solution (0.6M KCl, 50 mM CaCl₂, 10 mM Tris/HCl, pH 7.5). The protoplasts are transformed, for example in the conditioning solution. In some examples, at least 0.5 ug, at least 1 ug, or at least 2 ug of DNA (such as 1-2 ug DNA) is added to at least 10⁶ protoplasts (such as at least 10⁷ or 2x10⁷ protoplasts). Polyethylene glycol (PEG), such as PEG8000 is added (such as 25% PEG8000, 0.6M KCl, 50 mM CaCl₂, 10 mM Tris/HCl, and pH 7.5) and the reaction incubated for at least 5 min (such as at least 10 min, at least 20 min, or at least 30 min, such as 10-30 min, 15-20 min, or 20 min) on ice. Additional PEG solution can be added and the reaction incubated for at least 1 min, at least 3 min, or at least 5 min, on ice. Conditioning Solution is added to the reaction, and the protoplast suspension mixed with warm selection agar (Minimal media+0.6M KCl+1.5% Agar+100 ug/ml hygromycin) (such as at 50° C.), and poured directly onto petri dish plates and allowed to solidify. Solidified plates can be inverted and incubated overnight at room temperature (e.g., about 20-35° C., such as 30° C.). The following day, the plates can be overlaid with Minimal Medium containing a selection antibiotic, such as hygromycin. Colonies appear after 3-4 days. Transformants can be excised and transferred to MM plate containing the selection antibiotic.

B. Measuring Gene Inactivation

A fungus having an inactivated *cadA* gene can be identified using known methods. For example, PCR and nucleic acid hybridization techniques, such as Northern and Southern analysis, can be used to confirm that a fungus has a genetically inactivated *cadA* gene. In one example, real-time

reverse transcription PCR (qRT-PCR) is used for detection and quantification of targeted messenger RNA, such as mRNA of *cadA* gene in the parent and mutant strains as grown at the same culture conditions. Immunohisto-chemical and biochemical techniques can also be used to determine if a cell expresses *cadA* by detecting the expression of the *cadA* peptide encoded by *cadA*. For example, an antibody having specificity for *cadA* can be used to determine whether or not a particular fungus contains a functional nucleic acid encoding *cadA* protein. Further, biochemical techniques can be used to determine if a cell contains a *cadA* gene inactivation by detecting a product produced as a result of the lack of expression of the peptide. For example, production of aconitic acid by *A. terreus* or *A. pseudoterreus* can indicate that such a fungus contains an inactivated *cadA* gene.

C. Measuring Aconitic Acid Production

Methods of determining whether a genetic inactivation of *cadA* in *Aspergillus*, such as *A. terreus* or *A. pseudoterreus*, increases aconitic acid production, for example relative to the same strain of *A. terreus* or *A. pseudoterreus* with a native *cadA* sequence (such as a parental strain), are provided herein. Although particular examples are disclosed herein, the methods are not limiting.

For example, production of aconitic acid by *Aspergillus* (such as a Δ *cadA* strain) can be measured using a spectrophotometric assay, by liquid chromatography (LC), or high-pressure liquid chromatography (HPLC) methods. In some examples, the supernatant of the fungus is analyzed for the presence of aconitic acid. In some examples, the culture media containing the Δ *cadA* strain is filtered prior to measuring aconitic acid in the culture media (supernatant).

D. *cadA* Sequences

cadA protein and nucleic acid sequences are publicly available and specific examples are provided herein. In addition, *cadA* sequences can be identified using molecular biology methods.

Examples of *cadA* nucleic acid sequences are shown in SEQ ID NOS: 49, 51, 59 and 92. However, the disclosure also encompasses variants of SEQ ID NOS: 49, 51, 59 and 92 which encode a functional *cadA* protein. One skilled in the art will understand variants of the *cadA* nucleic acid sequences provided herein can be genetically inactivated. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). In addition, the degeneracy of the code permits multiple nucleic acid sequences to encode the same protein. Such variant *cadA* nucleic acid molecules can share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to any *cadA* nucleic acid sequence, such as SEQ ID NO: 49, 51, 59 or 92.

Examples of *cadA* protein sequences are shown in SEQ ID NOS: 50 and 52. However, the disclosure also encompasses variants SEQ ID NOS: 50 and 52 which retain *cadA* activity. One skilled in the art will understand that variants of these *cadA* enzyme sequences can be inactivated. Variant sequences can be identified, for example by aligning known *cadA* sequences. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). Such *cadA* peptides share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to a *cadA* protein sequence, such as SEQ ID NO: 50 or 52.

In some examples, a *cadA* sequence that is to be genetically inactivated encodes or includes one or more conservative amino acid substitutions. A conservative amino acid substitution is a substitution of one amino acid (such as one found in a native sequence) for another amino acid having similar biochemical properties. Typically, conservative substitutions have little to no impact on the activity of a resulting peptide. In one example, a *cadA* sequence (such as SEQ ID NO: 50 or 52) includes one or more amino acid substitutions, such as conservative substitutions (for example at 1, 2, 5 or 10 residues). Examples of amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative substitutions include: Ser for Ala; Lys for Arg; Gln or His for Asn; Glu for Asp; Ser for Cys; Asn for Gln; Asp for Glu; Pro for Gly; Asn or Gln for His; Leu or Val for Ile; Ile or Val for Leu; Arg or Gln for Lys; Leu or Ile for Met; Met, Leu or Tyr for Phe; Thr for Ser; Ser for Thr; Tyr for Trp; Trp or Phe for Tyr; and Ile or Leu for Val. Further information about conservative substitutions can be found in, among other locations in, Ben-Bassat et al., (*J. Bacteriol.* 169:751-7, 1987), O'Regan et al., (*Gene* 77:237-51, 1989), Sahin-Toth et al., (*Protein Sci.* 3:240-7, 1994), Hochuli et al., (*Bio/Technology* 6:1321-5, 1988), WO 00/67796 (Curd et al.) and in standard textbooks of genetics and molecular biology.

The *cadA* gene inactivated in a fungus, in particular examples, includes a sequence that encodes a *cadA* protein having at least at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a *cadA* protein sequence, such as SEQ ID NO: 50 or 52, wherein the protein can catalyze the decarboxylation of *cis*-aconitate to itaconate and CO₂ and vice versa. In a specific example, the *cadA* gene inactivated in a fungus encodes a *cadA* protein shown in SEQ NO: 50 or 52.

The *cadA* gene that is to be inactivated in a fungus, in particular examples, includes a sequence (such as a coding sequence) having at least at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a *cadA* nucleic acid sequence, such as SEQ ID NO: 49, 51, 59, or 92, and encodes a *cadA* protein that can catalyze the decarboxylation of *cis*-aconitate to itaconate and CO₂ and vice versa. In a specific example, *cadA* gene inactivated in a fungus is the sequence of SEQ ID NO: 2 or 4.

One skilled in the art will appreciate that additional *cadA* sequences can be identified. For example, *cadA* nucleic acid molecules that encode a *cadA* protein can be identified and obtained using molecular cloning or chemical nucleic acid synthesis procedures and techniques, including PCR. In addition, nucleic acid sequencing techniques and software programs that translate nucleic acid sequences into amino acid sequences based on the genetic code can be used to determine whether or not a particular nucleic acid has any sequence homology with known *cadA* sequences. Sequence alignment software such as MEGALIGN (DNASTAR, Madison, Wis., 1997) can be used to compare various sequences.

In addition, nucleic acid hybridization techniques can be used to identify and obtain a nucleic acid molecule that encodes a *cadA* protein. Briefly, any known *cadA* nucleic acid molecule, or fragment thereof, can be used as a probe to identify similar nucleic acid molecules by hybridization under conditions of moderate to high stringency. Such similar nucleic acid molecules then can be isolated, sequenced, and analyzed to determine whether the encoded protein is a *cadA* protein.

E. panD, BAPAT, and HPDH Sequences

panD, BAPAT, and HPDH protein and nucleic acid sequences are publicly available and specific examples are provided herein. In addition, panD, BAPAT, and HPDH sequences can be identified using molecular biology methods.

Exemplary of panD coding sequences are shown in SEQ ID NO: 53 and 65. However, the disclosure also encompasses variants of SEQ ID NO: 53 and 65 which encode a functional panD protein. Exemplary of BAPAT coding sequences are shown in SEQ ID NO: 55 and 71. However, the disclosure also encompasses variants of SEQ ID NO: 55 and 71 which encode a functional BAPAT protein. Exemplary of HPDH coding sequences are shown in SEQ ID NO: 57 and 80. However, the disclosure also encompasses variants of SEQ ID NO: 57 and 80 which encode a functional HPDH protein.

One skilled in the art will understand variants of the panD, BAPAT, and HPDH nucleic acid sequences provided herein can be introduced into an *Aspergillus* fungus, such as one that is Δ cadA, such as inserting panD, BAPAT, and HPDH expression sequences into the native cadA gene to inactivate it. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). In addition, the degeneracy of the code permits multiple nucleic acid sequences to encode the same protein. In some examples, a panD, BAPAT, and HPDH sequence that is to be expressed in an *Aspergillus* fungus is codon optimized for expression in *Aspergillus*, such as *Aspergillus terreus* or *pseudoterreus*. Such variant panD, BAPAT, and HPDH nucleic acid molecules in some examples share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to any panD, BAPAT, and HPDH nucleic acid sequence, such as SEQ ID NO: 53, 55, or 57, respectively, or SEQ ID NO: 65, 71, or 80, respectively.

Exemplary panD, BAPAT, and HPDH protein sequences are shown in SEQ ID NOS: 54, 56, and 58, respectively. However, the disclosure also encompasses variants SEQ ID NOS: 54, 56, and 58 which retain panD, BAPAT, and HPDH activity, respectively. One skilled in the art will understand that variants of these panD, BAPAT, and HPDH sequences can be expressed in an *Aspergillus* fungus, such as one that is Δ cadA. Variant sequences can be identified, for example by aligning known panD, BAPAT, and HPDH sequences. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). Such panD, BAPAT, and HPDH peptides expressed in Δ cadA fungus in some examples share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to a panD, BAPAT, and HPDH protein sequence, such as SEQ ID NO: 54, 56, or 58, respectively.

In some examples, a panD, BAPAT, and HPDH sequence that is to be expressed in an *Aspergillus* fungus encodes or includes one or more conservative amino acid substitutions. In one example, a panD, BAPAT, or HPDH sequence (such as SEQ ID NO: 54, 56, or 58, respectively) includes one or more amino acid substitutions, such as conservative substitutions (for example at 1, 2, 5, or 10 residues). Examples of conservative substitutions are provided elsewhere herein.

The panD, BAPAT, and HPDH gene expressed in a fungus, in particular examples, includes a sequence that

encodes a panD, BAPAT, and HPDH protein having at least at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to a panD, BAPAT, and HPDH protein sequence, such as SEQ ID NO: 54, 56, or 58, respectively, wherein the variant protein has the biological activity of panD, BAPAT, or HPDH, respectively. In a specific example, the panD, BAPAT, and HPDH gene expressed in a Δ cadA fungus encodes the protein shown in SEQ ID NO: 54, 56, and 58, respectively.

One skilled in the art will appreciate that additional panD, BAPAT, and HPDH sequences can be identified. For example, panD, BAPAT, and HPDH nucleic acid molecules that encode a panD, BAPAT, and HPDH protein, respectively can be identified and obtained using molecular cloning or chemical nucleic acid synthesis procedures and techniques, including PCR. In addition, nucleic acid sequencing techniques and software programs that translate nucleic acid sequences into amino acid sequences based on the genetic code can be used to determine whether or not a particular nucleic acid has any sequence homology with panD, BAPAT, or HPDH sequences. Sequence alignment software such as MEGALIGN (DNASTAR, Madison, Wis., 1997) can be used to compare various sequences.

In addition, nucleic acid hybridization techniques can be used to identify and obtain a nucleic acid molecule that encodes a panD, BAPAT, or HPDH protein. Briefly, any known panD, BAPAT, or HPDH nucleic acid molecule, or fragment thereof, can be used as a probe to identify similar nucleic acid molecules by hybridization under conditions of moderate to high stringency. Such similar nucleic acid molecules then can be isolated, sequenced, and analyzed to determine whether the encoded protein is a panD, BAPAT, or HPDH protein.

In one example, exogenous panD, BAPAT, and/or HPDH nucleic acid sequences are introduced into *A. pseudoterreus* or *A. terreus* using protoplast transformation, for example as described in Example 1 (and described above).

F. Methods of Increasing panD, BAPAT, and HPDH Expression

In some examples, a native *A. pseudoterreus* or *A. terreus* fungi does not have or express panD, BAPAT, and/or HPDH nucleic acid sequences. Thus, in some examples, expression of these genes is increased by introducing panD, BAPAT, and/or HPDH nucleic acid coding sequences (such may be codon optimized) into the *A. pseudoterreus* or *A. terreus* fungi.

In some examples, a native *A. pseudoterreus* or *A. terreus* fungi does express native panD, BAPAT, and/or HPDH nucleic acid sequences. Thus, in some examples, expression of these genes is upregulated by introducing additional copies of panD, BAPAT, and/or HPDH nucleic acid coding sequences (such may be codon optimized) into the *A. pseudoterreus* or *A. terreus* fungi. As used herein, "up-regulated" gene means that expression of the gene or gene product (e.g., protein) has been up-regulated, for example by introduction of additional copies of the appropriate gene or coding sequence into the fungus (or other molecular biology methods), such that the introduced nucleic acid sequence is expressed, resulting in increased expression or biological activity of the encoded gene product. In some embodiments, introduction of one or more transgenes including panD, BAPAT, and/or HPDH coding sequences into a native *A. pseudoterreus* or *A. terreus* fungi increases expression of panD, BAPAT, and/or HPDH by at least 20%, at least 40%, at least 50%, at least 100%, at least 150%, at least 200%, at least 300%, or at least 500%, for example relative to the

parental fungal strain without the introduced panD, BAPAT, and/or HPDH coding sequences. The term “increased” or “up-regulated” as used herein with respect to a cell and a particular gene or protein activity refers to a higher level of activity than that measured in a comparable cell of the same species. For example, a particular fungi having increased or up-regulated panD, BAPAT, and/or HPDH activity has increased panD, BAPAT, and/or HPDH activity if a comparable fungi having native panD, BAPAT, and/or HPDH activity has less detectable panD, BAPAT, and/or HPDH activity (for example as measured by gene or protein expression).

In one example, a strain of *Aspergillus* is transformed with a vector which has the effect of up-regulating a panD, BAPAT, and/or HPDH gene (such as a native or non-native panD, BAPAT, and/or HPDH gene). This can be done by introducing one or more panD, BAPAT, and/or HPDH coding sequences (such as a gene sequence), whose expression is controlled by elements such as promoters and the like which control gene expression, by introducing a nucleic acid sequence which itself (or its encoded protein) can increase panD, BAPAT, and/or HPDH protein activity in the fungus, or by introducing another molecule (such as a protein or antibody) increases panD, BAPAT, and/or HPDH protein activity in the fungus. For example, a panD, BAPAT, and/or HPDH gene can be up-regulated by introduction of a vector that includes one or more panD, BAPAT, and/or HPDH sequences (such as 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 panD, BAPAT, and/or HPDH sequences or copies of such sequences) into the desired fungus. In some examples, such panD, BAPAT, and/or HPDH sequences are from different fungal species, can be multiple copies from a single species, or combinations thereof, such as panD, BAPAT, and/or HPDH sequences from at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 different fungal species. In some examples, the panD, BAPAT, and/or HPDH sequence(s) introduced into the fungus is optimized for codon usage. Thus, the disclosure in some examples provides transformed fungi that include at least one exogenous nucleic acid molecule which includes a panD, BAPAT, and/or HPDH gene or coding sequence (such as a nucleic acid sequence encoding SEQ ID NO: 54, 56, or 58, respectively), for example in combination with Δ cadA. In one example, such transformed cells produce more 3HP, for example relative to a comparable fungus with a native cadA.

In one example, the cre-lox system is used for site specific recombination of DNA (for example see Steiger et al., *Appl. Environ. Microbiol.* 77(1):114, 2011). Using recombination techniques, the targeted gene of interest (e.g., cadA) can be deleted in the *Aspergillus* genome and replaced with one or more copies of a non-native panD, BAPAT, and/or HPDH sequence (for example in *A. terreus*, replacing one or both *A. terreus* cadA sequences with panD, BAPAT, and/or HPDH sequences from *A. nidulans* or *A. flavus*) flanked by the lox sites. Transient expression (by electroporation of a suicide plasmid containing the cre gene under control of a promoter that functions in *Aspergillus*) of the cre recombinase should result in efficient elimination of the lox flanked marker. This process will produce a fungus containing the desired insertion mutation and one copy of the lox sequence.

In one example, a transgene is generated and expressed in the desired fungal cell, such as an Δ cadA fungal cell, to increase panD, BAPAT, and HPDH expression. For example, one or more transgenes can include a panD, BAPAT, and HPDH genomic or cDNA sequence (such as one having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence

identity to any panD, BAPAT, and HPDH sequence provided herein), for example operably linked to one or more promoters, such as *gpdA* and *enol*. In one example, the promoter has at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 74 and/or 77. In some examples, the transgene further includes a *trpC* transcriptional terminator sequence of *A. nidulans*, for example downstream of the panD, BAPAT, and/or HPDH sequence. As an alternative to *trpC*, other transcriptional terminators can be used, such as promoters which include a transcriptional terminators (e.g., *ArsA7*, *ArsA-37*, polyubiquitin (*ubi4*)). In one example, the *trpC* transcriptional terminator has at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 83 or 86. In one example, the *trpC* transcriptional terminator comprises or consists of the sequence shown in SEQ ID NO: 83 or 86. In some examples, the transgene further includes a *ptrA* sequence, for example downstream of the *trpC* transcriptional terminator sequence. As an alternative to *ptrA*, the bleomycin gene or *bar* gene can be used. In one example, the *ptrA* sequence has at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 89. In one example, the *ptrA* sequence comprises or consists of the sequence shown in SEQ ID NO: 89.

In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 59, 62, 65, 68, 71, and/or 74. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 59, 62, 65, 68, 71, and/or 74.

In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 77, 80, and/or 83. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 77, 80, and/or 83.

In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 86, 89, and/or 92. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 86, 89, and/or 92.

In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 89 and/or 92. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 89 and/or 92.

In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 95 and/or 97. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 95 and/or 97.

In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 59, 62, 65, 68, 71, 74, 77, 80, 83, 86, 89, and/or 92. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 59, 62, 65, 68, 71, 74, 77, 80, 83, 86, 89, and/or 92.

In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity

to SEQ ID NO: 59, 62, 65, 68, 71, 74, 77, 80, 83, 89, and/or 92. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 59, 62, 65, 68, 71, 74, 77, 80, 83, 89, and/or 92.

G. Measuring Gene Expression

A Δ cadA fungus expressing panD, BAPAT, and/or HPDH can be identified using known methods. For example, PCR and nucleic acid hybridization techniques, such as Northern, RT-PCR, and Southern analysis, can be used to confirm that a fungus expresses panD, BAPAT, and/or HPDH such as an increase in the panD, BAPAT, and/or HPDH copy number. Immunohisto-chemical and biochemical techniques can also be used to determine if a cell expresses panD, BAPAT, and/or HPDH by detecting the expression of the panD, BAPAT, and/or HPDH peptide encoded by panD, BAPAT, and/or HPDH. For example, an antibody having specificity for panD, BAPAT, and/or HPDH can be used to determine whether or not a particular fungus has increased panD, BAPAT, and/or HPDH protein expression, respectively. Further, biochemical techniques can be used to determine if a cell has increased panD, BAPAT, and/or HPDH expression by detecting a product produced as a result of the expression of the peptide. For example, production of 3-HP by Δ cadA *A. terreus* or *A. pseudoterreus* can indicate that such a fungus expresses panD, BAPAT, and HPDH.

H. Measuring 3-HP Production

Methods of determining whether a genetic inactivation of cadA in combination with expression of panD, BAPAT, and HPDH in *Aspergillus* increases 3-HP production, for example relative to the same strain with a native cadA sequence, (such as a parental strain) include HPLC.

Methods of Producing Aconitic Acid

The recombinant Δ cadA fungi can be used to produce aconitic acid (for example for as a building block for other materials, such as polymers). Such fungi can be from any *Aspergillus* species, such as *Aspergillus terreus* or *pseudoterreus*. For example, the disclosure provides methods of making aconitic acid (such as cis-acconitic acid, trans-acconitic acid, or both), which can include culturing Δ cadA fungi under conditions that permit the fungus to make aconitic acid, for example in Riscaldati medium.

In some examples, the fungi are cultured at room temperature (e.g., 20-35° C.) at normal atmospheric pressure (e.g., 1 atm). In some examples, the method includes purifying or isolating the aconitic acid, for example from the culture media or from the cultured fungus. In some examples, the aconitic acid is isolated at least 2 days, at least 3 days, at least 5 days, at least 8 days or at least 10 days after the start of culturing, such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 days after the start of culturing.

Methods of making aconitic acid include culturing Δ cadA fungi under conditions that permit the fungus to make aconitic acid. In general, the culture media and/or culture conditions can be such that the fungi grow to an adequate density and produce aconitic acid efficiently. In one example the Δ cadA fungi are cultured or grown in an acidic liquid medium, such as Riscaldati medium (100 g Glucose, 0.11 g KH_2PO_4 , 2.36 g $(\text{NH}_4)_2\text{SO}_4$, 2.08 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.074 g NaCl, 0.13 g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1 ml of 1000× trace elements in 1000 ml DI water, adjust pH to 3.4 with H_2SO_4 , 1000× trace elements contains 1.3 g/L $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 5.5 g/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.7 g/L $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$). In one example the Δ cadA fungi are cultured or grown in a liquid medium having an initial pH of less than

4, such as less than 3.5, for example about pH 3 to 4, 3.5 to 4, 3.3 to 3.5, for example pH 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9 or 4. In some examples the Δ cadA fungi are cultured or grown in a liquid Riscaldati medium at about 20 to 35° C. (such as 20° C. to 30° C., 25° C. to 30° C., 28 to 32° C., or 30° C.) with rotation (such as at least 100 rpm, at least 120 rpm, such as 150 rpm) at normal pressure.


In one example, the fungi are grown in culture containers (such as baffled flasks, and in some examples are silanized (5% solution of dichlorodimethylsilane in heptane (Sigma, St. Louis, Mo.)). Each culture container is inoculated with spores (such as at least 10^6 spores/ml [agree?]) and incubated for at least 3 days, at least 4 days, at least 5 days, or at least 10 days at 30° C. and 100 to 200 rpm to obtain aconitic acid.

In one example, the Δ cadA fungi produce more aconitic acid than a corresponding fungus with wild-type cadA. In specific examples, the Δ cadA fungi produce at least 1 g/l of total aconitic acid after 4 days, for example at least 2 g/l, at least 3 g/l, at least 4 g/l, at least 5 g/l, at least 6 g/l, at least 7 g/l, at least 8 g/l, at least 9 g/l or at least 10 g/l after at least 5 days, at least 6 days, at least 7 days, at least 8 days, or at least 10 days, such as after 4 to 6 days, 8 to 10 days, or 4 to 5 days) when grown in Riscaldati medium at 30° C. with 150 rpm shaking. In specific examples, the Δ cadA fungi produce at least 1 g/l of cis-acconitic acid after 4 days, for example at least 2 g/l after at least 5 days, at least 6 days, at least 7 days, at least 8 days, or at least 10 days, such as after 4 to 6 days, 8 to 10 days, or 4 to 5 days when grown in Riscaldati medium at 30° C. with 150 rpm shaking. In specific examples, the Δ cadA fungi produce at least 1 g/l of trans-acconitic acid after 6 days, for example at least 2 g/l, at least 3 g/l, at least 4 g/l, at least 5 g/l, at least 6 g/l, at least 7 g/l, at least 8 g/l, at least 9 g/l or at least 10 g/l after at least 7 days, at least 8 days, or at least 10 days, such as after 6 to 12 days, 5 to 10 days, or 6 to 10 days) when grown in Riscaldati medium at 30° C. with 150 rpm shaking.

In some examples, the method further includes isolating the aconitic acid made by the Δ cadA fungi. Once produced, any method can be used to isolate the aconitic acid. For example, separation techniques (such as filtration) can be used to remove the fungal biomass from the culture medium, and isolation procedures (e.g., filtration, distillation, precipitation, electrodialysis, and ion-exchange procedures) can be used to obtain the aconitic acid from the broth (such as a fungi-free broth). In addition, the aconitic acid can be isolated from the culture medium after the aconitic acid production phase has been terminated.

Methods of Producing 3-HP

The recombinant Δ cadA fungi that also express panD, BAPAT, and HPDH can be used to produce 3-HP



(for example for as a building block for other materials, such as acrylonitrile, acrylic acid by dehydration, malonic acid by oxidation, esters by esterification reactions with alcohols, and reduction to 1,3 propanediol). Such fungi can be from any *Aspergillus* species, such as *Aspergillus terreus* or *pseudoterreus*. For example, the disclosure provides

methods of making 3-HP, which can include culturing Δ cadA fungi that also express panD, BAPAT, and HPDH under conditions that permit the fungus to make 3-HP, for example in Riscaldati medium (such as modified Riscaldati medium with 20 \times trace elements).

In some examples, the Δ cadA fungi that also express panD, BAPAT, and HPDH are cultured at room temperature (e.g., 20-35° C.) at normal atmospheric pressure (e.g., 1 atm). In some examples, the method includes purifying or isolating the 3-HP, for example from the culture media or from the cultured fungus. In some examples, the 3-HP is isolated at least 2 days, at least 3 days, at least 5 days, at least 7 days, at least 8 days or at least 10 days after the start of culturing, such as 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 days after the start of culturing.

Methods of making 3-HP include culturing Δ cadA fungi that also express panD, BAPAT, and HPDH under conditions that permit the fungus to make 3-HP. In general, the culture media and/or culture conditions can be such that the fungi grow to an adequate density and produce 3-HP efficiently. In one example the Δ cadA fungi that also express panD, BAPAT, and HPDH are cultured or grown in an acidic liquid medium, such as Riscaldati medium (100 g Glucose, 0.11 g KH_2PO_4 , 2.36 g $(\text{NH}_4)_2\text{SO}_4$, 2.08 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.074 g NaCl, 0.13 g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1 ml of 1000 \times trace elements in 1000 ml DI water, adjust pH to 3.4 with H_2SO_4 , 1000 \times trace elements contains 1.3 g/L $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 5.5 g/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.7 g/L $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, which may include 20 \times trace elements). In one example the Δ cadA fungi are cultured or grown in a liquid medium having an initial pH of less than 4, such as less than 3.5, for example about pH 3 to 4, 3.5 to 4, 3.3 to 3.5, for example pH 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9 or 4. In some examples the Δ cadA fungi that also express panD, BAPAT, and HPDH are cultured or grown in a liquid modified Riscaldati medium with 20 \times trace elements at about 20 to 35° C. (such as 20° C. to 30° C., 25° C. to 30° C., 28 to 32° C., or 30° C.) with rotation (such as at least 100 rpm, at least 120 rpm, such as 150 or 200 rpm) at normal pressure.

In one example, the fungi are grown in culture containers (such as baffled flasks, and in some examples are silanized (5% solution of dichlorodimethylsilane in heptane (Sigma, St. Louis, Mo.)). Each culture container is inoculated with spores (such as at least 10⁶ spores/ml) and incubated for at least 3 days, at least 4 days, at least 5 days, or at least 10 days at 30° C. and 100 to 300 rpm (such as 150 or 200 rpm) to obtain 3-HP.

In one example, the cadA fungi that also express panD, BAPAT, and HPDH produce more 3-HP than a corresponding fungus with wild-type cadA (either with or without panD, BAPAT, and HPDH expression). In specific examples, the Δ cadA fungi that also express panD, BAPAT, and HPDH produce at least 0.1 g/l of 3-HP after at least 4 days, for example at least 0.2 g/l, at least 0.25 g/l, at least 0.3 g/l, at least 0.4 g/l, at least 0.5 g/l, at least 0.6 g/l, at least 0.7 g/l, at least 0.8 g/l, at least 0.9 g/l, at least 1.1 g/l, at least 1.2 g/l, at least 1.5 g/l, or at least 1.6 g/l after at least 5 days, at least 6 days, at least 7 days, at least 8 days, or at least 10 days, such as after 4 to 6 days, 8 to 10 days, or 4 to 5 days, when grown in Riscaldati medium (such as modified Riscaldati medium with 20 \times trace elements) at 30° C. with 150 rpm shaking.

In some examples, the method further includes isolating the 3-HP made by the Δ cadA fungi. Once produced, any method can be used to isolate the 3-HP. For example, separation techniques (such as filtration) can be used to

remove the fungal biomass from the culture medium, and isolation procedures (e.g., filtration, distillation, precipitation, electro dialysis, and ion-exchange procedures) can be used to obtain the 3-HP from the broth (such as a fungi-free broth). In addition, the 3-HP can be isolated from the culture medium after the 3-HP production phase has been terminated.

Compositions and Kits

Also provided by the present disclosure are compositions that include isolated Δ cadA fungi (which in some examples also express panD, BAPAT, and HPDH, such as exogenous panD, BAPAT, and HPDH proteins), such as a medium for culturing, storing, or growing the fungus. In some examples, the Δ cadA fungi and Δ cadA fungi which express panD, BAPAT, and HPDH in the composition are freeze dried or lyophilized.

Also provided by the present disclosure are kits that include isolated Δ cadA fungi (which in some examples also express panD, BAPAT, and HPDH, such as exogenous panD, BAPAT, and HPDH proteins), such as a kit that includes a medium for culturing, storing, or growing the fungus. In some examples, the Δ cadA fungi and Δ cadA fungi which express panD, BAPAT, and HPDH in the kit are freeze dried or lyophilized.

Exemplary mediums include that can be in the disclosed compositions and kits include solid medium (such as those containing agar, for example complete medium (CM) or minimal medium (MM)) and liquid media (such as a fermentation broth, such as CM, MM, or CAP medium). In one example, the kit or composition includes Riscaldati medium (100 g Glucose, 0.11 g KH_2PO_4 , 2.36 g $(\text{NH}_4)_2\text{SO}_4$, 2.08 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.074 g NaCl, 0.13 g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 1 ml of 1000 \times trace elements in 1000 ml DI water, adjust pH to 3.4 with H_2SO_4 , 1000 \times trace elements contains 1.3 g/L $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 5.5 g/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g/L $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.7 g/L $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$), for example

	Conc. (g/L)	Amount	Notes
Glucose	100	100 g	
KH_2PO_4	0.11	0.11 g	
$(\text{NH}_4)_2\text{SO}_4$	2.36	2.36 g	
$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$	2.08	2.08 g	
NaCl	0.074	0.074 g	
$\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$	0.13	0.13 g	
$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$	0.0013	0.0013 g	Use 1000 X soln.
$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	0.0055	0.0055 g	"
$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	0.0002	0.0002 g	"
$\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$	0.0007	0.0007 g	"
DI Water (L)		1 L	
Autoclave Time		15 min for small flasks 30 min for large flasks 30-60 for fermenter	
Comments:		Adjust to pH = 3.4 with H_2SO_4	

In one example, the kit or composition includes a modified Riscaldati medium with 20 \times trace elements, for example 20 times of the following

$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$	0.0013	0.0013 g	Use 1000 X soln.
$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$	0.0055	0.0055 g	"
$\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$	0.0002	0.0002 g	"
$\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$	0.0007	0.0007 g	"

Materials and Methods

This example describes methods used in the experiments described in Examples 2-6 below.

Strains and Vector.

The parental wild type *A. pseudoterreus* strain ATCC 32359 was from ATCC. The hygromycin phosphotransferase (hph) marker cassette was amplified from vector pCB1003.

Growth Conditions.

All strains were maintained on complete medium (CM) agar and conidia of spore were harvested from cultures grown for five days on complete medium (CM) plate (10 g Glucose, 2 g Triptase peptone, 1 g yeast extract, 1 g casamino acids, 50 ml 20×NO₃ Salts, 1 ml of 1000×Trace elements, 1 ml of 1000× Vitamin stock, in 1000 ml DI water, pH to 6.5), 20×NO₃ Salts contains in g/l, Na₂NO₃, 120; KCl, 10.4 g; MgSO₄·7H₂O, 10.4 g; KH₂PO₄, 30.4 g. 1000× vitamin solution contains in per 100 ml H₂O: Biotin, 0.01 gm; pyridoxinHCl, 0.01 gm; thiamineHCl, 0.01 gm; riboflavin, 0.01 gm; paba, 0.01 gm; nicotinic acid, 0.01 gm, filtered and stock at 4° C. 1000× trace element contains in per 100 ml H₂O: ZnSO₄·7H₂O, 2.2 g; H₃B03, 1.1 g; MnCl₂·4H₂O, 0.5 g; FeSO₄·7H₂O, 0.5 g; CoCl₂·6H₂O, 0.17 g; CuSO₄·5H₂O, 0.16 g; Na₂MoO₄·2H₂O, 0.15 g; Na₂EDTA, 5 g, add the compounds in order, boil and cool to 60° C. Adjust pH to 6.5 with KOH. Cool to room temperature. Adjust volume to 100 ml with distilled water.

The transformants were selected for hygromycin resistance on minimum media (MM) (10 g Glucose, 50 ml 20×NO₃ Salts, 1 ml of 1000×Trace elements, 1 ml of 1000× Vitamin stock, 1000 ml DI water, pH to 6.25-6.5, hygromycin 100 ug/ml). 0.5×10⁸ conidia were inoculated into 50 ml of production media for itaconic acid production (Riscal-dati medium) as described previously (100 g Glucose, 0.11 g KH₂PO₄, 2.36 g (NH₄)₂SO₄, 2.08 g MgSO₄·7H₂O, 0.074

g NaCl, 0.13 g CaCl₂·2H₂O, 1 ml of 1000× trace elements in 1000 ml DI water, adjust pH to 3.4 with H₂SO₄, 1000× trace elements contains 1.3 g/L ZnSO₄·7H₂O, 5.5 g/L FeSO₄·7H₂O, 0.2 g/L CuSO₄·5H₂O, 0.7 g/L MnCl₂·4H₂O). Cultivation was performed at 30° C. on a rotary shaker at 150 rpm. At intervals during the incubation period, three single flasks were harvested for HPLC analysis, biomass measurement and RNA extraction. All experiments were replicated three times, and the standard deviation of the itaconic acid concentrations or dry weight was always less than 10% of the mean. For collecting samples for EST analysis, *A. pseudoterreus* was grown in 20 liter stirred tank bioreactor.

Construction of Deletion Mutants.

The deletion mutants were constructed by homologous double crossover with fusion PCR products. Synthetic oligos used for each construct are described in Table 1. Oligonucleotides were from IDT (Coraville, Iowa). Ex Taq polymerase (TaKaRa, Japan) was used to generate DNA constructs for making gene knockouts. Briefly, the 5' flanking region (~1.5 kb) of the target gene was amplified by primer pair F1 and R3. The 3' flanking region (~1.5 kb) of the target genes was amplified by primer pair F4 and R6. R3 and F4 carried 20-25 bases complementary to 5' and 3' ends of the hph cassette, respectively. The hph marker cassette was amplified from pCB1003 with the hphF and hphR primers that carried 30 bases complementary to the 3' end of the 5' flanking region and the 5' of the 3' flanking region, respectively. The three fragments, including the 5' flanking region, the hph marker cassette and the 3' flanking region were mixed in 1:3:1 molar ratio and combined by overlap PCR during the second round PCR. In the third round of PCR, the fusion PCR product was amplified with a nested primer pair (F2 and R5). This final PCR product carried a hygromycin marker cassette flanked by sequences homologous to the upstream and downstream regions of the target gene. 1-2 ug of the final product was used to transform strain *A. pseudoterreus* strain ATCC 32359.

TABLE 1

Primers for making deletion constructs		
gene	targeted primer name	primer sequence (SEQ ID NO)
tf	at tff1	gagccatagccatgcaagcg (1)
	at tff2	atagagtccttgatgagacg (2)
	at tfr5	gtggatttcgaggttccttgc (3)
	at tfr6	gaagtagaacctgtggatcg (4)
	at hphf tfr3	tgacctccactagctccagcactactagataggccggtttagagagtgc (5)
	at hphr tff4	aatagagttagtgccgaccggcgcttcgacgacagctctgcactctcc (6)
	at tfr3hphf	ggcactctctaaccggcctatctagttagtgcggagctagtggaggtca (7)
	at tff4hphr	ggagagtgcagagctgtcgtcgaagcggcgggctcggcactctactctatt (8)
mttA	at motf1	gctgcatactcggattacgc (9)
	at motf2	Gaaaaggactcggagtagc (10)
	at motr5	cagaccaaggagcttctctg (11)
	at motr6	cattaagccacaggcttgcg (12)
	athphfmotr3	tgacctccactagctccagcaaatatggatgctgttcggttcgctgctgg (13)
	athphrmotr4	aatagagttagtgccgaccgtgacgaggatgtgctgagtcacaacaagc (14)
	at motr3hphf	ccagcagcggcaacgaacagcatccatattgctggagctagtggaggtca (15)
	at motf4hphr	gctttgtttggactcagcaatcctcgtcagcggctcggcatctactctatt (16)
cadA	at cadf1	ctccagtaacagaaccgacc (17)
	at cadf2	gaacttcaactgcccattgg (18)
	at cadr5	ggacactccaagaggataagg (19)
	at cadr6	gctcatcacattgtttgccc (20)
	at hphfcadr3	tgacctccactagctccagcggcgaatttaagaggacgatcttcgctgcg (21)
	at hphrcadf4	Aatagagttagtgccgaccgtcagcctggacaggtccaccgacattagcc (22)
	at cadr3hphf	cgacgcaagatcgtcctcttaattgaccgctggagctagtggaggtca (23)
	at cadf4hphr	ggctaagtgcggtagcctgtccaggctgacggctcggcactctactctatt (24)

TABLE 1-continued

Primers for making deletion constructs		
gene	targeted primer name	primer sequence (SEQ ID NO)
mfsA	mfsf1	tgatgagctgaattcggttc (25)
	mfsf2	tatagccagcttttgctgtg (26)
	mfsr5	catagcgttcagagtgttg (27)
	mfsr6	ccatttcaatgctttgtgcg (28)
	mfsr3hphf	ccataccacccttacccctcttgagtggtccgctggagctagtgagggtca (29)
	mfsf4hphr	gctgtggcctcctcggcgattacgcaatattcggctcggcatctactotatt (30)
	hphf450r3	tgacctccactagctccagcggacactccaagaggtaaggggtgatgg (31)
	hphrmfsf4	aatagagttagtgccgaccgaatattgctgtaatcgccaggaggccacagc (32)
	p450	p450f1
p450f2		cttcaatcgcaccgacatcc (34)
p450r5		tcgtgtagacaagtcacagtc (35)
p450r6		ctataccactctagtgtg (36)
p450r3hphf		cctctgctcaggttgttttcgaacaggagcgtggagctagtgagggtca (37)
p450f4hphr		cggaaatgcagataggcatcacagtcacagaacggctcggcatctactotatt (38)
hphfp450r3		tgacctccactagctccagcgtcctcgttccgaaaacaacctgagcagagg (39)
hphrp450f4		aatagagttagtgccgaccgtctcgtgactgtgatgctatctgcattccg (40)

Transformation of *A. pseudoterreus* Protoplasts.

10⁸ conidia of *A. pseudoterreus* ATCC 32359 were used to inoculate 300 ml Erlenmeyer baffle flasks containing 100 ml of complete media. The cultures were grown overnight (16-18 hrs) at 30° C. and 150 rpm. The mycelia were then harvested by filtering the culture through miracloth and rinsing the mycelia mat with sterile water. The protoplasts were prepared by treating mycelia (mass of approximately 1-2 beans) with 20 mg/ml lysing enzyme (L1412, Sigma) dissolved in 20 ml of osmotic wash buffer (0.5M KCl, 10 mM sodium phosphate, pH 5.8) for 2 h. Protoplasts were collected by filtering protoplasts through sterile miracloth into a 50 ml screw cap centrifuge tube and centrifuging at 1000xg for 10 min at 4° C. Protoplasts were then washed twice with 20 ml Washing Solution (0.6M KCl, 0.1M Tris/HCl, pH 7.0) and once in 10 ml Conditioning Solution (0.6M KCl, 50 mM CaCl₂, 10 mM Tris/HCl, pH 7.5). For transformation, 1-2 ug DNA was added to 2x10⁷ protoplasts in 0.1 ml Conditioning Solution. A control reaction without added DNA was performed at the same time. 25 µl of PEG solution (25% PEG8000, 0.6M KCl, 50 mM CaCl₂, 10 mM Tris/HCl, and pH 7.5) was added and the protoplasts were incubated for 20 min on ice. An additional 500 µl of the PEG was added using a wide bore pipette tip and carefully mixed with the protoplasts by gently pipetting up and down 1-2 times. The protoplast solution was then incubated for 5 min on ice. 1 ml of cold Conditioning Solution was added and mixed by gently inverting the tube several times. Then the protoplast suspension was mixed with 12 ml of 50° C. selection agar (Minimal media+0.6M KCl+1.5% Agar+100 µg/ml hygromycin) contained in a 15 ml screwcap centrifuge tube. The tubes were then mixed by inverting the tubes 3-4 times and poured directly onto the petri dish plates. The control reaction was divided into a positive control plate (no selection antibiotics in the top agar and bottom plates) and a negative control (with selection hygromycin in top and bottom agar). The solidified plates were inverted and incubated overnight at 30° C. The next day, the plates were then overlaid with 15 ml of Minimal Medium (MM) containing 150 µg/ml hygromycin. Colonies should start to appear after 3-4 days. The transformants were excised and transferred to MM plate containing 100 µg/ml hygromycin. Correct transformants on the hygromycin plate were confirmed by PCR approaches and southern blot.

Dry Mass Measurement.

Dry mass at each time point was determined by harvesting the mycelium on miracloth by suction filtration and washed twice with 50 ml distilled water. Subsequently, the dry weight was determined by drying it overnight in pre-weighed tubes on lyophilizer.

HPLC.

The content of itaconic acid, aconitic acid, and glucose in each sample collected from filtration (0.22 µm) was assayed by a high-pressure liquid chromatography (HPLC) on a Bio-Rad Aminex HPX-87H ion exclusion column (300 mmx7.8 mm). Columns were eluted with Sulfuric acid (0.005 M) at a flow rate of 0.55 mL/min. The sample volume was 10-100 µl, and IA was detected at 210 nm with a Waters 2414 refractive index detector.

RNA Isolation and Transcript Analysis by Quantitative Real Time RT-PCR.

Wild type and tf deletion strains were grown in Riscaldati medium at 30° C. After 3 days growth, mycelia were harvested, pressed dry between paper towels and immediately flash frozen in liquid nitrogen. The entire sample was then ground in a mortar and pestle with liquid nitrogen. Approximately 100 mg samples (about 0.1 ml) were extracted using Trizol® reagent (Chomczynski, *BioTechniques* 1993, 15(3):532-534, 536-537) and the resulting RNA was converted to cDNA using high capacity RNA-to-DNA kit (Applied Biosystems). Quantitative RT-PCR were performed in 50 µl reactions containing 25 µl of Power SYBR green PCR master mix (Applied Biosystems), 50 ng cDNA (from 50 ng RNA) and 0.2 µM forward and reverse primers. The RT-PCR primers used for analysis of the mttA, cadA, mfsA genes and benA (β-tubulin) as endogenous control gene are listed in Table 3. There are two additional controls, one is a no RT (without adding RT enzyme mix) control to estimate contamination from genomic DNA, and the other is no-template controls for each primer pair to measure effect from primer dimer formation. Amplification was performed using 7900HT Fast Real-Time PCR system (Applied Biosystems) programmed to initially hold at 95° C. for 10 min and then to complete 45 cycles of 95° C. for 15 s, 60° C., for 60 s. The data were analyzed using the comparative C_T method (e.g., see Schmittgen et al., *Analytical biochemistry* 2000, 285(2):194-204)

Example 2

Expression Profile of Itaconic Acid Gene Cluster in *A. pseudoterreus*

RNA samples were prepared from three different growth stages of *A. pseudoterreus* in the itaconic acid production process. The stages were 1) "pre-production," before itaconic acid production begins, 2) "production onset," the beginning of itaconic acid production correlated with phosphate depletion, and 3) "production," early in the phase of maximum itaconic acid production rate (FIG. 2). EST data revealed four genes in the cluster having high expression frequency both in the onset phase and production phase, but not in the pre-production phase (Table 2). These genes were *tf*, *mttA*, *cadA*, and *mfsA*.

TABLE 2

Number of ESTs per gene at three stages of itaconic acid production				
Broad Institute Gene No.	Gene Description	Pre-Production	Production Onset	Production
ATEG_09968.1	upstream flanking gene; <i>lovE</i>	0	0	0
ATEG_09969.1	<i>tf</i>	0	4	4
ATEG_09970.1	<i>mttA</i>	0	81	93
ATEG_09971.1	<i>cadA</i>	0	77	110
ATEG_09972.1	<i>mfsA</i>	0	6	7
ATEG_09973.1	<i>p450</i>	0	7	11
ATEG_09974.1	downstream flanking gene	0	0	0
ATEG_09817.1	control; <i>gapdh</i>	31	51	43

cadA has 77 ESTs at the beginning of itaconic acid (IA) production and 110 ESTs during the IA production, while *mttA* has 81 and 93 ESTs respectively in each stage. Both have no transcript detected before IA is produced. Transcription factor (*tf*) and *mfsA*, like *cadA* and *mttA*, did not show any expression before IA production, but had significant levels of transcription following the initiation of itaconic acid production (Table 2).

When genes upstream and downstream of *tf*, *cadA*, *mttA* and *mfsA* were examined, a similar expression pattern was not observed. No transcript was detected for either upstream or downstream genes in any stage of IA production except for *p450*. Control gene *gapdh*, which is far away from this region, showed high expression through the whole growth stage. This EST data clearly demonstrated that four genes *tf*, *cadA*, *mttA* and *mfsA* have the same expression pattern and are closely related to the IA production process. In addition, these four genes are in the same cluster. They are turned on strongly at the onset of IA production and persists through the production phase (FIG. 2).

Example 3

Effect of *Tf*, *cadA*, *mttA* and *mfsA* Deletion on Itaconic Acid Production in *A. pseudoterreus*

A transformation system was developed to allow for transformation of *A. pseudoterreus* (see Example 1). This system was used to generate recombinant knockout strains for each of the endogenous *tf*, *cadA*, *mttA* and *mfsA* genes. The KO mutant strains were confirmed by PCR and southern blot. The transformation protocol gave very high frequency of homologous deletion, 8 out of 10 had the correct deletion.

This high deletion frequency may be due to the presence of a *ku* gene mutation in the genome of wild-type *A. pseudoterreus*.

Biomass accumulation and itaconic acid (IA) production of each of the four knockout mutants and wild type *A. pseudoterreus* were measured at day 5. All strains, including wild type, had similar biomass accumulation (FIG. 3A). There is no significant difference in biomass among these five strains, indicating that deletion of these genes does not cause a noticeable growth defect.

However, the yield of IA was significantly lower in all four deletion strains (Δtf , $\Delta cadA$, $\Delta mttA$ and $\Delta mfsA$) when compared to wild type *A. pseudoterreus*. After 5 days growth in the Riscaldati medium, the Δtf strain had only generated ~3 g/l IA, compared to the wild type strain, which generated ~24 g/l of IA (about an 8-fold decrease). No detectable IA was produced by the $\Delta cadA$ and $\Delta mttA$ strains. $\Delta mfsA$ produced around 16 g/L itaconic acid, about 2/3 of wild type *A. pseudoterreus*.

These observations demonstrate that *tf*, *mttA*, *cadA* and *mfsA* genes play a role in itaconic acid production.

Example 4

Production Kinetics of Itaconic Acid in Wild Type and *Tf* Deletion Strain

To test the production kinetics in the deletion strains, Δtf and wild type *A. pseudoterreus* strain ATCC 32359 were tested for IA production during the growth on a rotary shaker for 7 days. IA was analyzed by HPLC for 2, 4, 6 and 7 day cultures.

As shown in FIG. 4, the IA yield plateaued at day 7 in both Δtf and wild type strains. Interestingly, the IA yield in Δtf (50) is much lower than that of wild type (35 g/l), a decrease of about 7-fold. Thus, the Δtf strain produces IA at slower rate with a lower maximum IA yield than the wild type strain.

Example 5

Tf Regulation

The effects of *tf* gene deletion on the transcription level of other genes in the cluster were investigated by real-time reverse transcription PCR (RT-PCR). In the both Δtf and wild type strains, expression level of each gene was analyzed by RT-PCR by measuring *mttA*, *cadA*, *mfsA* mRNA levels using primers specific for those genes (Table 3).

TABLE 3

primers for real-time RT-PCR analysis of cluster gene transcript level		
Gene targeted	Primer name	Primer sequence (SEQ ID NO:)
<i>mttA</i>	<i>mttF</i>	Gctttcaatgtggttctact (41)
	<i>mttR</i>	ctocatcacctacccttctc (42)
<i>cadA</i>	<i>cadF</i>	gaagtgtgggatctggc (43)
	<i>cadR</i>	gggttcggattattgtgaag (44)
<i>mfsA</i>	<i>mfsF</i>	caagaacagtttggcctgag (45)
	<i>mfsR</i>	gcggacatcatacaatctgg (46)
<i>benA</i>	β -tubulinF	ttgtcgatgttctgctgc (47)
	β -tubulinR	tggcgttgtaaggctcaacc (48)

As shown in FIG. 5, in Δ tf strains, mRNA level of mttA decreased 57 fold, cadA mRNA level decreased 37 fold, and mfsA decreased 23 fold, as compared to their expression in wild type *A. pseudoterreus* 32359. Thus, inactivation of the tf gene dramatically reduced the level of mRNA of other genes in the cluster. Within the itaconic acid biosynthesis cluster, the transcription factor potentially controls expression of other genes.

Example 6

cadA Deletion Creates a Novel Strain that Produces Aconitic Acid

In *A. pseudoterreus*, when cadA was deleted, itaconic acid production was completely abolished (FIG. 3B). However, 3.5 g per liter aconitic acid in the Δ cadA strain was detected at day 5 (FIG. 6A). Aconitic acid was not produced by the wild type, Δ mttA or Δ mfsA strains (FIG. 6A). A time course analysis showed that aconitic acid started to appear in the supernatant at day 3, similar as IA in the wild type strain (FIG. 6B). At day 3, only cis-acconitic acid was detected in the supernatant. At day 4, both cis-acconitic acid and trans-acconitic acid were detected. From day 5 onward, cis-acconitic acid remained consistent at about 2 g/L, while trans-acconitic acid yield continued to increase (FIG. 6B). By day 10, 10 g/L trans-acconitic acid was detected in the supernatant from the Δ cadA strain (FIG. 6B). FIG. 6C shows a comparison of total aconitic acid production between wild type and Δ cadA fungi. Thus, Δ cadA strains of *A. pseudoterreus* and *A. terreus* can be used to produce cis- and trans-acconitic acid.

Example 7

Materials and Methods

This example describes methods used in the experiments described in Example 8.

Transgene Expression Vector for 3-HP Production

Isolation of DNA Fragments:

Fragment 1: *A. pseudoterreus* 5'-cadA gene, 987 bp (SEQ ID NO: 59) isolated by PCR with the oligo pair 1969 and 1970 (SEQ ID NOS: 60 and 61, respectively) and *A. pseudoterreus* genomic DNA;

Fragment 2: *A. niger* gpdA promoter, 813 bp (SEQ ID NO: 62) isolated by PCR with oligo pair of 1971 and 1972 (SEQ ID NOS: 63 and 64, respectively) and *A. niger* genomic DNA;

Fragment 3: aspartate 1-decarboxylase (panD) cDNA of *Tribolium castaneum* with codon optimization for *A. pseudoterreus*, 1617 bp (SEQ ID NO: 65) was isolated by PCR with the oligo pair of 1973 and 1974 (SEQ ID NOS: 66 and 67, respectively) and the plasmid DNA containing the synthesized panD cDNA;

Fragment 4: bidirectional terminator from *A. niger* elf3/ multifunctional chaperone (SEQ ID NO: 68) was isolated by PCR with oligo pair of 1975 and 1976 (SEQ ID NOS: 69 and 70, respectively) and the genomic DNA of *A. niger*;

Fragment 5: codon optimized synthetic cDNA of β -alanine-pyruvate aminotransferase (BAPAT) of *Bacillus cereus*, 1350 bp (SEQ ID NO: 71) was isolated by PCR with oligo pair of 1977 and 1978 (SEQ ID NOS: 72 and 73, respectively) and the plasmid DNA containing the synthesized BABAT cDNA;

Fragment 6: *A. niger* enol promoter, 704 bp (SEQ ID NO: 74) isolated by PCR with oligo pair of 1979 and 1980 (SEQ ID NOS: 75 and 76, respectively) and *A. niger* genomic DNA;

Fragment 7: *A. nidulans* gpdA promoter, 885 bp (SEQ ID NO: 77) was isolated by PCR with the oligo pair of 2002 and 1982 (SEQ ID NOS: 78 and 79, respectively) and *A. nidulans* genomic DNA;

Fragment 8: the codon optimized synthetic cDNA of *E. coli* 3-hydroxypropionate dehydrogenase (HPDH), 741 bp (SEQ ID NO: 80) was isolated by PCR with oligo pair of 1983 and 1984 (SEQ ID NOS: 81 and 82, respectively) and the plasmid DNA containing the codon-optimized synthesized HPDH DNA of *E. coli*;

Fragment 9: trpC terminator of *A. nidulans*, 473 bp (SEQ ID NO: 83) isolated by PCR with oligo pair of 1985 and 2004 (SEQ ID NOS: 84 and 85, respectively) and plasmid DNA of pAN7.1;

Fragment 10: trpC terminator of *A. nidulans*, 473 bp (SEQ ID NO: 86) isolated by PCR with the oligo pair of 2005 and 1986 (SEQ ID NOS: 87 and 88, respectively) and plasmid DNA of pAN7.1;

Fragment 11: *A. oryzae* ptrA selection marker gene, 2005 bp; SEQ ID NO: 89) isolated by PCR with the oligo pair of 1987 and 1988 (SEQ ID NOS: 90 and 91, respectively) and *A. oryzae* genomic DNA;

Fragment 12: *A. pseudoterreus* 3'-cadA gene, 908 bp (SEQ ID NO: 92) isolated by PCR with the oligo pair 1989 and 2003 (SEQ ID NOS: 93 and 94, respectively) and *A. oryzae* genomic DNA;

Fragment 13 (SEQ ID NO: 95): Combination of Fragments 7 to 9 (SEQ ID NOS: 77, 80, and 83, respectively), 2099 bp isolated by PCR with oligo pair of 1981 and 1986 (SEQ ID NOS: 96 and 88, respectively) and plasmid DNA of pZD-2; and

Fragment 14 (SEQ ID NO: 97): Combination of Fragments 11 to 12 (SEQ ID NOS: 89 and 92, respectively), 2913 bp was isolated by PCR with the oligo pair of 1987 and 1990 (SEQ ID NOS: 90 and 98, respectively) and plasmid DNA of pZD-3.

The oligonucleotide primers used are shown in Table 4.

TABLE 4

Primers used to generate vector for 3-HP production	
Name	Sequence (SEQ ID NO:)
1969cad1	ccctcgaggctgacggatcgataGATATCGGTTGTAGCAGCGTAAACAC (60)
1970cad2	tctttcatagtagCCTTGGTGAACATCTTGAGG (61)
1971gpdA1	atgttcaccaaggCTACTATGAAAGACCGCGATG (63)
1972gpdA2	cgccggtggcgggCATTGTTTGTAGATGTCTATGTG (64)
1973pan1	catctaaacaatgCCCCACCACCGCGAGGACCA (66)
1974pan2	atccaaccatcaGAGGTCGGAGCCAGGCGTTCG (67)
1975ter1	gggctccgacctGTGATGGGTTGGATGACGATG (69)
1976ter2	tctggcccagctcTGAGTCTAGATGGGTGGTG (70)
1977bap1	catctaggactcaGAGCTGGGCCAGACATTCCTTC (72)
1978bap2	gtccatcaacatgGAACTGATGATCGTCCAGGTCAC (73)
1979eno1	cgatcatcagttcCATGTTGATGGACTGGAGGG (75)
1980eno2	gaactagtgatcccccggtgcGttaaCTCGAGCTTAC AAGAGTAGCC (76)

TABLE 4-continued

Primers used to generate vector for 3-HP production	
Name	Sequence (SEQ ID NO:)
1981gpdA1	acaggctacttcttctaagctcgagttTCTGTACAGTGAC CGGTGAC (96)
1982gpdA2	tgaccagcagcatCATGGTGATGTCTGCTCAAG (79)
1983hpd1	agacatcaccatgATCGTCTGGTCACGGGCGC (81)
1984hpd2	gccatcggtcctaTTGGCGGTGGACGTTTCAGGC (82)
1985trp1	cgccccaccgcaaTAGGACCGATGGCTGTGTAG (84)
1986trp2	cccgtctgtcagaGAGCGGATTCCTCAGTCTCG (88)
1987ptrA1	gaggaatccgctcTCTGACAGACGGCAATTGATTAC (90)
1988ptrA2	gaatggtctgagGAGCCGCTCTTGCATCTTTG (91)
1989cad3	gcaagagcggctcCTCAGCAACATTCGCCATGTTC (93)
1990cad4	actaaagggaaacaaagctggagctCAGCTCCACTGCAT AGTCTTTG(98)
2002gpdA5	ccctcgaggtcgacggatcgataGTTAACTCTGTACAGTG ACCGGTGAC (78)
2003cad3	gaactagtggatccccgggctgcaCAGCTCCACTGCAT AGTCTTTG (94)
2004trpR	gaactagtggatccccgggctgcaGAGCGGATTCCTCAGT CTCG (85)
2005trpF	ccctcgaggtcgacggatcgataTAGGACCGATGGCTGTG TAG (87)

An overview of the arrangement of the Fragments is shown in FIG. 7. Fragments 1 to 6 (SEQ ID NOS: 59, 62, 65, 68, 71 and 74, respectively) were assembled into the plasmid DNA pBlueScript SK (-) linearized with HindIII and PstI via Gibson Assembly master kit to form plasmid pZD-1. A restriction enzyme site HpaI was introduced at the end of the fragment 6 for further cloning.

Fragments 7 to 9 (SEQ ID NOS: 77, 80, and 83, respectively) were assembled into the plasmid DNA pBlueScript SK (-) linearized with HindIII and PstI via Gibson Assembly master kit to form plasmid DNA pZD-2.

Fragments 10 to 12 (SEQ ID NOS: 86, 89, and 92, respectively) were assembled into the pBlueScript SK(-) vector linearized with restriction enzyme HindIII and PstI by

Gibson assembly to form the plasmid vector ZD-3. (Only fragments 11 and 12 were used in the next step; SEQ ID NOS: 89 and 92).

Fragments 13 and 14 (SEQ ID NOS: 95 and 97) were assembled together into the plasmid DNA vector ZD-1 linearized with restriction enzyme HpaI/SacI via Gibson Assembly master kit to form pZD-4.

Genomic DNA isolation and Southern blotting analysis were performed as described in Example 1 (and see Dai et al., 2017, *Appl Microbiol Biotechnol* 101:6099-6110).

Detection of 3-HP

The extracellular 3-HP in the culture supernatants was quantified with HPLC method as described in Example 1.

Example 8

Production of 3-HP

The constructs generated in Example 7 (FIG. 7) were transformed into wild type *A. pseudoterreus* strain ATCC 32359 using the methods describe in Example 1, thereby inactivating/disrupting the *cadA* gene in some examples.

As shown in FIG. 8, restriction fragment length polymorphism of selected transgenic strains show that the transgene expression cassette was inserted into the *cadA* locus in strain-2 (with one copy) and strain-6 (two copies), while the strain-4 and strain-5 carry the transgene expression cassette with random integration. No integration of transgene expression cassette was observed in strain-1 and strain-3.

3-HP production was measured in several transformants. As shown in FIG. 9A, the $\Delta cadA$ strain did not produce 3-HP, while insertion of the transgene expression cassette that allowed for expression of panD, BAPAT, and HPDH, into the *cadA* locus with one copy or two copies and resulted in 0.9 or 1.7 g/13-HP accumulation in the strains 3HP-2 or 3HP-6. In contrast, when the transgene expression cassette was randomly inserted into the chromosome, 3HP production was substantially lower (Strains 3HP-4 and 3HP-5). FIG. 9B shows 3-HP production over 8 days in Strains 3HP-2 and 3HP-6 (strains 2 and 6, respectively). Thus, genetically inactivating *cadA* can increase 3-HP production.

In view of the many possible embodiments to which the principles of the disclosure may be applied, it should be recognized that the illustrated embodiments are only examples of the invention and should not be taken as limiting the scope of the invention. Rather, the scope of the invention is defined by the following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 98
 <210> SEQ ID NO 1
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete *tf* gene in *A. pseudoterreus*

<400> SEQUENCE: 1

gagccatagc catgcaagcg

<210> SEQ ID NO 2

-continued

<211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete tf gene in A.
 pseudoterreus

 <400> SEQUENCE: 2

 atagagtcct tggatgagac g 21

 <210> SEQ ID NO 3
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete tf gene in A.
 pseudoterreus

 <400> SEQUENCE: 3

 gtggatttcg aggttccttg c 21

 <210> SEQ ID NO 4
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete tf gene in A.
 pseudoterreus

 <400> SEQUENCE: 4

 gaagtagaac catgtggatc g 21

 <210> SEQ ID NO 5
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete tf gene in A.
 pseudoterreus

 <400> SEQUENCE: 5

 tgacctccac tagctccagc actactagat aggcccgttt agagagtgcc 50

 <210> SEQ ID NO 6
 <211> LENGTH: 49
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete tf gene in A.
 pseudoterreus

 <400> SEQUENCE: 6

 aatagagtag atgccgaccg gccgcttcga cgacagctct gcactctcc 49

 <210> SEQ ID NO 7
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete tf gene in A.
 pseudoterreus

 <400> SEQUENCE: 7

 ggcactctct aaacgggctc atctagtagt gctggageta gtggagggtca 50

 <210> SEQ ID NO 8
 <211> LENGTH: 49

-continued

<212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete tf gene in A.
 pseudoterreus

 <400> SEQUENCE: 8

 ggagagtgca gagctgtcgt cgaagcggcc ggtcggcatc tactctatt 49

 <210> SEQ ID NO 9
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete mttA gene in A.
 pseudoterreus

 <400> SEQUENCE: 9

 gctgcatact cggattacgc 20

 <210> SEQ ID NO 10
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete mttA gene in A.
 pseudoterreus

 <400> SEQUENCE: 10

 gaaaaggtac tggagtacg 20

 <210> SEQ ID NO 11
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete mttA gene in A.
 pseudoterreus

 <400> SEQUENCE: 11

 cagaccaagg agctttcctg 20

 <210> SEQ ID NO 12
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete mttA gene in A.
 pseudoterreus

 <400> SEQUENCE: 12

 cattaagcca caggcttgcg 20

 <210> SEQ ID NO 13
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete mttA gene in A.
 pseudoterreus

 <400> SEQUENCE: 13

 tgacctccac tagctccagc aatatggatg ctgttcgttc gccgtgctgg 50

 <210> SEQ ID NO 14
 <211> LENGTH: 50
 <212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete mttA gene in A.
 pseudoterreus

 <400> SEQUENCE: 14

 aatagagtag atgccgaccg tgaccgaggat gtgctgagtc caaacaagaagc 50

 <210> SEQ ID NO 15
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete mttA gene in A.
 pseudoterreus

 <400> SEQUENCE: 15

 ccagcacggc gaacgaacag catccatatt gctggagcta gtggaggctca 50

 <210> SEQ ID NO 16
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete mttA gene in A.
 pseudoterreus

 <400> SEQUENCE: 16

 gctttgtttg gactcagcac atcctcgtca cggtcggcat ctactctatt 50

 <210> SEQ ID NO 17
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete cadA gene in A.
 pseudoterreus

 <400> SEQUENCE: 17

 ctccagtaac agaaccgacc 20

 <210> SEQ ID NO 18
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete cadA gene in A.
 pseudoterreus

 <400> SEQUENCE: 18

 gaacttcact gccgcattgg 20

 <210> SEQ ID NO 19
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete cadA gene in A.
 pseudoterreus

 <400> SEQUENCE: 19

 ggacactcca agaggataag g 21

 <210> SEQ ID NO 20
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete cadA gene in A.
 pseudoterreus

<400> SEQUENCE: 20

gctcatcaca ttgtttgccc 20

<210> SEQ ID NO 21
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete cadA gene in A.
 pseudoterreus

<400> SEQUENCE: 21

tgacctccac tagctccagc ggtcaattta agaggacgat cttegctgcg 50

<210> SEQ ID NO 22
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete cadA gene in A.
 pseudoterreus

<400> SEQUENCE: 22

aatagagtag atgccgaccg tcagcctgga caggctcacc gacattagcc 50

<210> SEQ ID NO 23
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete cadA gene in A.
 pseudoterreus

<400> SEQUENCE: 23

cgcagcgaag atcgtcctct taaattgacc gctggagcta gtggagggtca 50

<210> SEQ ID NO 24
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete cadA gene in A.
 pseudoterreus

<400> SEQUENCE: 24

ggctaagtgc ggtgagcctg tccaggctga cggtcggcat ctactctatt 50

<210> SEQ ID NO 25
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete mfsA gene in A.
 pseudoterreus

<400> SEQUENCE: 25

tgatgagctg aattcgttgc 20

<210> SEQ ID NO 26
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 26

tatagccagc ttttgctgtg 20

<210> SEQ ID NO 27
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 27

catagcggtc agagtgttg 19

<210> SEQ ID NO 28
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 28

ccatttcaat gctttgtgcg 20

<210> SEQ ID NO 29
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 29

ccataccacc cttaccctct tggagtgtcc gctggageta gtggaggta 50

<210> SEQ ID NO 30
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 30

gctgtggcct cctggcgatt acgcaatatt cggtcggcat ctactctatt 50

<210> SEQ ID NO 31
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.
pseudoterreus

<400> SEQUENCE: 31

tgacctccac tagctccagc ggacactcca agagggtaag ggtggatg 50

<210> SEQ ID NO 32
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to delete mfsA gene in A.

-continued

pseudoterreus

<400> SEQUENCE: 32

aatagagtag atgccgaccg aatattgcgt aatgccagg aggccacagc 50

<210> SEQ ID NO 33
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete p450 gene in A.
 pseudoterreus

<400> SEQUENCE: 33

tctccaaatc atcgatcatcg 20

<210> SEQ ID NO 34
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete p450 gene in A.
 pseudoterreus

<400> SEQUENCE: 34

cttcaatcgc accgacatcc 20

<210> SEQ ID NO 35
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete p450 gene in A.
 pseudoterreus

<400> SEQUENCE: 35

tcgtgtagac aagtccagtc 20

<210> SEQ ID NO 36
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete p450 gene in A.
 pseudoterreus

<400> SEQUENCE: 36

ctataccact ctagtgatgg 20

<210> SEQ ID NO 37
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete p450 gene in A.
 pseudoterreus

<400> SEQUENCE: 37

cctctgctca ggttgttttc gaacaggagc gctggagcta gtggagggtca 50

<210> SEQ ID NO 38
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete p450 gene in A.
 pseudoterreus

-continued

<400> SEQUENCE: 38
 cggaatgcag ataggcatca cagtccagaa cggtcggcat ctactctatt 50

<210> SEQ ID NO 39
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete p450 gene in A.
 pseudoterreus

<400> SEQUENCE: 39
 tgacctccac tagctccagc gctcctgttc gaaaacaacc tgagcagagg 50

<210> SEQ ID NO 40
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to delete p450 gene in A.
 pseudoterreus

<400> SEQUENCE: 40
 aatagagtag atgccgaccg ttctggactg tgatgcctat ctgcattccg 50

<210> SEQ ID NO 41
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to amplify mttA in A.
 pseudoterreus

<400> SEQUENCE: 41
 gctttcaatg tggttcctac 20

<210> SEQ ID NO 42
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to amplify mttA in A.
 pseudoterreus

<400> SEQUENCE: 42
 ctccatcacc taccctttc 19

<210> SEQ ID NO 43
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to amplify cadA in A.
 pseudoterreus

<400> SEQUENCE: 43
 gaagtgtggg atctggc 17

<210> SEQ ID NO 44
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to amplify cadA in A.
 pseudoterreus

-continued

<400> SEQUENCE: 44
 gggttcggta tttgtgaag 19

<210> SEQ ID NO 45
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to amplify mfsA in A.
 pseudoterreus

<400> SEQUENCE: 45
 caagaacagt ttggcctgag 20

<210> SEQ ID NO 46
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to amplify mfsA in A.
 pseudoterreus

<400> SEQUENCE: 46
 gcggacatca tacaatctgg 20

<210> SEQ ID NO 47
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to amplify beta-tubulin in A.
 pseudoterreus

<400> SEQUENCE: 47
 ttgtcgatgt tgttcgtcgc 20

<210> SEQ ID NO 48
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to amplify beta-tubulin in A.
 pseudoterreus

<400> SEQUENCE: 48
 tggcgttgta aggctcaacc 20

<210> SEQ ID NO 49
 <211> LENGTH: 2206
 <212> TYPE: DNA
 <213> ORGANISM: Aspergillus terreus

<400> SEQUENCE: 49
 gtgggtcttg aaatcgtatg ccacacttgc tccggatgaa acacattccg gagcgcgcat 60
 cgatattgct acacagtata gacccaatgg tctgcagatg ccctaaatgg tagttctcac 120
 tggcctgcat taagtcttgg ttgcagatca ttgtcggcct aacatcagtg taggttacgg 180
 tgtgagatth acttgcatag aagattocag accacaaggt tctagatcct ttgacggcgg 240
 actcccctcg aggtgcceggg cgccgacgtg tgcgttgctc cgggatttgt aggacgcagc 300
 tcggatacct agccggtatg ggaatcggag gttgtagcag cgtaaacaca tggatagtta 360
 aataatcgga tgtacacca ctgttgaaa tgacgggggc ctacaacacg agattatctg 420
 atccaatttc tgttcgttgg cattctatca ttcgcagcga aaattgtcct attaaattga 480

-continued

```

ccatgaccaa acaatctgcg gacagcaacg caaagtcagg agttacgtcc gaaatatgtc 540
attgggcatc caacctggcc actgacgaca tcccttcgga cgtattagaa agagcaaaat 600
accttattct cgacgggtatt gcatgtgcct gggttgggtc aagagtgcct tggtcagaga 660
agtatgttca ggcaacgatg agctttgagc cgccgggggc ctgcagggtg attggatatg 720
gacaggtaaa ttttattcac tctagacggt ccacaaagta tactgacgat ccttcgtata 780
gaaactgggg cctgttgagc cagccatgac caattccgct ttcatacagg ctacggagct 840
tgacgactac cacagcgaag cccccctaca ctctgcaagc attgtccttc ctgcggtctt 900
tgcagcaagt gaggtcttag ccgagcaggg caaaacaatt tccggtatag atgttattct 960
agccgccatt gtgggggttg aatctggccc acggatcggc aaagcaatct acggatcggg 1020
cctcttgaac aacggctggc attgtggagc tgtgtatggc gctccagccg gtgctgctggc 1080
cacaggaaa gctcctcggtc taactccaga ctccatggaa gatgctctcg gaattgcgtg 1140
cacgcaagcc tgtggtttaa tgcgcgca atacggaggc atggtaaagc gtgtgcaaca 1200
cggattcgca gcgcgtaatg gtctcttgg gggactggtg gcccatgggtg ggtacgagcc 1260
aatgaaaggt gtcttgagga gatcttacgg cggttctctc aagatgttca ccaagggcaa 1320
cggcagagag cctccctaca aagaggagga agtgggtggc ggtctcgggt cattctggca 1380
tacctttact attcgcatac agctctatgc ctgctgcgga cttgtccatg gtccagtcga 1440
ggctatcgaa aaccttcagg ggagatacc cagactcttg aatagagcca acctcagcaa 1500
cattcgccat gttcatgtac agctttcaac ggcctcgaac agtcaactgtg gatggatacc 1560
agaggagaga cccatcagtt caatcgcagg gcagatgagt gtcgcataca ttctcgcgt 1620
ccagctggtc gaccagcaat gtcttttgtc ccagttttct gagtttgatg acaacctgga 1680
gaggccagaa gtttgggatc tggccaggaa ggttacttca tctcaaagcg aagagtttga 1740
tcaagacggc aactgtctca gtgcgggtcg cgtgaggatt gagttcaacg atggttcttc 1800
tattacggaa agtgtcgaga agcctcttgg tgtcaaagag cccatgccaa acgaacggat 1860
tctccacaaa taccgaaccc ttgctggtag cgtgacggac gaatcccggg tgaagagat 1920
tgaggatctt gtctcggcc tggacaggct caccgacatt agcccattgc tggagctgct 1980
gaattgcccc gtgaaatgc cactggtata aatgggaagc gatatggaaa catttcatgt 2040
cacgggcaca aattctaggt catatcgtac ctggatggtg aaaccaccag cggtttagca 2100
gatagaagat agactccttc tgetctgctg tgcgtcttga atttagttcg ttcactggtc 2160
taagaactta gaatgcaata cagtctctct tatttcttat taaaat 2206

```

```

<210> SEQ ID NO 50
<211> LENGTH: 490
<212> TYPE: PRT
<213> ORGANISM: Aspergillus terreus
<400> SEQUENCE: 50

```

```

Met Thr Lys Gln Ser Ala Asp Ser Asn Ala Lys Ser Gly Val Thr Ser
 1             5             10             15
Glu Ile Cys His Trp Ala Ser Asn Leu Ala Thr Asp Asp Ile Pro Ser
          20             25             30
Asp Val Leu Glu Arg Ala Lys Tyr Leu Ile Leu Asp Gly Ile Ala Cys
          35             40             45
Ala Trp Val Gly Ala Arg Val Pro Trp Ser Glu Lys Tyr Val Gln Ala
 50             55             60

```

-continued

Thr Met Ser Phe Glu Pro Pro Gly Ala Cys Arg Val Ile Gly Tyr Gly
 65 70 75 80
 Gln Lys Leu Gly Pro Val Ala Ala Ala Met Thr Asn Ser Ala Phe Ile
 85 90 95
 Gln Ala Thr Glu Leu Asp Asp Tyr His Ser Glu Ala Pro Leu His Ser
 100 105 110
 Ala Ser Ile Val Leu Pro Ala Val Phe Ala Ala Ser Glu Val Leu Ala
 115 120 125
 Glu Gln Gly Lys Thr Ile Ser Gly Ile Asp Val Ile Leu Ala Ala Ile
 130 135 140
 Val Gly Phe Glu Ser Gly Pro Arg Ile Gly Lys Ala Ile Tyr Gly Ser
 145 150 155 160
 Asp Leu Leu Asn Asn Gly Trp His Cys Gly Ala Val Tyr Gly Ala Pro
 165 170 175
 Ala Gly Ala Leu Ala Thr Gly Lys Leu Leu Gly Leu Thr Pro Asp Ser
 180 185 190
 Met Glu Asp Ala Leu Gly Ile Ala Cys Thr Gln Ala Cys Gly Leu Met
 195 200 205
 Ser Ala Gln Tyr Gly Gly Met Val Lys Arg Val Gln His Gly Phe Ala
 210 215 220
 Ala Arg Asn Gly Leu Leu Gly Gly Leu Leu Ala His Gly Gly Tyr Glu
 225 230 235 240
 Ala Met Lys Gly Val Leu Glu Arg Ser Tyr Gly Gly Phe Leu Lys Met
 245 250 255
 Phe Thr Lys Gly Asn Gly Arg Glu Pro Pro Tyr Lys Glu Glu Glu Val
 260 265 270
 Val Ala Gly Leu Gly Ser Phe Trp His Thr Phe Thr Ile Arg Ile Lys
 275 280 285
 Leu Tyr Ala Cys Cys Gly Leu Val His Gly Pro Val Glu Ala Ile Glu
 290 295 300
 Asn Leu Gln Gly Arg Tyr Pro Glu Leu Leu Asn Arg Ala Asn Leu Ser
 305 310 315 320
 Asn Ile Arg His Val His Val Gln Leu Ser Thr Ala Ser Asn Ser His
 325 330 335
 Cys Gly Trp Ile Pro Glu Glu Arg Pro Ile Ser Ser Ile Ala Gly Gln
 340 345 350
 Met Ser Val Ala Tyr Ile Leu Ala Val Gln Leu Val Asp Gln Gln Cys
 355 360 365
 Leu Leu Ser Gln Phe Ser Glu Phe Asp Asp Asn Leu Glu Arg Pro Glu
 370 375 380
 Val Trp Asp Leu Ala Arg Lys Val Thr Ser Ser Gln Ser Glu Glu Phe
 385 390 395 400
 Asp Gln Asp Gly Asn Cys Leu Ser Ala Gly Arg Val Arg Ile Glu Phe
 405 410 415
 Asn Asp Gly Ser Ser Ile Thr Glu Ser Val Glu Lys Pro Leu Gly Val
 420 425 430
 Lys Glu Pro Met Pro Asn Glu Arg Ile Leu His Lys Tyr Arg Thr Leu
 435 440 445
 Ala Gly Ser Val Thr Asp Glu Ser Arg Val Lys Glu Ile Glu Asp Leu
 450 455 460
 Val Leu Gly Leu Asp Arg Leu Thr Asp Ile Ser Pro Leu Leu Glu Leu
 465 470 475 480
 Leu Asn Cys Pro Val Lys Ser Pro Leu Val

-continued

485

490

<210> SEQ ID NO 51
 <211> LENGTH: 1862
 <212> TYPE: DNA
 <213> ORGANISM: *Aspergillus vadensis*

<400> SEQUENCE: 51

```

gttttctgtg tgtctttggg gggttataaa taggggtgctg aatatctgga agatagggaa    60
ttctttctct ttcaatcaat caatcaagaa ttcttttagg gagtttctat actacatccg    120
atatggctgc catcaccgct aaatctgaag cggcttctgc tacttcgccc attcctacca    180
attctaatac taccatgact actaccctca acggggtaga tggttcaaaa gagaaagaaa    240
aagaccagat acccccaaac aaagaggaag gaacaaaagc agaagagaaa gaaaccgaag    300
catacaactc ctccaacggc gtcaccagcc aactctgcaa ctggatcgcc tctctccagc    360
tagaagacat tccagactct gtcgcacccc gcgccaagta cctctttctc gatggcatcg    420
cctgcgcact cgtcgggtgc cgcgtcccat ggtcgcagaa ggcttttgat gcgatggctg    480
ttttcgagga gaagggaaag catgtggtta ttgggtatga agagcgcctt ggtgctatcg    540
ccgcgcgaac cctcaacggc tctcggatcc aagcctgcga agtagacgac taccacagcg    600
tggcgccctt gactcgcag gccgtggtea tccctcctct cttcgctgcc gccgtcagtg    660
cgcggaacca tccgaccgca ccgcgcatca tcgacgggag aacacttctt ctgcctccg    720
tggtaggggt cgaggttggt ccgcgcgtgg gcatggcgtt acacggcacc gagatgctcg    780
cgaagggatg gactcgcggg tctgtgtttg gtggaccgcg gcccgaggc agttctgcaa    840
aactactcgg tttgtcggcg ggtcaagtgc aagacgcgat cggagtagca gcgacacaag    900
catgcggact catggcggcg cagtacgacg ggatggtgaa gcggatgcat catggcttcg    960
cggcaaggaa tggactgttg ggcacgatgt tagcgtgggg aggttatgaa gggatcaaga   1020
agggtgttga gcggccgctat ggaggatttc tggcaatggt tggcctaggg tcgaagcaca   1080
cgcctagttc gaagccggag gaggtggcaa aggatttggg gacgttcttg cacacggcgg   1140
agtggattcg gttgaagtgt catgcgtgct gtggggggat tcatggcacc attgagtgtt   1200
tggcggagat gcaggagatg tatccagacg gatttggacg ggagaaacta ggagagatca   1260
aggagattcg gatccagttg agtcatgctg tgtttcatca ttgtggatgg gcgccggaga   1320
cgaggccgtt gaccccgacg ggggcgcaga tgaatacggc gtttgtggcg gcctcgcagt   1380
tgggtggatg acaagtgttg ttggagcagt tctcgtcggg gaagttggat cgggatgagg   1440
tttgggaatt gattgggaag acgagttgta ttcatacggc ggagttggac aagccgaata   1500
ttggttggg tgcgttgatc tccatcacgt ttgcggatgg cagtcagggt cagcattcgt   1560
tgttgaagcc gaaggggggtg gatgaacca tttcgaatga ggagatcttg gagaagtttc   1620
gtcggttgac gggcgggttg attggggtgg agaggcagga gaagattgaa aaggccgtgc   1680
tggggatgga ggagttgcag gatgtggatg agttgattga gttgctgagt gtgaatgtgg   1740
tcaatccgtt gcagtagtat actagtoatc tgttttgatg cttctggcgt tggctcgtgtt   1800
gggatagtat ctcataatct tgaattaata aatcattcaa catggtgaaa atcatatctg   1860
tg

```

<210> SEQ ID NO 52
 <211> LENGTH: 544
 <212> TYPE: PRT
 <213> ORGANISM: *Aspergillus vadensis*

-continued

<400> SEQUENCE: 52

Met Val Ala Ile Thr Ala Lys Ser Glu Ala Ala Ser Ala Thr Ser Pro
1 5 10 15
Ile Pro Thr Asn Ser Asn Thr Thr Met Thr Thr Thr Leu Asn Gly Val
20 25 30
Asp Gly Ser Lys Glu Lys Glu Lys Asp Gln Ile Pro Pro Asn Lys Glu
35 40 45
Glu Gly Thr Lys Ala Glu Glu Lys Glu Thr Glu Ala Tyr Asn Ser Ser
50 55 60
Asn Gly Val Thr Ser Gln Leu Cys Asn Trp Ile Ala Ser Leu Gln Leu
65 70 75 80
Glu Asp Ile Pro Asp Ser Val Arg Thr Arg Ala Lys Tyr Leu Phe Leu
85 90 95
Asp Gly Ile Ala Cys Ala Leu Val Gly Ala Arg Val Pro Trp Ser Gln
100 105 110
Lys Ala Phe Asp Ala Met Ala Val Phe Glu Glu Lys Gly Lys His Val
115 120 125
Val Ile Gly Tyr Glu Glu Arg Leu Gly Ala Ile Ala Ala Thr Leu
130 135 140
Asn Gly Ser Trp Ile Gln Ala Cys Glu Val Asp Asp Tyr His Ser Val
145 150 155 160
Ala Pro Leu His Ser Gln Ala Val Val Ile Pro Pro Leu Phe Ala Ala
165 170 175
Ala Val Ser Ala Arg Asn His Pro Thr Ala Pro Arg Ile Ile Asp Gly
180 185 190
Arg Thr Leu Leu Leu Ala Ser Val Val Gly Phe Glu Val Gly Pro Arg
195 200 205
Val Gly Met Ala Leu His Gly Thr Glu Met Leu Ala Lys Gly Trp His
210 215 220
Cys Gly Ser Val Phe Gly Gly Pro Ala Ala Ala Gly Ser Ser Ala Lys
225 230 235 240
Leu Leu Gly Leu Ser Ala Gly Gln Val Glu Asp Ala Ile Gly Val Ala
245 250 255
Ala Thr Gln Ala Cys Gly Leu Met Ala Ala Gln Tyr Asp Gly Met Val
260 265 270
Lys Arg Met His His Gly Phe Ala Ala Arg Asn Gly Leu Leu Gly Thr
275 280 285
Met Leu Ala Trp Gly Gly Tyr Glu Gly Ile Lys Lys Val Phe Glu Arg
290 295 300
Pro Tyr Gly Gly Phe Leu Ala Met Phe Gly Leu Gly Ser Lys His Thr
305 310 315 320
Pro Ser Ser Lys Pro Glu Glu Val Ala Lys Asp Leu Gly Thr Phe Trp
325 330 335
His Thr Ala Glu Trp Ile Arg Leu Lys Leu His Ala Cys Cys Gly Gly
340 345 350
Ile His Gly Thr Ile Glu Cys Leu Ala Glu Met Gln Glu Met Tyr Pro
355 360 365
Glu Arg Phe Gly Arg Glu Lys Leu Gly Glu Ile Lys Glu Ile Arg Ile
370 375 380
Gln Leu Ser Asp Ala Val Phe His His Cys Gly Trp Ala Pro Glu Thr
385 390 395 400
Arg Pro Leu Thr Pro Thr Gly Ala Gln Met Asn Thr Ala Phe Val Ala

-continued

	405		410		415
Ala Ser Gln Leu Val Asp Gly Gln Val Leu Leu Glu Gln Phe Ser Ser	420		425		430
Gly Lys Leu Asp Arg Asp Glu Val Trp Glu Leu Ile Gly Lys Thr Ser	435		440		445
Cys Ile His Thr Ala Glu Leu Asp Lys Pro Asn Ile Gly Cys Gly Ala	450		455		460
Leu Ile Ser Ile Thr Phe Ala Asp Gly Ser Gln Val Gln His Ser Leu	465		470		475
Leu Lys Pro Lys Gly Val Asp Glu Pro Ile Ser Asn Glu Glu Ile Leu	485		490		495
Glu Lys Phe Arg Arg Leu Thr Gly Gly Leu Ile Gly Val Glu Arg Gln	500		505		510
Glu Lys Ile Glu Lys Ala Val Leu Gly Met Glu Glu Leu Gln Asp Val	515		520		525
Asp Glu Leu Ile Glu Leu Leu Ser Val Asn Val Val Asn Pro Leu Gln	530		535		540

<210> SEQ ID NO 53
 <211> LENGTH: 1797
 <212> TYPE: DNA
 <213> ORGANISM: Tribolium castaneum
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (41)..(1663)

<400> SEQUENCE: 53

acttgtgaat cagtcgtgcc cccacgagga tccacacacg atg ccg gcc aca ggc	55
Met Pro Ala Thr Gly	
1 5	
gaa gac caa gac ctg gtg caa gac ctc atc gag gag ccc gcc acc ttc	103
Glu Asp Gln Asp Leu Val Gln Asp Leu Ile Glu Glu Pro Ala Thr Phe	
10 15 20	
agc gac gcc gtc ctc tcc tcc gac gag gaa ctc ttc cac cag aag tgc	151
Ser Asp Ala Val Leu Ser Ser Asp Glu Glu Leu Phe His Gln Lys Cys	
25 30 35	
ccc aaa ccc gcc ccc att tac tcc ccg gtc tcg aaa ccg gtc tcc ttc	199
Pro Lys Pro Ala Pro Ile Tyr Ser Pro Val Ser Lys Pro Val Ser Phe	
40 45 50	
gag agc ctc ccc aac agg cgc ctc cac gag gag ttc ctc cgc agc tcg	247
Glu Ser Leu Pro Asn Arg Arg Leu His Glu Glu Phe Leu Arg Ser Ser	
55 60 65	
gtg gac gtc ctc ctc cag gag gcg gtg ttc gag gga acg aac cgc aag	295
Val Asp Val Leu Leu Gln Glu Ala Val Phe Glu Gly Thr Asn Arg Lys	
70 75 80 85	
aac cgg gtg ctg caa tgg cgg gag ccg gag gag ttg agg cgt ctg atg	343
Asn Arg Val Leu Gln Trp Arg Glu Pro Glu Glu Leu Arg Arg Leu Met	
90 95 100	
gac ttt ggg gtg cgg agt gcg ccc tcc acg cac gag gag ttg ttg gag	391
Asp Phe Gly Val Arg Ser Ala Pro Ser Thr His Glu Glu Leu Leu Glu	
105 110 115	
gtg ttg aag aag gtt gta act tat tcg gtt aaa acc gga cat ccg tac	439
Val Leu Lys Lys Val Val Thr Ser Val Lys Thr Gly His Pro Tyr	
120 125 130	
ttc gtg aac cag ttg ttc tcg gcg gtg gat ccg tac ggt ttg gtg gca	487
Phe Val Asn Gln Leu Phe Ser Ala Val Asp Pro Tyr Gly Leu Val Ala	
135 140 145	
caa tgg gcc acg gat gcg ctc aat ccg agt gtt tac acc tac gag gtt	535
Gln Trp Ala Thr Asp Ala Leu Asn Pro Ser Val Tyr Thr Tyr Glu Val	

-continued

150	155	160	165	
tcg ccg gtt ttt gtt ctg atg gag gaa gtg gtt ttg agg gag atg agg Ser Pro Val Phe Val Leu Met Glu Glu Val Val Leu Arg Glu Met Arg 170 175 180				583
gcc att gtg ggg ttc gag ggg gga aag ggc gat ggg att ttt tgc cca Ala Ile Val Gly Phe Glu Gly Gly Lys Gly Asp Gly Ile Phe Cys Pro 185 190 195				631
gga ggg tcc att gcc aat gga tat gcc atc agt tgt gcc aga tac agg Gly Gly Ser Ile Ala Asn Gly Tyr Ala Ile Ser Cys Ala Arg Tyr Arg 200 205 210				679
ttt atg ccc gat att aag aaa aaa ggc ctc cac tct ctc ccc cgt ttg Phe Met Pro Asp Ile Lys Lys Lys Gly Leu His Ser Leu Pro Arg Leu 215 220 225				727
gtc ctc ttc acc tct gaa gat gcc cac tat tcc atc aaa aaa ctc gcc Val Leu Phe Thr Ser Glu Asp Ala His Tyr Ser Ile Lys Lys Leu Ala 230 235 240 245				775
tct ttc caa ggc atc ggc acc gac aac gtc tac ttg ata cga acg gac Ser Phe Gln Gly Ile Gly Thr Asp Asn Val Tyr Leu Ile Arg Thr Asp 250 255 260				823
gcc cga ggt cgc atg gac gtc tcg cac ctg gtg gag gaa atc gag cgt Ala Arg Gly Arg Met Asp Val Ser His Leu Val Glu Glu Ile Glu Arg 265 270 275				871
tcg ctc cgt gaa ggc gcc gct cct ttc atg gtc agt gcc acc gct gga Ser Leu Arg Glu Gly Ala Ala Pro Phe Met Val Ser Ala Thr Ala Gly 280 285 290				919
acc aca gtg att ggt gcc ttt gac ccc atc gaa aaa atc gca gat gtg Thr Thr Val Ile Gly Ala Phe Asp Pro Ile Glu Lys Ile Ala Asp Val 295 300 305				967
tgc caa aaa tac aaa ctg tgg ttg cac gtg gat gcc gcc tgg gga ggt Cys Gln Lys Tyr Lys Leu Trp Leu His Val Asp Ala Ala Trp Gly Gly 310 315 320 325				1015
ggc gcg ctt gtc tct gcc aaa cac cgc cac ctc ctc aaa ggg att gag Gly Ala Leu Val Ser Ala Lys His Arg His Leu Leu Lys Gly Ile Glu 330 335 340				1063
agg gcc gac tcg gtc acc tgg aac cct cac aaa ctc cta aca gcc ccc Arg Ala Asp Ser Val Thr Trp Asn Pro His Lys Leu Leu Thr Ala Pro 345 350 355				1111
cag caa tgt tcc aca ctt tta ctg cga cat gag ggt gtc ctc gcc gag Gln Gln Cys Ser Thr Leu Leu Arg His Glu Gly Val Leu Ala Glu 360 365 370				1159
gcg cat tcc acg aac gcc gct tac ctc ttc caa aaa gac aaa ttc tac Ala His Ser Thr Asn Ala Ala Tyr Leu Phe Gln Lys Asp Lys Phe Tyr 375 380 385				1207
gac acc aaa tac gac acg ggc gac aag cac atc cag tgc ggc cgc agg Asp Thr Lys Tyr Asp Thr Gly Asp Lys His Ile Gln Cys Gly Arg Arg 390 395 400 405				1255
gcc gac gtc ctc aag ttc tgg ttc atg tgg aag gcg aag gga aca tca Ala Asp Val Leu Lys Phe Trp Phe Met Trp Lys Ala Lys Gly Thr Ser 410 415 420				1303
ggg ttg gag aaa cac gtc gat aaa gtg ttc gaa aat gcg aga ttt ttc Gly Leu Glu Lys His Val Asp Lys Val Phe Glu Asn Ala Arg Phe Phe 425 430 435				1351
acc gat tgt ata aaa aat cgg gaa ggg ttt gaa atg gtg ata gcg gag Thr Asp Cys Ile Lys Asn Arg Glu Gly Phe Glu Met Val Ile Ala Glu 440 445 450				1399
ccc gaa tac aca aac atc tgc ttt tgg tac gtg ccg aag agt ctg agg Pro Glu Tyr Thr Asn Ile Cys Phe Trp Tyr Val Pro Lys Ser Leu Arg 455 460 465				1447
ggg gcg aag gac gaa gcc gat tac aaa gac aag ctg cat aag gtg gcc				1495

-continued

```

attcataaat ccaaacggct tccctgttcg tccaaatcca catataactt catcacaat 600
gagcaacgca ccatgcttct cgcaaatttc ttttactttt tccatatatc catcaggagg 660
cattaaaatt cggccccagc taatgattgg ctccataatc acaccggcta ctgtttggct 720
taactcccat gtcatgacac gatcgatttc ctcagcactt gccagtgtat gaacatcctc 780
tggattgcga tacgtatcag gcggtgctac atgcaaaaaa ccttgcctca atggctcata 840
tttatacttt ctttgtgctt gcocctgttc tgcaagagca cccattgagt taccgtgata 900
agcgcggtag cgggaaataa acttatagcg tccatgatca cctttttgct gatgatattg 960
acgagcaatt ttaaatgctg ttccatttgc ttctgatcca ctgtagaaa agaaaatgac 1020
gtattcatca tccagccatt cattcaattt ctctgctaat ttaatggcag gaacatgact 1080
ttgtgtcaga gggaaatagc gcatttcttc aagttgctca aatgccgctc ttgcaagctc 1140
ttttcggcgc tatccaacat tcacacacca aagaccagac ataccgtcta aataacggtt 1200
tccatcaata tccgtcacc c atgccccttc tgcttttgtg ataattaaat tcgttggact 1260
aggggcccgt cctctcatcg catgccaaag gtacttttca tctgtttttt tcaaactttg 1320
tgtttgcctc gtcacttgca caatcatcag ctccat 1356

```

```

<210> SEQ ID NO 56
<211> LENGTH: 451
<212> TYPE: PRT
<213> ORGANISM: Bacillus cereus

```

<400> SEQUENCE: 56

```

Met Glu Leu Met Ile Val Gln Val Thr Glu Gln Thr Gln Ser Leu Lys
1 5 10 15
Lys Thr Asp Glu Lys Tyr Leu Trp His Ala Met Arg Gly Ala Ala Pro
20 25 30
Ser Pro Thr Asn Leu Ile Ile Thr Lys Ala Glu Gly Ala Trp Val Thr
35 40 45
Asp Ile Asp Gly Asn Arg Tyr Leu Asp Gly Met Ser Gly Leu Trp Cys
50 55 60
Val Asn Val Gly Tyr Gly Arg Lys Glu Leu Ala Arg Ala Ala Phe Glu
65 70 75 80
Gln Leu Glu Glu Met Pro Tyr Phe Pro Leu Thr Gln Ser His Val Pro
85 90 95
Ala Ile Lys Leu Ala Glu Lys Leu Asn Glu Trp Leu Asp Asp Glu Tyr
100 105 110
Val Ile Phe Phe Ser Asn Ser Gly Ser Glu Ala Asn Glu Thr Ala Phe
115 120 125
Lys Ile Ala Arg Gln Tyr His Gln Gln Lys Gly Asp His Gly Arg Tyr
130 135 140
Lys Phe Ile Ser Arg Tyr Arg Ala Tyr His Gly Asn Ser Met Gly Ala
145 150 155 160
Leu Ala Ala Thr Gly Gln Ala Gln Arg Lys Tyr Lys Tyr Glu Pro Leu
165 170 175
Gly Gln Gly Phe Leu His Val Ala Pro Pro Asp Thr Tyr Arg Asn Pro
180 185 190
Glu Asp Val His Thr Leu Ala Ser Ala Glu Glu Ile Asp Arg Val Met
195 200 205
Thr Trp Glu Leu Ser Gln Thr Val Ala Gly Val Ile Met Glu Pro Ile
210 215 220

```

-continued

Ile Thr Gly Gly Gly Ile Leu Met Pro Pro Asp Gly Tyr Met Glu Lys
 225 230 235 240

Val Lys Glu Ile Cys Glu Lys His Gly Ala Leu Leu Ile Cys Asp Glu
 245 250 255

Val Ile Cys Gly Phe Gly Arg Thr Gly Lys Pro Phe Gly Phe Met Asn
 260 265 270

Tyr Gly Val Lys Pro Asp Ile Ile Thr Met Ala Lys Gly Ile Thr Ser
 275 280 285

Ala Tyr Leu Pro Leu Ser Ala Thr Ala Val Arg Arg Glu Val Tyr Glu
 290 295 300

Ala Phe Val Gly Ser Asp Asp Tyr Asp Arg Phe Arg His Val Asn Thr
 305 310 315 320

Phe Gly Gly Asn Pro Ala Ala Cys Ala Leu Ala Leu Lys Asn Leu Glu
 325 330 335

Ile Met Glu Asn Glu Lys Leu Ile Glu Arg Ser Lys Glu Leu Gly Glu
 340 345 350

Arg Leu Leu Tyr Glu Leu Glu Asp Val Lys Glu His Pro Asn Val Gly
 355 360 365

Asp Val Arg Gly Lys Gly Leu Leu Leu Gly Ile Glu Leu Val Glu Asp
 370 375 380

Lys Gln Thr Lys Glu Pro Ala Ser Ile Glu Lys Met Asn Lys Val Ile
 385 390 395 400

Asn Ala Cys Lys Glu Lys Gly Leu Ile Ile Gly Lys Asn Gly Asp Thr
 405 410 415

Val Ala Gly Tyr Asn Asn Ile Leu Gln Leu Ala Pro Pro Leu Ser Ile
 420 425 430

Thr Glu Glu Asp Phe Thr Phe Ile Val Lys Thr Met Lys Glu Cys Leu
 435 440 445

Ala Gln Leu
 450

<210> SEQ ID NO 57
 <211> LENGTH: 747
 <212> TYPE: DNA
 <213> ORGANISM: Escherichia coli

<400> SEQUENCE: 57

```

atgatcgttt tagtaactgg agcaacggca ggttttggtg aatgcattac tcgtcgtttt    60
attcaacaag ggcataaagt tatcgccact ggcctcgccc aggagcgggt gcaggagtta    120
aaagacgaac tgggagataa tctgtatata gcccaactgg acgttcgcaa ccgocgcgct    180
attgaagaga tgctggcatc gcttctcgcc gagtgggtgca atattgatat cctggtaaat    240
aatgcccggct tggcgttggg catggagcct gcgcataaag ccagcgttga agactgggaa    300
acgatgattg ataccaacaa caaaggcctg gtatatatga cgcgcgccgt cttaccgggt    360
atggttgaac gtaatcatgg tcatattatt aacattggct caacggcagg tagctggccc    420
tatgcccgtg gtaacgttta cggtcgcgac aaagecgtttg ttcgtcagtt tagcctgaat    480
ctgcgtacgg atctgcatgg tacggcgggt cgcgtcaccg acatcgaacc gggctcgtgtg    540
gggtggcacc agttttccaa tgtccgcttt aaaggcgatg acggtaaagc ggaaaaaacc    600
tatcaaaata ccgttgcatg gacgccagaa gatgtcagcg aagccgtctg gtgggtgtca    660
acgctgcctg ctcacgtcaa tatcaatacc ctggaatga tgccggttac ccaaagctat    720
gccggactga atgtccaccg tcagtaa                                     747
    
```

-continued

<210> SEQ ID NO 58
 <211> LENGTH: 248
 <212> TYPE: PRT
 <213> ORGANISM: Proteobacteria

<400> SEQUENCE: 58

Met Ile Val Leu Val Thr Gly Ala Thr Ala Gly Phe Gly Glu Cys Ile
 1 5 10 15
 Thr Arg Arg Phe Ile Gln Gln Gly His Lys Val Ile Ala Thr Gly Arg
 20 25 30
 Arg Gln Glu Arg Leu Gln Glu Leu Lys Asp Glu Leu Gly Asp Asn Leu
 35 40 45
 Tyr Ile Ala Gln Leu Asp Val Arg Asn Arg Ala Ala Ile Glu Glu Met
 50 55 60
 Leu Ala Ser Leu Pro Ala Glu Trp Cys Asn Ile Asp Ile Leu Val Asn
 65 70 75 80
 Asn Ala Gly Leu Ala Leu Gly Met Glu Pro Ala His Lys Ala Ser Val
 85 90 95
 Glu Asp Trp Glu Thr Met Ile Asp Thr Asn Asn Lys Gly Leu Val Tyr
 100 105 110
 Met Thr Arg Ala Val Leu Pro Gly Met Val Glu Arg Asn His Gly His
 115 120 125
 Ile Ile Asn Ile Gly Ser Thr Ala Gly Ser Trp Pro Tyr Ala Gly Gly
 130 135 140
 Asn Val Tyr Gly Ala Thr Lys Ala Phe Val Arg Gln Phe Ser Leu Asn
 145 150 155 160
 Leu Arg Thr Asp Leu His Gly Thr Ala Val Arg Val Thr Asp Ile Glu
 165 170 175
 Pro Gly Leu Val Gly Gly Thr Glu Phe Ser Asn Val Arg Phe Lys Gly
 180 185 190
 Asp Asp Gly Lys Ala Glu Lys Thr Tyr Gln Asn Thr Val Ala Leu Thr
 195 200 205
 Pro Glu Asp Val Ser Glu Ala Val Trp Trp Val Ser Thr Leu Pro Ala
 210 215 220
 His Val Asn Ile Asn Thr Leu Glu Met Met Pro Val Thr Gln Ser Tyr
 225 230 235 240
 Ala Gly Leu Asn Val His Arg Gln
 245

<210> SEQ ID NO 59
 <211> LENGTH: 987
 <212> TYPE: DNA
 <213> ORGANISM: Aspergillus pseudoterreus

<400> SEQUENCE: 59

ggttgtagca gcgtaaacac atggatagtt aaataatcgg atgtacaccc actggttgaa 60
 atgacggggg cctacaacac gagattatct gatccaattt ctggtcgttg gcattctatc 120
 attcgcagcg aaaattgtcc tattaattg accatgacca aacaatctgc ggacagcaac 180
 gcaaagtcag gagttacgtc cgaatatgt cattgggcat ccaacctggc cactgacgac 240
 atcccttcgg acgtattaga aagagcaaaa taccttattc tcgacgggat tgcattgtgc 300
 tgggttggtg caagagtgc ttggtcagag aagtatgttc aggcaacgat gagctttgag 360
 ccgccggggg cctgcagggt gattggatat ggacaggtaa attttattca ctctagacgg 420
 tccacaaagt atactgacga tccttcgat agaaactggg gcctggttga gcagccatga 480

-continued

```

ccaattccgc tttcatacag gctacggagc ttgacgacta ccacagcgaa gccccctac 540
actctgcaag cattgtcctt cctgcggtct ttgacgcaag tgaggtctta gccgagcagg 600
gcaaaacaat ttccggata gatgttattc tagccgcat tgtggggtt gaatctggcc 660
cacggatcgg caaagcaatc tacggatcgg acctcttgaa caacggctgg cattgtggag 720
ctgtgtatgg cgctccagcc ggtgcgctgg ccacaggaaa gtcctcgggt ctaactccag 780
actccatgga agatgctctc ggaattgctg gcacgcaagc ctgtggttta atgtcggcgc 840
aatacggagg catggtaaag cgtgtgcaac acggattcgc agcgcgtaat ggtctcttg 900
ggggactgtt ggcccatggt gggtagcagg caatgaaagg tgcctcggag agatcttacg 960
gcggtttcct caagatgttc accaagg 987

```

```

<210> SEQ ID NO 60
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate A. pseudoterreus 5'-cadA
gene

```

```

<400> SEQUENCE: 60

```

```

ccctcgaggt cgacggtatc gatagatc ggtttagca gcgtaaacac 50

```

```

<210> SEQ ID NO 61
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate A. pseudoterreus 5'-cadA
gene

```

```

<400> SEQUENCE: 61

```

```

tctttcatag tagccttggg gaacatcttg agg 33

```

```

<210> SEQ ID NO 62
<211> LENGTH: 813
<212> TYPE: DNA
<213> ORGANISM: Aspergillus niger

```

```

<400> SEQUENCE: 62

```

```

ctactatgaa agaccgcgat gggccgatag tagtagttac ttccattaca tcatctcatc 60
cgcccgggtc ctgcctccg cggcagteta cgggtaggat cgtagcaaaa acccggggga 120
tagaccgctc gtcccagct ggagttcctg ataacctagg tagaaggtat caattgaacc 180
cgaacaactg gcaaaacatt ctgcgatcgc taggagtgag taccggcgt gatggagggg 240
gagcacgctc attggtcctg acggcagctg ccgaggggga gcaggagatc caaatatcgt 300
gagtctctg ctttgcctgg tgtatgaaac cggaaaggac tgctggggaa ctggggagcg 360
gcgcaagccg ggaatcccag ctgacaattg acccctctc atgcccggc agagcttgag 420
gtagcttttg ccccgctgt ctecccggtg tgcgcattcg actgggcgcg gcatctgtgc 480
ctctccagg agcggaggac ccagtagtaa gtaggctga cctggctggt gcgtcagtc 540
agaggttccc tcccctacc ttttctact tcccctccc cgcgctcaa cttttcttc 600
ccttttactt tctctctctc ttctcttca tccatctct ctctcact tccctcttc 660
cttcatcaa ttcattctcc aagtgagct tctcccct ctgtccctcc atctttcca 720
tcatcatctc cctcccagc tctcccctc ctctctctc ctcacgaagc ttgactaac 780

```

-continued

attacccccg cacatagaca catctaaca atg 813

<210> SEQ ID NO 63
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. niger gpdA promoter

<400> SEQUENCE: 63

atggtcacca aggtactat gaaagaccgc gatg 34

<210> SEQ ID NO 64
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. niger gpdA promoter

<400> SEQUENCE: 64

cgccgggtggc gggcattggt tagatgtgtc tatgtg 36

<210> SEQ ID NO 65
 <211> LENGTH: 1617
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: panD cDNA of Tribolium castaneum with codon optimization for A. pseudoterreus

<400> SEQUENCE: 65

cccgccaccg gcgaggacca ggacctggtg caggacctga tcgaggaacc cgccaccttc 60
 tccgacgcgc tctctgcttc cgacgaggaa ctgttcacc agaagtgcc caagccggct 120
 ccgatctaca gccccgtcag caagccgctc agcttcgagt cctgcccga ccgcccctg 180
 cacgaagagt tcctccgctc ctccgtgac gtcctgctgc aagaggccgt gttcgagggc 240
 accaacgcga agaaccgct cctgcagtgg cgcgagccc aagaactgcg ccgcctgat 300
 gacttcggcg tccgcagcgc ccgctccacg catgaggaac tgctcgaggt cctgaagaag 360
 gtcgtcacct actccgtaaa gaccggccat ccgtaactcg tcaaccagct gttctccgcc 420
 gtgatccct acggcctggt cgcccagtgg gccaccgacg cgctgaaccc ctccgtctac 480
 acctacgagg tcagccccgt gttcgtctcg atggaagagg tcgtctcgcg cgagatgcgc 540
 gccatcgtcg gcttcgaagg cggcaaaagg gacggcatct tctgccctgg cggtcgcac 600
 gccaacggct acgccatcag ctgcgcccc taccgcttca tgcccacat caagaagaag 660
 ggctgcact ccctgcccgc cctggtctcg ttcacctccg aggacgccc ctactcgatc 720
 aagaagctgg cctcgttcca aggcacggc accgacaacg tctacctgat ccgcaccgac 780
 gctcgcggtc gcatggacgt cagccacctg gtcgaagaga tcgagcgtc cctccgcgag 840
 ggcgctgcc cgttcatggt cagcgcacc gccggcacca ccgcatcgg cgccttcgat 900
 cccatcgaga agatcgccga cgtctgccag aagtacaagg tctggctgca cgtcgacgcc 960
 gcctggggcg gaggcgtct ggtgtccgcc aagcaccgcc atctgctgaa gggcatcgag 1020
 cgcccgact ccgtcacctg gaatccccac aagctgctga ccgctccgca gcagtgcagc 1080
 accctgctgc tgcgccacga gggcgtctcg gccgagcgc actccaccaa cgccgctac 1140
 ctgttcaga aggacaagtt ctacgacacc aagtacgaca ccggcgacaa gcacatccag 1200

-continued

tgcggccgctc gcgccgacgt gctgaagttc tggttcatgt ggaaggccaa gggcacctcc	1260
ggcctcgaga agcacgtgga caaggtgttc gagaacgccc gcttcttcac cgactgcatc	1320
aagaaccgtg agggcttoga gatggtgatc gccgagcctg agtacaccaa catctgtttc	1380
tggtacgtcc ccaagagcct gcgccgacgc aaggacgagg ccgactacaa ggacaagctg	1440
cacaaggtcg cccctcgcat caaagaacgc atgatgaagg aaggctccat gatggtcacc	1500
taccaggcgc agaagggcca tccgaatttc ttccgcatcg tctttcagaa ctccggcctg	1560
gacaaggcgc acatggtcca tctggtcgag gaaatcgaac gcctgggctc cgacctc	1617

<210> SEQ ID NO 66
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate panD cDNA of Tribolium
 castaneum with codon optimization for A. pseudoterreus

<400> SEQUENCE: 66
 catctaaaca atgcccgcca ccggcgagga cca 33

<210> SEQ ID NO 67
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate panD cDNA of Tribolium
 castaneum with codon optimization for A. pseudoterreus

<400> SEQUENCE: 67
 atccaaccca tcagaggctc gagcccaggc gtctc 35

<210> SEQ ID NO 68
 <211> LENGTH: 1040
 <212> TYPE: DNA
 <213> ORGANISM: Aspergillus niger

<400> SEQUENCE: 68
 tgatgggttg gatgacgatg acttcatgtg atttgttat ttagaatatt ttatatttcc 60
 ttttcttctt ctcaccaccg atccccctaa cactcttctt tcatttgctt cagatttctc 120
 gggttcttct ttttcttctt ccccgattat ccaactatct tttgctagac cggcctgcgc 180
 cctggcatgc atcataaaat catgtccggt ggtcatcctc tgttttgat atccgtcata 240
 taaagtattc ttttattccc tccccctcg gtcgtcttcc gctgtcccgc ttctacctc 300
 cggtttatag agcatggctc atctcttccg tacatttccg ttggtactag catttatgtc 360
 ttcagctagt atagaagctg ccgcagttgt tcgcttacta cctgcctaag tccttaactt 420
 tttaaagtgt ttaacctata cgtagtgtta aacgagtact gggaggtggt gaggtagaaa 480
 atgttctgca cgggcagtggt gtatttggta gtgtgtaagg cggttattta tcaggctgac 540
 gctaagaact tctatgggag cagtatggga tcgcggtcca tagaagtaca caaaatctaa 600
 gagtcgtttg ataattaatt gattcccggc agggctctct tgggattgag agaactggtt 660
 actttgattt gagatattgt aaagcttaag gctcttaaca cgtaacgagc aaacagcagg 720
 ggggaaatcg ggaaaagggt cgtgggggtga ataaaaaagt tgaataaga cactgtatct 780
 tgctgggggt gaataaagag agaataaag agaggtaaat tccactcagc cccttttctt 840
 cgctctccaa acatcaaaact ccgcccggcc acccacagga tcccgaacaa gtggaagata 900
 tgtgccggctc cagacccttc gcacagctaa aagcagacct tcataagcgt ttccgggtag 960

-continued

tattcgacaca cctgaactgg cacgtcgggg acacaactgt ttttgatata caagaacaca 1020
caccacccat ctaggactca 1040

<210> SEQ ID NO 69
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate bidirectional terminator
from A. niger elf3/multifunctional chaperone

<400> SEQUENCE: 69
gggctccgac ctctgatggg ttggatgacg atg 33

<210> SEQ ID NO 70
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate bidirectional terminator
from A. niger elf3/multifunctional chaperone

<400> SEQUENCE: 70
tctggcccag ctctgagtcc tagatgggtg gtg 33

<210> SEQ ID NO 71
<211> LENGTH: 1350
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: codon optimized synthetic cDNA of beta-alanine-
pyruvate aminotransferase (BAPAT) of Bacillus cereus

<400> SEQUENCE: 71
gagctgggcc agacattcct tcatagtctt gacgatgaag gtgaagtcct cttcggatgat 60
ggacagcggga ggggcgagct gcaggatggt gttgtagccg gccacgggtg cgcggttctt 120
gccgatgac agacccttct ctttgcaggc gttgatgacc ttgttcatct tttcgatgga 180
ggcgggctct ttggtctgct taccctcgac gaggctcgata ccacgcagga ggccttgcc 240
gcggacgtcc ccgacgttgg ggtgctcttt gacgtcctcc aactcgtaca gcaggcgctc 300
gcccagttct ttggaccgct cgatgagctt ctcgttttcc atgatctcga ggttcttcag 360
ggccagcgcg caggcggcag ggttgccgcc gaaggtgttg acatggcggga agcggtcgta 420
gtcgtcggag ccgacgaagg cctcgtagac ctccgcccgg accgcccgtg cagacagcgg 480
caggtaggcc gaggtgatac ccttgcccat ggtgataatg tcgggcttga cgcgtagtt 540
catgaagcgg aaggccttgc cgtgctgacc gaagccgcag atgacctcgt cgcagatcag 600
caggcgcgcg tgcttttcgc agatctcttt gaccttttcc atgtagccgt ccggcggcat 660
caggatgccca ccaccggtga tgatgggttc catgatgacg ccggcgcaggg tctgggacag 720
ctcccaggtc atgacgcggt cgatttcttc ggccgaggcc aggggtgtgca cgtcctcggg 780
gttgcgatag gtgtccggag gggccacgtg caggaagccc tgaccgaggg gctcgtactt 840
gtacttgcgc tgggcctgac cgttcgcggc cagggcaccc atggagtgc cgtggtaggc 900
gcggtagcga gagatgaact tgtageggcc gtggtcaccc ttctgctggt ggtactggcg 960
ggcgatcttg aaggcggttt cgttggcctc cgaccggag ttggagaaga agatgacgta 1020
ctcgtcgtcc agccactcgt tcagcttctc ggccagcttg atggcgggga cgtgcgactg 1080
cgtcagcggg aagtacggca tctcttcacg ctgctcgaag gcagcgcgag ccagctcttt 1140

-continued

```

geggccgtag cgcacgttga cgcaccacag gccggacatg ccgtccaggt agcgggttgc 1200
gtcgatgtcg gtgaccacag cgccttcggc cttggtgatg atcaggttgg tcggactcgg 1260
agcggcaccg cgcacgtggc gccacaggta cttctcgtcg gttttcttca ggctctgggt 1320
ctgctcggtg acctggacga tcatcagttc 1350

```

```

<210> SEQ ID NO 72
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate codon optimized
synthetic cDNA of beta-alanine-pyruvate aminotransferase (BAPAT)
of Bacillus cereus

```

```

<400> SEQUENCE: 72
catctaggac tcagagctgg gccagacatt ccttc 35

```

```

<210> SEQ ID NO 73
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate codon optimized
synthetic cDNA of beta-alanine-pyruvate aminotransferase (bapat)
of bacillus cereus

```

```

<400> SEQUENCE: 73
gtccatcaac atggaactga tgatcgtcca ggtcac 36

```

```

<210> SEQ ID NO 74
<211> LENGTH: 704
<212> TYPE: DNA
<213> ORGANISM: Aspergillus niger
<400> SEQUENCE: 74
catgttgatg gactggaggg ggatgagtta tggatcagtg aaactgggag aaaacaaaga 60
tggcaaaggg agaacatggc ccagatatag gaaaaaacgg aggaggcaaa aatgtaagcg 120
ctccggactt gctgtttcgg tgtgcactag cagcagcggg ggggaagggtg gtgagtgttc 180
accgaggacc caaaaagaat gagcggatgg cggatgagtg acggagaagg gaaggacggg 240
gggggaatta gaggtggaga ggtccgatcc atcaaataga ccaggctcgg cacagccaag 300
tttcccaaat gatcaactaa tcaatgggac ttggtgctaa atccggagat gccagatcat 360
tgatagacag acaggatgga gtgatggcat atagacagga ggatggatgg atggatagat 420
ggaggggtca agcacaacat ggtgggatga tggcggggtc atgactagca gctaagagga 480
agaagaggag gatgaaatgg acagagaaag atgggagggg tgataaaatg agtatatggg 540
acaagtcata cttacaggac cttgaagatg gtggtgttac tatctaagaa aggctttttt 600
tgagagtact cttaacacaa gaggaggagg gaggaggggg aagtagtaga taaataataa 660
acacgaccac agacttgcta caggctactt cttgtaagct cgag 704

```

```

<210> SEQ ID NO 75
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate an A. niger eno1
promoter

```

```

<400> SEQUENCE: 75

```

-continued

cgatcatcag ttccatggtg atggactgga ggg 33

<210> SEQ ID NO 76
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. niger enol promoter

<400> SEQUENCE: 76

gaactagtgg atccccggg ctgctgtaac tcgagcttac aagaagtagc c 51

<210> SEQ ID NO 77
 <211> LENGTH: 885
 <212> TYPE: DNA
 <213> ORGANISM: Aspergillus nidulans

<400> SEQUENCE: 77

tctgtacagt gaccggtgac tctttctggc atgcggagag acggacggac gcagagagaa 60
 gggctgagta ataagccact ggccagacag ctctggcggc tctgaggtgc agtggatgat 120
 tattaatccg ggaccggcgc cccctccgcc ccgaagtgga aaggctggtg tgcccctcgt 180
 tgaccaagaa tctattgcat catcgagaa tatggagctt catcgaatca ccggcagtaa 240
 gcgaaggaga atgtgaagcc aggggtgat agccgtcggc gaaatagcat gccattaacc 300
 taggtacaga agtccaattg cttccgatct ggtaaaagat tcacgagata gtaccttctc 360
 cgaagtaggt agagcgagta cccggcgcgt aagctcccta attggccat ccggcatctg 420
 tagggcgtcc aaatategtg cctctcctgc ttgcccgggt gtatgaaacc gaaaggccg 480
 ctcaggagct ggccagcggc gcagaccggg aacacaagct ggcagtcgac ccatccggtg 540
 ctctgcactc gacctgctga ggtccctcag tccctggtag gcagctttgc cccgtctgtc 600
 gcgccggtgt gtcggcgggg ttgacaaggt cgttgcgtea gtccaacatt tgttgccata 660
 ttttctgct ctecccacca gctgctcttt tctttctctt ttcttttccc atcttcagta 720
 tattcatctt cccatccaag aacctttatt tcccctaagt aagtactttg ctacatccat 780
 actccatcct tcccatccct tattcctttg aacctttcag ttcgagcttt cccacttcat 840
 gcgagcttga ctaacagcta ccccgcttga gcagacatca ccatg 885

<210> SEQ ID NO 78
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. nidulans gpdA promoter

<400> SEQUENCE: 78

ccctcgaggt cgacggatc gatagttaac tctgtacagt gaccggtgac 50

<210> SEQ ID NO 79
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate an A. nidulans gpdA promoter

<400> SEQUENCE: 79

tgaccagcac gatcatggtg atgtctgctc aag 33

-continued

<210> SEQ ID NO 80
 <211> LENGTH: 741
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: codon optimized synthetic cDNA of E. coli HPDH

<400> SEQUENCE: 80

```

atcgtgctgg tcacggggcgc gaccgcccgt ttcggcgagt gcatcaccgc ccgcttcac 60
cagcagggcc acaaggtgat cgctaccgga cgccgccaag agcgctcca agagctgaag 120
gatgagctgg gcgacaacct gtacattgcc cagctggacg tgcgcaaccg ggctgccac 180
gaagaaatgc tcgcctcgct gcccgccgag tggtgcaaca tcgacatcct ggtcaacaac 240
gccggtctgg ccctcgcat ggaaccggcg cacaaggcca gcgtcgagga ctgggaaacc 300
atgatcgaca ccaacaacaa gggactcgtc tacatgaccg gcgctgtgct gcccgcatg 360
gtcagcgcga accacggcca catcatcaac atcggctcca ccgctggcag ctggccctac 420
gctggcggca acgtctatgg cgcgaccaag gcgttcgtcc gccagttctc cctgaaactg 480
cgcaccgacc tgcacggcac cgccgtccgc gtgaccgaca ttgagcccgg tctggtcggc 540
ggcaccgagt tcagcaacgt ccgcttcaag ggcgacgacg gcaaggccga gaaaacctac 600
cagaacaccg tcgctctgac cctgaggat gtcagcgagg ccgtctggtg ggtcagcaact 660
ctgcccgcgc acgtcaacat caacaccctc gagatgatgc ccgtcacgca gtcctacgcc 720
ggcctgaacg tccaccgcca a 741

```

<210> SEQ ID NO 81
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate a codon optimized synthetic cDNA of E. coli HPDH

<400> SEQUENCE: 81

```

agacatcacc atgatcgtgc tggtcacggg cgc 33

```

<210> SEQ ID NO 82
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate a codon optimized synthetic cDNA of E. coli HPDH

<400> SEQUENCE: 82

```

gccatcggtc ctattggcgg tggacgttca ggc 33

```

<210> SEQ ID NO 83
 <211> LENGTH: 473
 <212> TYPE: DNA
 <213> ORGANISM: Aspergillus nidulans

<400> SEQUENCE: 83

```

taggaccgat ggctgtgtag aagtactcgc cgatagtgga aaccgacgcc ccagcactcg 60
tccgagggca aaggaataga gtatagtcgc accgcgggat ccaactaacg ttactgaaat 120
catcaaacag cttgacgaat ctggatataa gatcgttggg gtcgatgtca gctccggagt 180
tgagacaaat ggtgttcagg atctcgataa gatacgttca tttgtccaag cagcaaagag 240

```

-continued

tgccttctag tgatttaata gctccatgtc aacaagaata aaacgcgttt tcgggtttac	300
ctcttocaga tacagctcat ctgcaatgca ttaatgcatt gactgcaacc tagtaacgcc	360
ttcaggctcc ggccaagaga agaatagctt agcagagcta ttttcatttt cgggagacga	420
gatcaagcag atcaacggtc gtcaagagac ctacgagact gaggaatccg ctc	473

<210> SEQ ID NO 84
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate the trpC terminator of
 A. nidulans

<400> SEQUENCE: 84

cgtcaccgc caataggacc gatggctgtg tag	33
-------------------------------------	----

<210> SEQ ID NO 85
 <211> LENGTH: 45
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate the trpC terminator of
 A. nidulans

<400> SEQUENCE: 85

gaactagtgg atccccggg ctgcagagcg gattcctcag tctcg	45
--	----

<210> SEQ ID NO 86
 <211> LENGTH: 510
 <212> TYPE: DNA
 <213> ORGANISM: Aspergillus nidulans

<400> SEQUENCE: 86

ccctcgaggt cgacggtatc gatataggac cgatggctgt gtagaagtac tcgccgatag	60
tggaaaccga cgccccagca ctcgctcgag ggcaaaggaa tagagtagat gccgaccgcg	120
ggatccactt aacgttactg aaatcatcaa acagcttgac gaatctggat ataagatcgt	180
tgggtgctgat gtcagctccg gagttgagac aaatgggtgtt caggatctcg ataagatacg	240
ttcatttgtc caagcagcaa agagtgcctt ctagtgattt aatagctcca tgtcaacaag	300
aataaaacgc gttttcgggt ttacctcttc cagatacagc tcatctgcaa tgcattaatg	360
cattgactgc aacctagtaa cgccttcagg ctccggcgaa gagaagaata gcttagcaga	420
gctattttca ttttcgggag acgagatcaa gcagatcaac ggctcgtcaag agacctacga	480
gactgaggaa tccgctctct gacagacggg	510

<210> SEQ ID NO 87
 <211> LENGTH: 44
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer used to isolate a trpC terminator of A.
 nidulans

<400> SEQUENCE: 87

ccctcgaggt cgacggtatc gatataggac cgatggctgt gtag	44
--	----

<210> SEQ ID NO 88
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: primer used to isolate a trpC terminator of *A. nidulans*

<400> SEQUENCE: 88

cccgtctgtc agagagcgga ttctcagtc tcg 33

<210> SEQ ID NO 89

<211> LENGTH: 2005

<212> TYPE: DNA

<213> ORGANISM: *Aspergillus oryzae*

<400> SEQUENCE: 89

tctgacagac gggcaattga ttacgggatc ccattggtaa cgaaatgtaa aagctaggag 60
atcgctccgcc gatgtcagga tgatttcact tgtttcttgt ccggctcacc ggtcaaagct 120
aaagaggagc aaaaggaacg gatagaatcg ggtgccgctg atctatacgg tatagtgtccc 180
ttatcacggt gactcaaccc atgctattta actcaacccc tccttctgaa cccaccatc 240
ttcttccttt tcctctcatc ccacacaatt ctctatctca gatttgaatt ccaaaagtcc 300
tcggacgaaa ctgaacaagt ttctctccct tcgataaaacc tttgggtgatt ggaataactg 360
accatcttct atagtccca aaccaaccga caatgtaaat aactcctcg attagcctc 420
tagagggcat acgatggaag tcattggaata cttttggctg gactctcaca atgatcaagg 480
tatcttaggt aacgtctttg gcgtgggccc gtgttcgttc ccagtcacg atgcattcac 540
atgccctccc taagctgggc cctagactct aggatcctag tctagaagga catggcatcg 600
atggactggg ttcgtttctga gattatacgg ctaaaacttg atctggataa taccagcgaa 660
aagggtcatg ccttctctcg ttcttctgtg tgatggaatg gctaacagat gatagtcatt 720
gcaacttgaa acatgtctcc tccagctgcc atctacgaac cactgtggc cgctaccggc 780
ctcaagggta aggtcgtggt ttctgagacc gtccccgttg agggagcttc tcagaccaag 840
ctggtggacc atttcgttgg caagtgggac gagttcaagt tcgcccctat ccgcgaaagc 900
caggtctctc gtgccatgac cagacgttac tttgaggacc tggacaagta cgctgaaagt 960
gacgttgtea ttgttggtgc tggttcctgc ggtctgagca ctgctgacgt cttggccaag 1020
gctcgtccgg acctgaagat tgctatcgtc gaggccagcg tctctcctgg tcagtagtcc 1080
atgatggatt gccttgcaact cagctttccg gaactaacgt gcaatagggt gcggtgctg 1140
gttgggtggc caactctttt ctgctatggt catgcgccc cccgcggaag tcttctgaa 1200
cgagctgggt gttccttacg aagaggacgc aaaccccaac tacgttgtcg tcaagcacgc 1260
ctccctgttt acctcgacac tcattgtgaa ggttctctcc tccccaatg tcaagctctt 1320
caatgctacc gctgttgagg acttgatcac ccgtccgacc gagaacggca accccagat 1380
tgctgggtgt gtcgtcaact ggacgtggt cacccttcac cacgatgac actcctgcat 1440
ggaccccaac actatcaacg ctctgtcat catcagtacc actggtcacg atgggcccatt 1500
cggcgccttc tgtcgaagc gcttggtgtc catgggcagc gtcgacaagc taggtggcat 1560
gcgtgggtct gacatgaact cggccgagga tgccatcgtc aagaacccc gcgaggttac 1620
taagggtctg ataactggcg gtatggagct gtctgaaatt gatggcttta accgcatggg 1680
ccctaccttc ggtgccatgg ttctcagtggt tgtcaaggct gccgaggagg cattgaaggt 1740
gttcgacgag cgtcagcgcg agtgtgctga gtaaatgact cactaccga atgggttcag 1800
tgcatgaacc ggatttctct tacggtcttt gacgataggg gaatgatgat tatgtgatag 1860
ttctgagatt tgaatgaact cgttagctcg taatccacat gcatatgtaa atggctgtgt 1920

-continued

```

cccgtatgta acggtggggc attctagaat aattatgtgt aacaagaaag acagtataat 1980
acaaacaaag atgcaagagc ggctc 2005

```

```

<210> SEQ ID NO 90
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate the A. oryzae ptrA
selection marker gene

```

```

<400> SEQUENCE: 90
gaggaatccg ctctctgaca gacgggcaat tgattac 37

```

```

<210> SEQ ID NO 91
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate the A. oryzae ptrA
selection marker gene

```

```

<400> SEQUENCE: 91
gaatgttgct gaggagccgc tcttgcatct ttg 33

```

```

<210> SEQ ID NO 92
<211> LENGTH: 908
<212> TYPE: DNA
<213> ORGANISM: Aspergillus pseudoterreus

```

```

<400> SEQUENCE: 92
ctcagcaaca ttcgccatgt tcatgtacag ctttcaacgg cctcgaacag tcaactgtgga 60
tggataccag aggagagacc catcagttca atcgcagggc agatgagtggt cgcatacatt 120
ctcgcctgac agctggtcga ccagcaatgt cttttgtccc agttttctga gtttgatgac 180
aacctggaga ggccagaagt ttgggatctg gccaggaagg ttacttcac tcaaagcgaa 240
gagtttgatc aagacggcaa ctgtctcagt gcggtctcgc tgaggattga gttcaacgat 300
ggttcttcta ttacggaaag tgcgagaag cctcttggtg tcaaagagcc catgccaac 360
gaacggattc tccacaaata ccgaaccctt gctggtagcg tgacggacga atcccgggtg 420
aaagagattg aggatcttgt cctcggcctg gacaggetca ccgacattag cccattgctg 480
gagctgctga attgcccgt gaaatcgcca ctggtataaa tgggaagcga tatggaaca 540
tttcatgtca cgggcacaaa ttctaggtca tctcgtacct ggatggtgaa accaccagcg 600
gtttagcaga tagaagatag actccttctg ctctgcgttg cgtcttgaat ttagttcgtt 660
cactggctta agaacttaga atgcaataca gtctctctta tttcttatta aaatcacgta 720
tccccacatt cggcgactgg aggatacga agcagtggtg gtggtgctcc ccgtaatgga 780
tatgatattg ctgactggac tattctatga ccattccctc caacggagat cctttctcga 840
cactttagat gttgacgctg tctggaggaa ctacttttgc gctgcaaaga ctatgagcag 900
tggagctg 908

```

```

<210> SEQ ID NO 93
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer used to isolate an A. pseudoterreus 3'-
cad1 gene fragment

```

-continued

<400> SEQUENCE: 93

gcaagagcgg ctctctcagca acattcgcca tgttc 35

<210> SEQ ID NO 94

<211> LENGTH: 49

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to isolate an A. pseudoterreus 3'-
cad1 gene fragment

<400> SEQUENCE: 94

gaactagtgg atcccccggg ctgcacagct ccaactgctca tagtctttg 49

<210> SEQ ID NO 95

<211> LENGTH: 2099

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: combination of SEQ ID NOS: 77, 80, and 83

<400> SEQUENCE: 95

tctgtacagt gaccggtgac tctttctggc atgctggagag acggacggac gcagagagaa 60

gggctgagta ataagccact ggccagacag ctctggcggc tctgaggtgc agtggatgat 120

tattaatccg ggaccggcgc cccctccgcc ccgaagtgga aaggctgggtg tgcccctcgt 180

tgaccaagaa tctattgcat catcgagaa tatggagctt catcgaatca ccggcagtaa 240

gcgaaggaga atgtgaagcc aggggtgat agccgtcggc gaaatagcat gccattaacc 300

taggtacaga agtccaattg ctccgatct ggtaaaagat tcacgagata gtaccttctc 360

cgaagtaggt agagcgagta cccggcgcgt aagctcccta attggcccat ccggcatctg 420

tagggcgtcc aaatategtg cctctcctgc ttgcccgggt gtatgaaacc gaaaggccg 480

ctcaggagct ggccagcggc gcagaccggg aacacaagct ggcagtcgac ccatccgggtg 540

ctctgcactc gacctgctga ggtccctcag tccctggtag gcagctttgc cccgtctgtc 600

cgcccgggtg gtcggcgggg ttgacaaggt cgttgctca gtccaacatt tgttgccata 660

ttttcctgct ctecccacca getgetcttt tcttttctct ttcttttccc atcttcagta 720

tattcatctt cccatccaag aacctttatt tcccctaagt aagtactttg ctacatccat 780

actccatcct tccatccct tattcctttg aacctttcag ttcgagcttt cccacttcat 840

cgagcttga ctaacagcta ccccgcttga gcagacatca ccatgatcgt gctggtcacg 900

ggcgcgaccg ccggtttcgg cgagtgcac acccgccgct tcatccagca gggccacaag 960

gtgatcgcta ccggacgcgc ccaagagcgc ctccaagagc tgaaggatga gctgggcgac 1020

aacctgtaca ttgccagct ggacgtgcgc aaccgggctg ccatcgaaga aatgctcgcc 1080

tcgctgcccg ccgagtggtg caacatcgac atcctggtea acaacgcccg tctggccctc 1140

ggcatggaac cggcgcacaa ggccagcgtc gaggactggg aaacctgat cgacaccaac 1200

aacaagggac tcgtctacat gaccgcgct gtgtgccccg gcaggtgca ggcgaaccac 1260

ggccacatca tcaacatcgg ctccaccgct ggcagctggc cctacgctgg cggcaacgctc 1320

tatggcgcga ccaaggcgtt cgtccgccag ttctccctga acctgcgcac cgacctgcac 1380

ggcaccgccc tccgctgac cgacattgag cccggtctgg tcggcggcac cgagttcagc 1440

aacgtccgct tcaagggcga cgacggcaag gccgagaaaa cctaccagaa caccgtcgct 1500

ctgacccctg aggatgtcag cgaggcgtc tgggtgggta gcaactctgcc cgcgcacgctc 1560

-continued

```

aacatcaaca ccctcgagat gatgccgctc acgcagtcct acgccggcct gaacgtccac 1620
cgccaatagg accgatggct gtgtagaagt actcgccgat agtggaaacc gacgccccag 1680
cactcgctccg agggcaaagg aatagagtag atgcccgaccg cgggatccac ttaacgttac 1740
tgaaatcatc aaacagcttg acgaatctgg atataagatc gttgggtgctg atgtcagctc 1800
cggagttgag acaaatggtg ttcaggatct cgataagata cgttcatttg tccaagcagc 1860
aaagagtgcc ttctagtgat ttaatagctc catgtcaaca agaataaaac gcgttttcgg 1920
gtttacctct tccagatata gctcatctgc aatgcattaa tgcattgact gcaacctagt 1980
aacgccttca ggctccggcg aagagaagaa tagcttagca gagctatctt cattttcggg 2040
agacgagatc aagcagatca acggctcgtca agagacctac gagactgagg aatccgctc 2099

```

```

<210> SEQ ID NO 96
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer to isolate Fragments 7 to 9

```

```

<400> SEQUENCE: 96
acaggctact tcttgtaagc tcgagtttct gtacagtgac cggtgac 47

```

```

<210> SEQ ID NO 97
<211> LENGTH: 2913
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: combination of SEQ ID NOS: 89 and 92

```

```

<400> SEQUENCE: 97
tctgacagac gggcaattga ttacgggac ccattggtaa cgaaatgtaa aagctaggag 60
atcgctccgc gatgtcagga tgatttact tgtttctgt cggctcacc ggtcaaagct 120
aaagaggagc aaaaggaacg gatagaatcg ggtgccgctg atctatacgg tatagtgcc 180
ttatcacggt gactcaacc atgctattta actcaacccc tccttctgaa ccccaccatc 240
ttcttcctt tcctctcacc ccacacaatt ctctatctca gattgaatt ccaaagtcc 300
tcggacgaaa ctgaacaagt ctctctccct tcgataaacc tttggtgatt ggaataactg 360
accatcttct atagtccca aaccaaccga caatgtaaat aactcctcg attagccctc 420
tagagggcat acgatggaag tcattggaata cttttggctg gactctcaca atgatcaagg 480
tatcttaggt aacgtctttg gcgtgggccc gtgttcgctc ccagtcacg atgcattcac 540
atgcctccc taagctgggc cctagactct aggatcctag tctagaagga catggcatcg 600
atggactggg ttcgttctga gattatcagg ctaaaacttg atctggataa taccagcgaa 660
aagggtcatg ccttctctcg ttcttctctg tgatggaatg gctaacagat gatagtcatt 720
gcaacttgaa acatgtctcc tccagctgcc atctacgaac ccaactgtggc cgctaccggc 780
ctcaagggta aggtcgtggt ttctgagacc gtcccgttg agggagcttc tcagaccaag 840
ctggttgacc atttcggtgg caagtgggac gagttcaagt tcgcccctat ccgcaaaagc 900
caggtctctc gtgccatgac cagacgttac tttgaggacc tggacaagta cgctgaaagt 960
gacgttgcca ttgttggtgc tggttcctgc ggtctgagca ctgcgtacgt cttggccaag 1020
gctcgteccg acctgaagat tgctatctgc gaggccagcg tctctcctgg tcagtagtcc 1080
atgatggatt gccttgcaat cagcttccg gaactaacgt gcaataggtg gcggtgctg 1140
gttgggtggc caactctttt ctgctatggt catgcgccgt cccgcaagag tcttctctgaa 1200

```

-continued

```

cgagctgggt gttccttacg aagaggacgc aaacccaac tacgttgtcg tcaagcacgc 1260
ctccctgttt acctogacac tcatgtogaa ggttctctcc ttcccgaatg tcaagctctt 1320
caatgctacc gctggtgagg acttgatcac ccgtecgacc gagaacggca acccccagat 1380
tgctgggtgt gtcgtcaact ggacgctggc cacccttcac cacgatgatc actcctgcat 1440
ggacccaac actatcaacg ctctgtcat catcagtacc actggtcacg atgggccatt 1500
cgggcccttc tgtgcgaagc gcttggtgtc catgggcagc gtcgacaagc taggtggcat 1560
gcgtggcttc gacatgaact cggccgagga tgccatcgtc aagaacaccc gcgaggttac 1620
taaggccttg ataactcggc gtatggagct gtctgaaatt gatggcttta accgcatggg 1680
ccctaccttc ggtgccatgg ttctcagtgg tgtcaaggct gccgaggagg cattgaaggc 1740
gttcgacgag cgtcagcgcg agtgtgctga gtaaagtact cactaccgca atgggttcag 1800
tgcatgaacc ggatttgtct tacggtcttt gacgataggg gaatgatgat tatgtgatag 1860
ttctgagatt tgaatgaact cgttagctcg taatccacat gcatatgtaa atggctgtgt 1920
cccgtatgta acggtggggc attctagaat aattatgtgt aacaagaaag acagtataat 1980
acaaacaaag atgcaagagc ggctcctcag caacattcgc catgttcatg tacagctttc 2040
aacggcctcg aacagtcact gtggatggat accagaggag agacccatca gttcaatcgc 2100
agggcagatg agtgtcgcac acattctcgc cgtccagctg gtcgaccagc aatgtctttt 2160
gtcccagttt tctgagtttg atgacaacct ggagaggcca gaagtttggg atctggccag 2220
gaaggttact tcactcctaaa gcgaagagtt tgatcaagac ggcaactgtc tcagtgcggg 2280
tcgctgaggg attgagttca acgatggctc ttctattacg gaaagtgtcg agaagcctct 2340
tggtgtcaaa gagcccatgc caaacgaaac gattctccac aaataccgaa cccttgctgg 2400
tagcgtgacg gacgaatccc gggtgaaaga gattgaggat cttgtcctcg gcctggacag 2460
gctcacccac attagcccat tgctggagct gctgaattgc cccgtgaaat cgccactggt 2520
ataaatggga agcgatatgg aaacatttca tgtcacgggc acaaattcta ggcatatcg 2580
tacctggatg gtgaaaccac cagcggttta gcagatagaa gatagactcc ttctgctctg 2640
cgttgcgtct tgaatttagt tcgttcactg gcttaagaac ttagaatgca atacagtctc 2700
tcttatttct tattaaaac acgtattccc acattcggcg actggaggat acgaaagcag 2760
tgttgggtgt gctcccgcga atggatatga ttttgctgac tggactatc tatgaccatt 2820
ccctccaacg gagatccttt ctgcacactt tagatgttga cgctgtctgg aggaactact 2880
tttgcgctgc aaagactatg agcagtggag ctg 2913

```

<210> SEQ ID NO 98

<211> LENGTH: 49

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: primer used to isolate Fragments 11-12

<400> SEQUENCE: 98

actaaagga acaaaagctg gagctcagct ccaactgctca tagtctttg

49

We claim:

1. An isolated recombinant *Aspergillus* fungus capable of producing 3-hydroxypropionic acid (3-HP), comprising: a genetic inactivation of an endogenous cis-aconitic acid decarboxylase (*cadA*) gene, an exogenous nucleic acid molecule encoding aspartate 1-decarboxylase (*panD*), an exogenous nucleic acid molecule encoding β -alanine-pyruvate aminotransferase (*BAPAT*), and an exogenous nucleic acid molecule encoding 3-hydroxypropionate dehydrogenase (*HPDH*).
2. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the *Aspergillus* fungus is *Aspergillus pseudotereus*.
3. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the *Aspergillus* fungus is *Aspergillus terreus*.
4. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the endogenous *cadA* gene is genetically inactivated by complete or partial deletion mutation or by insertional mutation.
5. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the *cadA* gene prior to its genetic inactivation encodes a cis-aconitic acid decarboxylase with an amino acid sequence having at least 80% sequence identity to the amino acid sequence of SEQ ID NO: 50 or 52.
6. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the *cadA* gene prior to its genetic inactivation comprises a coding sequence having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO:49, 51, 59 or 92.
7. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the exogenous nucleic acid molecule encoding *panD* comprises a nucleotide sequence having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO: 53 or 65, and/or encodes an amino acid sequence having at least 80% sequence identity to the amino acid sequence of SEQ ID NO: 54.
8. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the exogenous nucleic acid molecule encoding *BAPAT*

- comprises a nucleotide sequence having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO: 55 or 71, and/or
- encodes an amino acid sequence having at least 80% sequence identity to the amino acid sequence of SEQ ID NO: 56.
9. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the exogenous nucleic acid molecule encoding *HPDH* comprises a nucleotide sequence having at least 80% sequence identity to the nucleotide sequence of SEQ ID NO: 57 or 80, and/or encodes an amino acid sequence having at least 80% sequence identity to the amino acid sequence of SEQ ID NO: 58.
10. The isolated recombinant *Aspergillus* fungus of claim 1, wherein the exogenous nucleic acid molecule encoding *panD*, the exogenous nucleic acid molecule encoding *BAPAT*, and the exogenous nucleic acid molecule encoding *HPDH* are part of a single exogenous nucleic acid molecule.
11. A composition comprising the isolated recombinant *Aspergillus* fungus of claim 1.
12. A kit, comprising:
 - the isolated recombinant *Aspergillus* fungus of claim 1; and
 - a medium for culturing the fungus.
13. A method of making 3-hydroxypropionic acid (3-HP), comprising:
 - culturing the isolated *Aspergillus* fungus of claim 1 in a culture media under conditions that permit the isolated *Aspergillus* fungus to make 3-HP; thereby making 3-HP.
14. The method of claim 13, wherein the isolated *Aspergillus* fungus is cultured in Riscaldati medium or modified Riscaldati medium comprising 20x trace elements.
15. The method of claim 13, further comprising isolating the 3-HP from the culture media or from the isolated *Aspergillus* fungus.

* * * * *