



US011316812B2

(12) **United States Patent**
Hsu et al.

(10) **Patent No.:** **US 11,316,812 B2**
(45) **Date of Patent:** ***Apr. 26, 2022**

(54) **RNA TARGETING METHODS AND COMPOSITIONS**

- (71) Applicant: **Salk Institute for Biological Studies**, La Jolla, CA (US)
- (72) Inventors: **Patrick D. Hsu**, San Francisco, CA (US); **Silvana Konermann**, Menlo Park, CA (US)
- (73) Assignee: **Salk Institute for Biological Studies**, La Jolla, CA (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.
- (21) Appl. No.: **17/352,564**
- (22) Filed: **Jun. 21, 2021**

(65) **Prior Publication Data**

US 2021/0344627 A1 Nov. 4, 2021

Related U.S. Application Data

- (60) Continuation of application No. 16/257,493, filed on Jan. 25, 2019, now Pat. No. 1,122,857, which is a division of application No. 15/937,699, filed on Mar. 27, 2018, now Pat. No. 10,476,825.
- (60) Provisional application No. 62/639,178, filed on Mar. 6, 2018, provisional application No. 62/572,963, filed on Oct. 16, 2017, provisional application No. 62/548,846, filed on Aug. 22, 2017.

(51) **Int. Cl.**

C12N 15/85 (2006.01)
H04L 51/00 (2022.01)
C12N 15/10 (2006.01)
C12N 15/113 (2010.01)
C12N 15/82 (2006.01)
C12N 9/22 (2006.01)
H04L 9/08 (2006.01)
H04L 9/30 (2006.01)
H04L 45/745 (2022.01)
H04L 51/08 (2022.01)
H04L 51/18 (2022.01)

(52) **U.S. Cl.**

CPC **H04L 51/12** (2013.01); **C12N 9/22** (2013.01); **C12N 15/102** (2013.01); **C12N 15/113** (2013.01); **C12N 15/8201** (2013.01); **C12N 15/8213** (2013.01); **C12N 15/85** (2013.01); **H04L 9/08** (2013.01); **H04L 9/30** (2013.01); **H04L 45/745** (2013.01); **H04L 51/08** (2013.01); **H04L 51/14** (2013.01); **H04L 51/18** (2013.01); **C12N 2310/16** (2013.01); **C12N 2310/20** (2017.05); **C12N 2310/3519** (2013.01)

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

8,871,445 B2	10/2014	Cong et al.
10,392,616 B2	8/2019	Cheng et al.
10,476,825 B2	11/2019	Hsu et al.
10,666,592 B2	5/2020	Hsu et al.
10,931,613 B2	2/2021	Hsu et al.
11,005,799 B2	5/2021	Hsu et al.
11,025,574 B2	6/2021	Hsu et al.
11,032,224 B2	6/2021	Hsu et al.
11,032,225 B2	6/2021	Hsu et al.
2015/0218573 A1	8/2015	Loque et al.
2016/0208243 A1	7/2016	Zhang et al.
2017/0145394 A1	5/2017	Yeo et al.
2018/0282715 A1	10/2018	Carter et al.
2018/0334685 A1	11/2018	Yeo et al.
2019/0002875 A1	1/2019	Cheng et al.
2019/0002889 A1	1/2019	Cheng et al.
2019/0169595 A1	6/2019	Hsu et al.
2019/0207890 A1	7/2019	Hsu et al.
2020/0199556 A1	6/2020	Hsu et al.
2020/0322297 A1	10/2020	Hsu et al.
2020/0351231 A1	11/2020	Hsu et al.
2020/0351232 A1	11/2020	Hsu et al.

FOREIGN PATENT DOCUMENTS

WO	WO 2007/044468 A2	4/2007
WO	WO 2016/205764 A1	12/2016
WO	WO 2017/070605 A1	4/2017
WO	WO 2018/172556 A1	9/2018
WO	WO 2018/183403 A1	10/2018
WO	WO 2019/094969 A1	5/2019

OTHER PUBLICATIONS

Abudayyeh et al., "C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector," *Science* 353:aaf5573, 2016.

Abudayyeh et al., "Supplementary Material for C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector," *Science* 353:aaf5573, 2016.

Accession No. EM_GSS:ER054570, Mar. 16, 2007, XP055784868, retrieved from the Internet: RL:http://ibis.internal.epo.org/exam/dbfetch.jsp?id=EM_GSS:ER054570, retrieved on Mar. 11, 2021.

Accession No. GSN:BGJ29232, May 16, 2019, XP055784879, retrieved from the Internet: URL:http://ibis.inernal.epo.org/exam/dbfetch.jsp?id=GSN:BGJ29232, retrieved on Mar. 11, 2021.

Anantharaman et al., "Comprehensive analysis of the HEPN superfamily: identification of novel roles in intra-genomic conflicts, defense, pathogenesis and RNA processing," *Biol. Direct* 8:15, 2013.

(Continued)

Primary Examiner — Celine X Qian

(74) Attorney, Agent, or Firm — Klarquist Sparkman, LLP

(57) **ABSTRACT**

Provided herein are CRISPR/Cas methods and compositions for targeting RNA molecules, which can be used to detect, edit, or modify a target RNA.

30 Claims, 87 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

- Barrangou and Gersbach, "Expanding the CRISPR Toolbox: Targeting RNA with Cas13b," *Mol. Cell* 65:582-584, 2017.
- Bork, "Powers and Pitfalls in Sequence Analysis: The 70% Hurdle," *Genome Res.* 10:398-400, 2000.
- Brenner, "Errors in Genome Annotation," *Trends Genet.* 15:132-133, 1999.
- Broun et al., "Catalytic Plasticity of Fatty Acid Modification Enzymes Underlying Chemical Diversity of Plant Lipids," *Science* 282:1315-1317, 1998.
- Cebrian-Serrano and Davies, "CRISPR-Cas Orthologues and Variants: Optimizing the Repertoire, Specificity and Delivery of Genome Engineering Tools," *Mamm. Genome* 28:247-261, 2017.
- East-Seletsky et al., "RNA Targeting by Functionally Orthogonal Type VI-A CRISPR-Cas Enzymes," *Mol. Cell* 66:373-383, 2017.
- East-Seletsky et al., "Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA processing and RNA detection," *Nature* 538:270-273, 2016.
- Fonfara et al., "The CRISPR-associated DNA-cleaving enzyme Cpf1 also processes precursor CRISPR RNA," *Nature* 532:517-521, 2016.
- GenBank Accession No. HHH545617.1 "Ruminococcus bicirculans chromosome II, complete genome," Feb. 27, 2015.
- Gootenberg et al., "Nucleic acid detection with CRISPR-Cas13a/C2c2," *Science* 356:438-442, 2017.
- Grissa et al., "CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats," *Nucleic Acids Res.* 35:W52-W57, 2007.
- Kindsmüller et al., "Intranuclear Targeting and Nuclear Export of the Adenovirus E1B-55K Protein Are Regulated by SUMO 1 Conjugation," *Proc Natl Acad Sci.* 104:6684-6689, 2007.
- Konermann et al., "Transcriptome Engineering with RNA-Targeting Type VI-D CRISPR Effectors," *Cell* 173:1-12, 2018.
- Makarova et al., "An Updated Evolutionary Classification of CRISPR-Cas Systems," *Nat Rev Microbiol.* 13:722-736, 2015.
- Nelles et al., "Programmable RNA Tracking in Live Cells with CRISPR/Cas9," *Cell* 165:1-9, 2016.
- O'Connell et al., "Programmable RNA recognition and cleavage by CRISPR/Cas9," *Nature* 516:263-266, 2014.
- O'Connell, Mitchell R., "Molecular Mechanisms of RNA 11-Targeting by Cas13-containing Type VI CRISPR-Cas Systems," *J Mol Biol.* 431:66-87, 2018.
- Sapranaukas et al., "The *Streptococcus thermophilus* CRISPR/Cas system provides immunity in *Escherichia coli*," *Nucleic Acids Res.* 39:9275-9282, 2011.
- Shmakov et al., "Discovery and Functional Characterization of Diverse Class 2 CRISPR-Cas Systems," *Mol. Cell* 60:385-397, 2015.
- Shmakov et al., "Diversity and evolution of class 2 CRISPR-Cas systems," *Nat Rev Microbiol.* 15:169-182, 2017.
- Smargon et al., "Cas13b Is a Type VI-B CRISPR-Associated RNA-Guided RNase Differentially Regulated by Accessory Proteins Csx27 and Csx28," *Mol. Cell* 65:618-630, 2017.
- Smith et al., "The challenges of genome sequence annotation or 'the devil is in the details'," *Nat Biotechnol.* 15: 1222-1223, 1997.
- Stendahl, "Biotech Startup Co-Founded by MIT's Feng Zhang Unveils New CRISPR Tool," *Boston Business Journal*, <https://www.bizjournals.com/boston/news/2018/03/15/biotech-startup-co-founded-by-mit-s-feng-zhang.html>, accessed Mar. 15, 2018.
- Van de Loo et al., "An oleate 12-hydroxylase from *Ricinus communis* L. is a fatty acyl desaturase homolog," *Proc. Natl. Acad. Sci.* 92:6743-6747, 1995.
- Wegmann et al., "Complete genome of a new *Firmicutes* species belonging to the dominant human colonic microbiota ('Ruminococcus bicirculans') reveals two chromosomes and a selective capacity to utilize plant glucans", *Environ Microbiol.* 16:2879-2890, 2014.
- Wessels et al., "Massively Parallel Cas13 Screens Reveal Principles for Guide RNA Design," *Nat Biotechnol.* 38:722-727, 2020.
- Yan et al., "Cas13d Is a Compact RNA-Targeting Type VI CRISPR Effector Positively Modulated by a WYL-Domain-Containing Accessory Protein," *Mol. Cell* 70:327-339, 2018.
- Zhang et al., "Structural basis for the RNA-guided ribonuclease activity of CRISPR-Cas13d," *BioRxiv*, doi: <http://dx.doi.org/10.1101/314401>, published May 4, 2018 [retrieved from Internet on Oct. 12, 2018].
- Zhang et al., "Structural Basis for the RNA-Guided Ribonuclease Activity of CRISPR-Cas13d," *Cell* 175:212-223, 2018, with 4 pages of Supplemental Information.
- PCT/US2018/047597 International Search Report and Written Opinion dated Oct. 18, 2018 (14 pages).
- Accession No. GSN:BGJ29097, May 16, 2019, XP055816784, retrieved from the Internet: URL:<http://ibis.internal.epo.org/exam/dbfetch.jsp?id=GSN:BGJ29097>, retrieved on Jun. 22, 2021.
- Accession No. GSN:BGJ29098, May 16, 2019, XP055816789, retrieved from the Internet: URL:<http://ibis.internal.epo.org/exam/dbfetch.jsp?id=GSN:BGJ29098>, retrieved on Jun. 22, 2021.
- Accession No. GSN:BGJ29231, May 16, 2019, XP055816793, retrieved from the Internet: URL:<http://ibis.internal.epo.org/exam/dbfetch.jsp?id=GSN:BGJ29231>, retrieved on Jun. 22, 2021.

FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 3A





FIG. 3B






FIG. 3C




FIG. 3D

FIG. 4C




FIG. 4B




FIG. 4A




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 6F




FIG. 6E




FIG. 6G




FIG. 6H




FIG. 7A




FIG. 7B




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 8D




FIG. 9A



FIG. 9B




FIG. 9C




FIG. 9D




FIG. 9E




FIG. 9F




FIG. 9G




FIG. 9H




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 10D




FIG. 11A




FIG. 11B




FIG. 12A




FIG. 12B




FIG. 12C




FIG. 12D




FIG. 12E




FIG. 12F




FIG. 13




FIG. 14




FIG. 15A




FIG. 15B




FIG. 15C




FIG. 15D




FIG. 16A




FIG. 16B




FIG. 17A





FIG. 17 B



CLUSTAL O(1.2.4) multiple sequence alignment

```

037...emb|OIZA01000315.1|          -----MAKKRMTAKERKQHRESL-----MK      22
emb|OCTW011587266.1|          -----
k87...11092736                -----
BMZ...11E_GL0037915          -----
EMZ...11B_GL0037771          -----
EMZ...11B_GL0069617          -----
Ga0099364_10024192          -----
530373_GL0023589            -----
emb|ODAI011611274.1|          -----MKKISLKEQRNKKRAEN-----KL      20
EMG_10003641                -----
CasR_P1E0...metageno        -----
emb|OIZX01000427.1|          -----MAKKKTARQLREEMQQQR-----KQ      21
160582958_gene49834          -----MKNVVFVKLIQAQENKEA-----AR      20
gi|1198542314|gb|NFIR01000008.1| -----MAKLLRPKELREKRRMAE-----KE      20
emb|OCNF01009141.1          -----
Ga0129306_1000735          -----
MH0288_GL0082219            -----MKRMTAKERREQEKKQR-----AE      19
FIG-022_GL0026351          -----
FIG-028_GL0185479          -----
250twins...35838_GL0110300  -----MGNKQVSAQRRRENAKLCN-----QQ      22
FIG-014_GL0226364          -----
250twins...36050_GL0158985  -----MKRKHQSAAEKQVKKLKN-----QE      21

```

FIG. 18A

emb OJMM01002900.1MGKHLNAKQKLEKLN.....	QQ	20
emb OGPN01002610.1MAKKITAKQREKERLN.....	KQ	20
emb OHCP01000044.1MAKKITAKQREKERLN.....	KQ	20
PIG-046_GL0077813MAKKITAKQREERQN.....	KQ	20
piq_chimeraMAKKITAKQREERQN.....	KQ	20
emb CIEN01002196.1		0
O2_UC29-0_GL0096317		0
CasR_Anaerobic_dig		0
emb OGDF01008514.1		0
emb OGZC01000639.1MKKNIRATREALKQK.....	IK	19
Ga0224415_10048792_chimera		0
emb OCVV012889144.1		0
Ga0187910_10006931		0
R_flav_XPD...		0
Ga0224415_10007274		0
Ga0187911_10069260		0
Ga0187910_10015336MENVKNPKEIVKCAKVFEEIYENPKFKQKQ		31
OPAI_chimera	MLKKSVCCKKFEITNYGGKPVCRNCRSNKSRKPVNPKSRKPEISDK.....	AI	49
3109_GL0034153		0
DLF014_GL0011914_MAKKNAKQREERENL.....	KA	20
CasR_R_albusMAKKSQMSLREKRELEKQKR.....	IQ	23
Ga0187910_10040531MAKKKAKQREEQEAAE.....	MN	20
tpg DXD01000002.1		0
CasR_E_siraeumMGKKIHARDLREQRTDR.....	TE	20

FIG. 18B

PIG-018_GL0023397MAKKLSPRELREQREER.....QA	20
PIG-025_GL0099734MAKKNNAKEQRAAIQAEN.....NK	20
Ga0129317_1008067_chimera	0
EYZ-362B_GL0088915MKKLNIREQREAKKAAA.....KQ	20
uncultured_Ru...spMAKKNKMPRELREAQKAR.....QL	22
R_flav,_FD-1MKKMSLREKREAEKQAK.....KA	20
tpgIDBYI01000091.1.1MKKLIKARDLREAKKQEK.....LA	20
037_...emb OIZA01000315.1KADSNAEKKEKAKKPVVENK.....PDIATISKDNKPKPNKEIKKSSAK	64
emb OCTW011587266.1	18
k87_11092736	9
BMZ-11B_GL0037915	11
BMZ-11B_GL0037771	10
BMZ-11B_GL0069617	10
Ga0099364_10024192	10
530373_GL0023589	23
emb ODAI011611274.1KYQKAQAEBAAAQQIACGAASEENPCFDVVVDI...KPKALNPLHVEIEAPSAKKSSVK	77
EMG_10003641	30
CasR_P1E0_metageno	15
emb CIZX01000427.1AIQKQEQEQK.....ECAA.....AAR...ETAAPQPAAPVPPKQRKSLAK	61
160582958_gene49834KKAKDIAEQARIAKRNQVYKKEENRI...NRI...QIEIQTKKNTQNYHLKSLAK	72
gi 1198542314 gb INFIK01000008.1EHKKQFK.....LRKEQELRKKQEKQ.....RED.....QKELEKIKKEEGEGEKKSSGAK	68
emb OGNF01009141.1	12

FIG. 18C

Ga0129306_1000735MOK.....QREQQVTDESERKKKFLKSGAK	26
MH0288_GL0082219	V-----LFGKQEAN-----RKK---AQAATQDATAPVHHKNAKSLAK	54
PIG-022_GL0026351	-----MAESYKIKSKKTRIK	15
PIG-028_GL0185479	-----MSESDKIKIKSKKTRIK	17
250twins_35838_GL0110300	KARQAESQDKIKMNVK-----M-----KNINTNDIKHKITTKAK	58
PIG-014_GL0226364	-----MTKPEKKQIKQAITLK	16
250twins_36050_GL0158985	KAQKYASEPSPLOSDTAGV-----EC-----SOKKTVVSHIASSKIYAK	60
emb:IOJMM01002900.1:	KDMMYTKSDAVSVPTLKA-----APTKAEMSQDTAEASTLITPGTLKTKAK	67
emb:IOGPN01002610.1:	KWAKQDTPVPKSKTEKP-----V-----AASDDKLLKTTQVKKVVQIKSKAK	63
emb:IOHCF01000044.1:	KWAKNDSVIIVPEIKKIK-----TGEIQDNRRKRSRQKSOAK	58
PIG-046_GL0077813	KWAKKQDDATAVFECEANI-----KP-----ADSKDEDCANT--SIKRRKTQAK	62
pig_chimera	KWAKKQADATAVFECEADI-----KP-----ADSKDEDCINIYIKREKKKTQAK	64
emb:IOLEN01002196.1:	-----MERQKRKMKSKSK	13
O2_UC29-0_GL0096317	-----MERQKRKMKSKSK	13
CasR_Anaerobic_dig	-----MNNKPKTKAK	10
emb:IOGDF01008514.1:	-----MIETKPKREDIAKTPAAKSRSK	22
emb:IOGZC01000639.1	-----KRREELEKKNLACWEEITSAEGRRSRVK	65
Ga0224415_10048792_chimera	-----MSKKENRKSYYK	12
emb:IOCVV012889144.1:	-----MSKNKESYAK	10
Ga0187910_100006931	-----MIEKKKSYAK	10
R_Flav_XPD_	-----MIEKKKSPAK	10
Ga0224415_100007274	-----MIEKKKSYAK	10
Ga0187911_100069260	-----MSTKKEFRYSVAAK	14
Ga0187910_100153336	RFSAINVEVITDETHKEAKIQKNEFKITPDQFTKE-LQETQKVNGETKKEHITKNKHTNVK	90

FIG. 18D

ODAI_chimera	ELITTEKLIIDCKK.....NDCASECCGGRKRKSKAK	79
31009_GL0034153	-----MGRGVIIMTEKRSRAK	18
DLF014_GL0011914...	KIEFKIKYAEQEVTKI-----TEEIAQSDKIMKTESILVQHTKKS LAK	65
CasR_R,_albus	KAANVSVNDIPENTEAN-----VVSVNVRISAENKHSKKSAAK	62
Ga0187910_10040531	KIQSAVKAKAEIAPA-----VSSAFVEKRRKQSKKTFAK	55
tpg!DJXD01000002.1!	-----MKKQSKKTVSK	12
CasR_E,_siraenum	KFADQKKREAEKAVPKK-----DAAVSVKSVSSVSSKKDNVTKSMAK	63
FIG-018_GL0023397	KFAKQVQEREAEKAKREK-----QVSVSDTVGAAAAVQENEGKKSMAK	63
FIG-025_GL0099734	KIQMANAAREKQAEAEARLKAEAEERIEKENEKREAEELKRFKLDQETRGKLPKNERKSLAK	80
Ga0129317_1008067_chimera	-----MKKQKKS LVK	10
EYZ-362B_GL0088915	IYAAKASAAQKKADETV-----VAAQPAQDTANSKPIKSRNK	57
uncultured_Ru_sp	KAAEINNWAAPAI-----AMPAAEVIAPAAEKKS SVK	56
R_...flav,_PD-1	AYSAASKNTDSKFAEKKA-----ETPKPAEIIISDNSENKTVAK	58
tpg!DEYI01000091.1!	AFSAKANTVIYENEDK-----NVEAFPEALNLRSIKKS MNK	55
	*	
037_...emb!OIZA01000315.1!	LAGVKWIIKAN---DD-----VTYISSFGKGN-----NSVLEKRIIGDVSQDVN-----	105
emb!OCTW011587266.1!	AVGVKSLARLS---DG-----STVVSFGKGA-----AAELES LIYGGEIKLS-----	59
k87_11092736	RIGIKSTVAYG---QG-----KYAITTFKGS-----KAEI AVRSADPPPEELP TESD	54
BMZ--11B_GL0037915	RVGIKSVIAHD---GG-----KLA VTFKKS-----AAKLAYRDGDQTFDEKKSE	56
BMZ--11B_GL0037771	RMGVKALLAHG---ED-----KLIMTTFKGN-----RSKIEFTEGYHGRALET PKHF	55
BMZ--11B_GL0069617	RIGVKS AF SHG---NS-----RLTIITQFRGN-----TADIAVEIDSRGRSLSLPHRV	55
Ga0099364_10024192	RMGVKSVLAHG---KDEKGHIKLAIYAFKGN-----KAELAIQTDEKGSNLA KTYKE	60
530373_GL0023589	ANGLKSTFKLA---EN-----TYXMTSFGKNN-----IADPEKNIVRMVVENIR-----	64

FIG. 18E

emb ODAI011611274.1	ANGLKSLLLT-----DG-----KVTMTSFGSGS-----EANVEKRFEDETGTKTFD-----	117
EMG_10003641	MSGLKSALVIN--NH-----EMLLTSFGKGN-----NALAEKRYILDGDIETI-----N	72
CasR_P1E0_metageno	AAGLKSTFVISPQEK-----ELAMTAFGRGN-----DALLQKRIVDGVVRDVA-----	58
emb OIZX01000427.1	AAGLKSNFILDQRR-----TTVMTAFGGGS-----TALLEKQIVDRAISDLQ-----	104
160582958_gene49834	AAGKSVFAI---GN-----DLLMTSGPGN-----DATIEKRVFQNRALIEHL-----	112
gi 1198542314 gb NFIR01000008.1	ALGLKSTFILDREQ-----KMLMTSFGKGN-----KAVRDKYIIGDKVSDIDDSWE	115
emb OGNF01009141.1	AAGLKSTFVLE---NN-----KLLMTSFGNGN-----KAVIEKIIDEKVDSDIN-----	52
Ga0129306_1000735	AAGLKSVFVLS---EG-----KELLTSFGKGN-----EAVPEKRVTGGTIANAR-----T	68
MH0288_GL0082219	AAGLKSTFTLP---TG-----ELLMTSFGSGS-----EALIQKRIMKDVISDVQ-----	95
PIG-022_GL0026351	PAGLKSSFVN---EN-----KGLMNTAFGKGN-----NAVIEKRISGNLIENVS-----	55
PIG-028_GL0185479	PAGLKSSFVS---EG-----KGLMNTAFGKGN-----NAVIEKRISGDLIENVS-----	57
250twins_35838_GL0110300	KLGLKSTIIAD---K-----KIILTSFINEQ-----SSKTANIEKVAGFKGDILT---IS	103
PIG-014_GL0226364	RMGLKSTLVFD---N-----KITVTSFANSS-----NSEKASAHIEKRIDFWCKII---M	60
250twins_36050_GL0158985	AMGLKSTLVMG---D-----KLVITSFAASKAVGGAGYKSANIEKITDLQGRVI---E	107
emb OJMM01002900.1	AMGLKSTLVFD---D-----KIVVTSFLNSKI---EENEKCAHIEKIAADCNGQII-----V	112
emb OGPN01002610.1	AMGLKIVLSFD---D-----KIAIASFVNDK-----KIKLPHIERIIDKSGTII---H	105
emb OHCPO1000044.1	AMGLKAVLSFD---N-----KIAIASFVSSK-----NAKSSHIERIIDKEGTTI---S	100
PIG-046_GL0077813	AMGLKIVLGFQ---N-----KIAIASFTSSK-----DSKSSHIERIIDPNGKII-----R	104
pig_chimera	AMGLKIVLGFQ---N-----KIAIASFMSSK-----DSKSSHIERIIDPNGKII-----R	106
emb OIEN01002196.1	MAGVKSFFVIG---D-----ELLMTSFGDGD-----DAVLEKIDIDENGVVNDC---R	54
O2_UC29-0_GL0096317	MAGVKSFFVID---S-----DLLMTSFGERD-----KAVPEKYIAQGNVKSUR---K	54
CasR_Anaerobic_dig	AAGLKSVFFDQ---K-----QAVLITFAKGN-----NSQIEKVVNSEVVKDLR---Q	51
emb OGDF01008514.1	AAGLKSTFAVN---G-----SVLLNLSFGKGN-----DAVPEKLIITEKAVSEIN---T	63
emb OGZC01000639.1	AVGVKSFFVVG---D-----DLXLATFNGN-----EIVLEKKIITPDGKITTF---P	106

FIG. 18F

Ga0224415_10048792_chimera	GLGLKSTLVSD---S-----KVYLTFADGS-----NAKLEKCVENKIKICIS-----N	53
emb OCVV012889144.1	GMGLKSALVSG--S-----KVYMTSFEQGN-----DAKLEKVVENSEIVSLA-----	50
Ga0187910_10006931	GMGLKSTLVSD--S-----KVYMTSFGNGN-----DARLEKVVENNAISCLV----D	51
R_...flav_XPD...	GMGVKSTLVSG---S-----KVYMTYFAEGS-----DARLEKIVEGDSIRSVN-----	50
Ga0224415_10007274	GMGLKSTIVSG--S-----KVYMTTFDGS-----EARLEKVVENDSIKTLH-----	50
Ga0187911_10069260	AAGLKSSLAVD---ID-----RIVMTSFGHGN-----AAILEKEIVDGEISVLN----I	56
Ga0187910_10015336	AAGVKSVFVAVD---DG-----NVLITSFGRGN-----AADIEILKSDDDKTIINL-----F	132
ODAI_chimera	AAGLKSAIDLG--NG-----QALITSFGKGN-----EAVFEKRVEGDKIISLR-----E	121
31009_GL0034153	ALGVKSFAFLKD--D-----DVIMTSFGRGN-----DAVVEKIIKNNEVQIN----I	59
DLF014_GL0011914	ASGVKSVFVKD--D-----EVIMTSFGRGN-----DAVVEKIIKDNVDEK-----N	105
CasR_...albus	ALGLKSGLVIG--D-----ELYLTSFGRGN-----EAKLEKISGDIYVEKLG---I	103
Ga0187910_10040531	ASGLKSTLAVD--N-----SAVMIVFGRGN-----EAKLDHRINADLQSESL-----H	96
tpg DUXD01000002.1	TSCLKEALSVQ--G-----TVIMTSFGKGN-----MANLSYKIPSSQKFNL-----N	53
CasR_E_...siraenum	AAGVKSVFVAVG--N-----TVYMTSFGNGN-----DAVLEQKIVDTSHEPLN----I	104
FIG-018_GL0023397	AAGVKSVFVAVG--D-----IVYMTSFGNGN-----DAVLEKKIVDESHESLN----K	104
FIG-025_GL0099734	AAGLKSTFAVG---T-----DLYMTSFGKGN-----DAIVEKKSINKKQVTCIN-----T	121
Ga0129317_1008067_chimera	AAGLKSAFVVG---D-----SVYLTSFGKGN-----AARLDTKINPDNSTERY-----V	51
EYZ-362B_GL0088915	AAGLKSTFVVG--D-----SLYMTSFGKGN-----LSVPEYRVDIGDAYEVD---P	98
uncultured_Ru_sp	AAGMKSILVSE--N-----KMYIITSFGKGN-----SAVLEYEVDNNDYNGIQ---L	97
R_...flav_FD-1	AAGLKSTIIISG--D-----KLYMTSFGKGN-----AAVIEQKIDINDYSFSA-----	98
tpg DBYI01000091.1	AAGLKSTLIIDG--K-----SLYLTAFGKGN-----NAVVERHMIATDDSYSLK---T	96
	; : *	
037_...emb OIZA01000315.1	---KDSHMYVNPKYTKKNEYIKNGF-----SSGSSLVAVY	136

FIG. 18G

```

emb|OCTW011587266.1|
k87_11092736
EMZ-11B_GL0037915
BMZ-11E_GL0037771
BMZ-11E_GL0069617
Ga0099364_10024192
530373_GL0023589
emb|ODAI011611274.1|
EMG_10003641
CasR_P1E0_metagene
emb|OIZX01000427.1|
160582958_gene49834
gi|1198542314|gb|NFIR01000008.1|
emb|CGNF01009141.1
Ga0129306_1000735
MH0288_GL0082219
PIG-022_GL0026351
FIG-028_GL0185479
250twins_35838_GL0110300
PIG-014_GL0226364
250twins_36050_GL0158985
emb|OJMM01002900.1|
emb|OGPN01002610.1|
emb|OHCPO1000044.1|
-----D--KAILEITDDTQNKMYNVKS-----SRIPNLTAR      89
ATLSIHAKFAKAGRDGREFKCGVD-----EIRI-----HTSR      88
-----LGGFVTVGVDENTVDAERGI-----FEGGLEASVAN      88
G-----KRGFEVRKIDENVLDYGLD-----EEGK      79
-----EKSFVVKIDDEIDVQRS-----
-----RNITANKIVSEGIQTSGTI-----AGE      82
-----NIFSATFISERIVDITGEY-----
-----RDFELFSAKLELTGYRIQRFNA-----S-----PKDAG-----  145
-----NKNKFDANNDKVVVVKGISN-----PNGQLIN      101
-----GEKQOFQVQRFQDESFRRLQNSR-----LADRIVTADD      90
-----PVOQFQVEFASAAKYRLKNSR-----VRFNVTADD      135
-----SSPEQYSAEFQNKQFKIKGNIK-----VLNHS-IQKMEELIQTE  149
-----NKAALSVEVCGKSFNISKKEN-----DDCEPVKVVNN      147
-----EPEVFSVTECDKKFELQPAKR-----GLAADSIVDN      83
-----DNKEAFSAALQNKRFVEVFG-----TAGSSDD      95
-----EGDPAFSVLFPFKQFEIQGRAI-----DVQKEGSSVPEVLTDD  133
-----NKPRFSVEFGNKEYIINDK-----KLGIVSDD      82
-----NKPRFSVEFGNKEYIINDK-----KLGIVSDD      84
-----YTPRMFRSEINPGEIVISKGD-----DLSEFAN      131
-----QDEKMFRIYKQIYIEKEFFDENNKII-----I-----QIAMPEN  97
-----EHERMFSADVGEKNIELSKND-----CHTNVNN      135
-----ERPRMNTISNAKKVDLSKDN-----DEINYPN      140
-----ENARMPDSSVDEQNVNIEKRMVIEEKQND-----GTFKKDKVKATICN  150
-----VNSEMFESSVNRKRDINIEKRITIEEPQOD-----GTIKKEKGVKSTTCN  145

```

FIG. 18H

FIG-046_GL0077813EDVRMFDNVDECSINLEKRMVVEERQKD.....GTIKKDEKDVKSTICN	149
pig_chimera	----EDVRMFDNVDECSINLEKRMVVEERQKD----	----GTIKKDEKDVKSTICN	151
emb!O1EN01002196.1!	---N-PAAYDAVYGTDSIRVKKTN-----	---NNIRAKVNN	83
O2_UC29-0_GL0096317Q-EQAFDAQYKKEKIFVTNAD.....NKLHVAINN	83
CasR_Anaerobic_dig	---P--PAFDLELKEKTFYISGKNINITSRENPLASASLPLSKRQRIARAERAKARENR		106
emb!OGDF01008514.1!V-KFRFSVEKAPATSYSSSF.....GIKSHISATADN	93
emb!OGZC01000639.1E-EEFTTAAIKLFAQTEPTVATSI.....GISNGRIYVPEISVDN	144
Ga0224415_10048792_chimera	----D-KEAFAASIAKNKVGKIKND-----	----EKFRHPKGYDIISNN	89
emb!OCVV012889144.1!	----E-KESFSAEIEFKKNIGCKIE-N-----	----KKFKHPKPYDVIADN	85
Ga0187910_10006931	----K-KEAFVAEITDKNAGYKII-N-----	----KKFGHPKGYDVVANN	86
R_...flav_XPD_	---E-GEAFSAEMADKNAGYKIG-N-----	---AKFSHPKGYAVVANN	85
Ga0224415_10007274E-GEAFSAEMDNNAGYKIG-N.....VKFSHPNGYDVVANN	85
Ga0187911_10069260	----E-NPAFDAVIND----KKYALTG-----	----HH-----AGVHALVDO	85
Ga0187910_10015336	----ETENQ-----KKYVVTN-----	----K-----RSNVKGLADN	155
ODAI_chimera	----G---EAFALAKEPVMHKLVLKSG-----	----RIKAEQGFVEFDG	154
31009_GL0034153	----D-KPVYDVVGVKEKTEIGIINWVG----	----TKRGINTINKKQYAN	95
DLE014_GL0011914_T-KPAYEIVDIEKNGDLIKVQ.....SRRCKNIESAN	136
CasR_R_...albusG---AFVVAERD---ESTLILES.....GRI---KDKTARPKD	132
Ga0187910_10040531	----PQAALKN--VHAPNKQKIHFIG-----	----RM-----QDMNLTADH	127
tpg!DJXD01000002.1!SSAGLKN---VEVS--GKKIKFOG-----R-----HPKIATTDN	82
CasR_E_...siraecum	----D-----DPAYQINVVVTMNGYSVIG.....	----HR---G---ETVSAVIDN	135
FIG-018_GL0023397	----A---NPAYEITDISSVNVYSVEG-----	----HR--G--QSVCATADN	135
FIG-025_GL0099734E-KENFTVDDGITDIIVPVQS-----	----KR---I---ACUNIQADN	154
Ga0129317_1008067_chimera	----SDSEKHTLKINSITDTELRUSG-----	----P-----FFKQAEAKN	83

FIG. 18I

EYZ--362E_GL0088915	-----IS--EHSELHVLGADDLAVSFG-----TRRMK-----DVSELKADN	133
uncultured_Rb_sp	----SSKDNSNIQLGGVNEVNIYFSS-----KHGFE--SGVEINTSN	133
R...flav...FD-1	---M-KDTPSLEVDKAESKEISFSS-----HFFV--KNDKLITYN	133
tpg:DEYI01000091.1:	-----L--ENEPSLKVRAADELKVTFMS-----RRPFV---QESEL SAVN	131
037...__emb OIZA01000315.1	PNK-----PDKNSGMDALCLKPYFEIDFFG-----HIF--TDNMRICAIYNIID	178
emb OCTW011587266.1:	TDK-----LSDKSGMDDLGEKRELELEVFG-----CCF---DDSIHQIAHAVFD	131
k87_11092736	EYES-----LISNPAESPREDYLGKGLKGTLEKRFPG-----DEY--PKDNLRIQIYYSILD	136
BMZ--11E_GL0037915	PVND-----LIDESKKEYGQDYLMLKDKOLEKVFVG-----QTF--PGDNLRIQIYNIILD	136
BMZ--11B_GL0037771	TIEA-----LLINPSEKVEDYLGKGLKGTLEKRFPG-----REF--PHDNIRIQLIYNIILD	127
BMZ--11B_GL0069617	SLLG-----TLINPAENDPVDYLGKGLKSTLEKEFFG-----QEF--FNDSIRIQLIHNILD	121
Ga0099364_10024192	CHAT-----FLNPAEHVCTDYLKLEKETELEMEFFG-----KSF--PGDSVRIQIHHQILD	130
530373_GL0023589	-----KKTACVLPACSDNQLHAKDTIEQMYFG-----KTY---TDNIHQIAYNIMD	128
emb ODAI011611274.1:	-----LAYRPAGVRPDQIGAKAALKRYFG-----KET--FGDNIHVQIAYQIQD	188
EMG...10003641	PLFD-----QSPTAIQPNRTSGNDMIGIRRMLEKRYFVHNEENKEF---QDNIRIQLIAYCILD	156
CasR_P1E0_metageno	PLHR-----A-ETPRQPLGAGMDQLRKAILEQKYFG-----RTF--DDNIHQIQLIYNIILD	139
emb OIZX01000427.1:	PLYR-----R--KD---CGFVPGMDALRRKNVLEQRFFG-----KSF---ADNIHQIHYYSILD	181
160582958_gene49834	LQD-----NYNRPHFDDLGCNKVLEQKYFG-----RTF--SDNIHVQIAYNIMD	191
gi 1198542314 gb NFIR01000008.1	PVL-----SGNKKDDDLIHCKPKNLEEMYFG-----CQF--KDNHQIQLIYNIILD	189
emb OGNF01009141.1	PLK-----SKKTAGDDAIHSKFLERQFFDG-----NTF---NDNIHQIQLIYNIILD	126
Ga0129306_1000735	PLA-----VSRAPGQDLIGAKTALEERYFG-----PAF--ADNIHQIYIYAIQD	137
MH0288_GL0082219	PRA-----SQKPAGADLIGAKPALEKRYFG-----KTF--TDNIHQIQLIYNIILD	175
PIG-022_GL0026351	PRE-----SQKPVGEDSIGHCKRALEERYFG-----KTF--NDNIHQIQLIYYSILD	124

FIG. 18J

```

FIG-028_GL0185479      PRE-----SQPVGEDGHCRALEERYFG-----KTF--NDNIHQIYIYSILD      126
250twins_35938_GL0110300  PAN-----FPIGRDYVXIRSALEKQYFG-----KEF--PEDNLHVQIAYNVAD      172
FIG-014_GL0226364      PAL-----RNCCKDYIGMKNSEKVMFG-----KEF--PNDNLHICAIYAINILD      138
250twins_36050_GL0158985  PVV-----INICKDYIGIKSRLECEFFG-----KTF--ENDNLHVQIAYNILD      176
emb!OJMM01002900.1:    PAF-----EDCGRDYINVSALAEKRVFG-----KTY--NKDNLHVQIAYNIFD      181
emb!OGPN01002610.1:    PYP-----KTCCKDYIGIKDVAEKYFFG-----KTF--FENLNRVQIAYNVFD      191
emb!OICP01000044.1:    PYP-----KVGCKDYIGIKEIAEEHFFG-----RAF--PNEHLRVQIAYNIFD      186
FIG-046_GL0077813      PYS-----NECGKDYIGIKSVAEELFFG-----RTF--PNDNLRVQIAYNIFD      190
pig_chimera            PYS-----NECGKDYIGIKSVAEELFFG-----RTF--PNDNLRVQIAYNIFD      192
emb!OIEN01002196.1:    PLAKSNIRSESALEFRTRVNEYKREQDKYETLFFG-----KTF--DDNIHQIYSKILD      136
O2_UC29-0_GL0096317    PLCGNKKQSPAEQYSRRVLEHKKAINQKYETIIVFG-----KPS--NDNIHQVVSKIMD      136
CasR_Anaerobic_dig     PYHNV-----KRVGEDDLRAKADLEKHFFG-----KEY----SDNLKIQIYYNILD      148
emb!OGDF01008514.1:    PLAGR-----APVGEDAIHAKEVLEQRVFG-----KTF--SDDNIHQIYYNILD      136
emb!OGZC01000639.1    PLHTT--MQK--NTIKRSAGEDILQLKDVLENRYFD-----RSF--RDDLHRLIYYNILD      193
Ga0224415_10048792_chimera  PLLHN-----NSVQQDMLGLKNVLEKRYFG-----KSSGGDNNLCIQIHHNILD      133
emb!OCVV012889144.1:    PLYK-----GSVRQDMLGLKETLEKRYFN-----SAD--GTDNVCIQVIHNNILD      127
Ga0187910_10006931    PLYT-----GPVQDMLGLKETLEKRYFG-----SSVSGDNDNICIQVIHNNILD      129
R_...flav_XFD_        PLYT-----GPVQDMLGLKETLEKRYFG-----ESADGNDNICIQVIHNNILD      128
Ga0224415_10007274    PFYTT-----GPVQDMLGLKEILELERRYFG-----SSTGDNNTICIQIHHNILD      128
Ga0187911_10069260    PQN-----RSDAVHIRGALEKKYFG-----DTF--ADNIHVQIAYNILD      122
Ga0187910_10015336    PTKV-----ESIIPGETIQIGFSILEKHFFG-----RTF--NDNIHQIYYNILD      198
ODAI_chimera          POKNS-----SMSAVRLDEYNNQIERKFFRGN--KVL--QDNIRIQIAHNIMD      198
31009_GL0034153      TQV-----SQKRMGMDLIQRKDKLEKVFVN-----ENF--NDNIHQIYYNILD      138
DLF014_GL0011914...   TEL-----PPEKIGMDLIRCKDKLEIYFG-----RTF--NDNIHQIYYNILD      176

```

FIG. 18K

CasR_R_...albus	PRHI-----T----VDTQKFKEDMLGIRSVLEKKIFG-----KTF---DDNIHVQLAYNILD	178
Ga0187910_10040531	PLHSH-----DGERAVGADLLCAKDKLEQLYFG-----RIF---NDNIHQIQLIYQILD	172
tpg!DJXD01000002.1	PLFK-----PQFGMDLLCLKDKLEMHYFG-----KTF---DDNIHQIQLIYQILD	123
CasR_E_...siraenum	PLRRFNGRKK---DEPEOSVPTDMLCKPTELEKFFG-----KEF---DDNIHQIQLIYNILD	186
FIG-018_GL0023397	PLHHD-----GGSSVPTDMLCKPTELEKFFG-----KTF---DDNLHQIQLIYNILD	179
FIG-025_GL0099734	PLYRK-----DSENQVQPKLLKDKTLEKVFYFG-----KTF---NDTLHQIQLIYNILD	199
Ga0129317_1008067_chimera	PTHK-----K---DNEQKNTQDMLGKSTLEKFFG-----STF---DDNIRIQIHHNIQD	129
EYZ-362B_GL0088915	PLHSG-----K---NNLVPKNGTDILGKLGILEQRYFG-----RNF---DDNVHQIQLIYNILD	180
uncultured_Ru_sp	PTHRS-----GESSFVRGDMGLKSELEKRFYFG-----KTF---DDNIHQIQLIYNILD	178
R_...flav_...FD-1	PLYGG-----K---DNPEKPVGRDMLGKDKLEKRYFG-----CTF---NDNLRIQIQLIYNILD	180
tpg!DEYI01000091.1	PLHSG-----K---DSPNKSAGQDMLGKSELEKRYFG-----KIF---DDNLHQIQLIHHNILD	178
	. * : ; : : :	
037_...__emb!OIZA01000315.1	IEKILAKHITWIIYTVNSFDRNYNQ-----SGNDITIGF-----DLNYRV	217
emb!OCTW011587266.1	IQKSLAAVIPNVLYTLNNLDRSYST-----DNTSDRKDIIGN-----TLNYQH	174
k87_...11092736	IQKILGLYVEDILHFVDGLQDEP-----EDLVGL-----GLG---D	159
BMZ--11B_GL0037915	MQKAFGLYVNDVLYAVNNLQNTGRL-----LIDDDWTSDEVIGL-----ALG---K	178
BMZ--11B_GL0037771	IYKILGMNVADILYALGNMQDIE-----LIDDMFGQ-----SLN---N	162
BMZ--11B_GL0069617	IQKILGIYINDVLYSIANLQNVF-----YVEDCLGQ-----SLS---K	156
Ga0099364_10024192	IQKLLGIYIITDIIYICINNLREDETHL-----DHESDIVGL-----SMS---N	168
530373_GL0023589	MKKIFGVYSNNIIYSINNLLEE-----REGSDFLGV-----FYTEN	164
emb!ODAI011611274.1	IEKLLAVYIISNIIYAVNNVTGVSAM-----KDSKGRFVDLLGD-----YGIIGEE	233
EMG_10003641	IEKILMPHINNICFEINMLRL---E-----GYQELSEFMGS-----FNIYK	194
CasR_P1E0_metageno	IHKMLAVFANHIVHTLNLGGY-----GETDFVGM-----LPAGL	174

FIG. 18L

emb OIZX01000427.1	IHKILAAASGHIVHLLNIVNGS-----KDRDFIGM-----LAAHV	216
160582958...gene49834	IEKLLTFYINNIYYTLNELMRD-----NSKDDFFGCDSHFSVAYLYDELKA	237
gi 1198542314 gb NFIR01000008.1	IEKILAVQINNVIFILNNLRW-S-----GEEEDLIGS-----LGVNH	227
emb OGNF01009141.1	IEKILSVHVDIVYSVNNILSRGE-----GMEYNDYIGT-----L---NLK	164
Ga0129306...1000735	INKILAVHANNIVYTLNNDRE-A-----DPEIDDFIGS-----GYLTLKN	177
MH0288_GL0082219	IKKVLAIHANNIVYILDNLSRR-E-----GTEGIDFIGI-----GGLTLKN	215
FIG--022_GL0026351	LEKILAVHSNNIVYTFDNLRRK-E-----TDEYDDFIGY-----MSTVN	152
FIG--028_GL0185479	LEKILAVHSNNIVYTFDNLRRK-E-----TDEYDDFIGY-----MSTVN	164
250twins_35838_GL0110300	IKKILSVYINNIYMFYFNLRSEEDIFYNSQSENSGRDCDVIG-----SLYYQA	222
FIG--014_GL0226364	IKKIICPYINNVNIFYNLGRGKLG-----DDKNDLLG-----KQIGP--	176
250twins_36050_GL0158985	IKKILGTIVVNNIYIFYNLNRAGTG-----RDERMYDDLIG-----TLYAYK	218
emb OJMM01002900.1	IKKIIGAYINNIYIFYNLGRREEYD-----AKKDIIG-----TQ-----	215
emb OGPN01002610.1	IQKILGTIVVNNIYISFYFNLRDGG-K-----SDVDIIG-----SLYAFA	228
emb OHCFO1000044.1	VQKILGTIVVNNIYISFYFNLSRDEVQ-----SDNDVIG-----MLYSIS	224
FIG--046_GL0077813	IQKIFGTIYINNIYISFYFNLSRDEFQ-----SDNDVIG-----TLYMLK	228
pig...chimera	IQKILGTIYINNIYISFYFNLSRDESQ-----SDNDVIG-----TLYMLK	230
emb OIEN01002196.1	IEKTFSVVIGNIVYAINNLSLEQS-----IDRPIDIFGD-----KNTQ-----	174
O2_UC29--0_GL0096317	IEKTFSFVMGNIVYAINNLDPEQN-----IDNPIDYMSN-----KART-----	174
CasR...Anaerobic...dig	INKIISPYINDIVYSMNNLARNDY-----IDGKIDVIG-----SLSSTT	188
emb OGDF01008514.1	IRKILSTYANNVYFTINSMRRLDE-----YDREQDYLG-----YLYTGN	175
emb OGZC01000639.1	IEKILAEYTTNAVFAIDNVSGCS-----DDFLS-----NFSTRN	227
Ga0224415...10048792...chimera	IEKILSEYIIPNVVYAFNFIAGFKDE-----HNNIIDIIGT-----QTYNSSY	175
emb OCVV012889144.1	IEKILAEYITNAVYFDNIAGFG-----EDIIGM-----GGFKPIY	163
Ga0187910...10006931	IEKILAEYITNAAYAVNNIAGLD-----KDIIGF-----GKFSTVY	165

FIG. 18M

R_...flav_XPD_	IEKILAEYITNAAYAVANNISGLD.....KDIIGF.....GKFSYVY	164
Ga02244415_10007274	IEKILAEYITNAAYATNNIIDPD.....NDVIGG.....KKFYSIK	164
Ga0187911_10069260	ITKILTYANNVYALNNLVHADD.....T-QADELDSL.....NFSAGI	163
Ga0187910_10015336	IEKILAVHNNIVYALDNIHERGRE.....NSAEKPIDMIGA.....GGISTSK	242
ODAI...chimera	IEKILAVHNNIVYSVNLFVQG.....NEGDFLIG.....YRGLDK	236
31009_GL0034153	IEKILSVYVNNIVYALGNLERDNS.....DDEKDLIG.....YASILN	176
DLF014_GL0011914_	IEKILSVYINNAVYALGNLERKEN.....DKKEDLVG.....YLSARA	216
CasR_R_...albus	VEKIMAQVSDIVYMLHNTDKTER.....NDNLMG.....YMSIRN	214
Ga0187910_10040531	IQKILALHANNIFALDNLHKKK.....DELSDFVGM.....GMRATI	213
tpg!DUXD01000002.1	IEKILAVHVNNIVFTLDNVLHFQK.....EELTEDFIGA.....GGWRINL	164
CasR_E_...siraem	IEKILAVYSTNAVYALNMSADEN.....IENSEFFM.....KRTIDE	224
FIG-018_GL0023397	IEKILAIYSTNAVYALNNMIADED.....DESDLPQ.....NMTIDY	217
FIG-025_GL0099734	IQKILTVYSVNTVYTNLNFCKEI.....SERGDFIG.....ALSFOQ	237
Ga0129317_1008067_...chimera	IAKILAAHSNNAGYALDNMLAYQG.....VESEDMIG.....YMGTSR	167
EYZ-362B_GL0088915	IEKNLAIYVTVVAAALDHMCDEIL.....DTEHEDFIG.....YMNIMN	219
uncultured_Pu...sp	IEKILAVYVTVINIVYALNMLGVKG.....SESHDFIG.....YLSINN	217
R_...flav,_FD-1	IEKILAVHSANITATLDHMVDEDE.....KYLNSDYIG.....YMNITIN	220
tpg!DBYI01000091.1	IEKILAVYATNITAAALDHMVDDNE.....QYLGDFIG.....YMNILN	218
	: * :	
	: * :	
037_...emb O1ZA01000315.1	PLYEYGGKDSNGK.....PNKISAWKKRENFINFYN.....EAKPHLGY--	257
emb OCTW011587266.1	SYESFNVE.....KRGFEITYYN.....AAKDRFSY--	200
k87_11092736EKMQKLLS.....KALPYMGF--	185
BMZ-11B_GL0037915GRGKEIIE.....SMSDALGF--	194

FIG. 18N

BMZ-11B_GL0037771P..EKANDVQ.....RMKPYMGF..	179
BMZ-11B_GL00969617GQDEKLLKILN.....SIFYLGF..	175
Ga0099364_10024192INVLALN.....QMFYFYG..	184
530373_GL0023589IKSKLLTKNSKYLNDMNVWF-NK-EVSELSFRKVV..	211
emb ODAI011611274.1Q.....ADEEKALCAFL.....CSERLSY-	264
EMG_10003641KE.....SSRRDNFAKLM.....TSKQVRY..	225
CasR_P1E0_metagenoNGDY.....VDIKADIAAYA.....KRPQLAY..	207
emb OIZX01000427.1KRSIADFC.....KSPRLIY..	240
160582958_gene49834LS.....KNIDRIWNNFCNYMN-SDSGNTEARLAY..	277
gi 1198542314 gb NFIR01000008.1NYGKFSLEI.....KQSQMRY..	253
emb CGNF01009141.1NKKKEDLDRVK.....KIPQLAY..	192
Ga0129306_1000735EREKVTVSKQHFDAFM.....QNERLAY..	214
MH0288_GL0082219FDEGKEKNIENIEKSHGSELEYM.....KNPRLAY..	257
PIG-022_GL0026351L-FGKDQSVYDNIINRSRETFLEYR.....KNPRLGY..	203
FIG-028_GL0185479L-PGKEQSVYDNIINRSQETLEYR.....KNERLGY..	205
250twins_35838_GL0110300KD.....GKKKAIDSLID.....DTRAYTY..	254
PIG-014_GL0226364KSEEMVEDLLK.....NSSAYFVY..	195
250twins_36050_GL0158985GDKDMRRFRFEVKQLIQ.....NTSAYVVY..	253
emb OJMM01002900.1DSAVISKILN.....NTSAYFTY..	233
emb GGPN01002610.1PAFREAKDLLK.....NTEAYFSY..	256
emb OHCPO1000044.1ETFLQAKSLIK.....QTEAYYAY..	251
PIG-046_GL0077813DIFRQAFALLE.....RTEAYYSY..	255
pig_chimeraDIFRQARALLE.....RTEAYYSY..	257
emb OIEN01002196.1GISLRENDNYLK.....TMLPRCEYL	195

FIG. 180

O2.UC29-0_GL0096317DENLK.....RVLRRMYL	188
CasR_Anaerobic_digLEKEKFSFHRENKKFVE.....ASKPYMRY-	226
emb!OGDF01008514.1	SYERLLDIADYAVDGEDWRNTAAGISNDFEKKQFCITINGEFDLLD-----MIEFYMCY-	229
emb!CGZC01000639.1	QWDEFONPEQH.....REHF.....GNKDNVICSVKKQOQLFFNFFK.....NNRIGY--	270
Ga0224415_10048792_chimera	TYADFSKDKSD-----K-----KYIEFQKLLK-----NKRLGY-	203
emb!OCVV012889144.1	TYKQFKEPDK.....YKKFDDILN.....NSRLGY--	189
Ga0187910_10006931	TFDEFAEPDRH.....KERFIKDGKLDIKLINQLKNQYDEEFAFLD.....DTEFGY--	212
R_...flav_XPD_	TYDEFKDPHEH-----RAAFN-----NNDKCLINAIKAQYDEEFDNFDL-----NPRLGY--	207
Ga0224415_10007274	TFAQFSASD.....SSNDFEQFLK-----NPRLGY--	189
Ga0187910_10069260	SYAKSKSKSKS.....KQDFVELFI-----KKKEIHGY--	192
Ga0187910_10015336	EYEQYCSSEKSD-----YEDNFKQLI-----NNERLAIY-	270
ODAI_chimera	NYSDFCKNKPE.....DREFFNFNFKGDKGKVTNPRLSY--	270
31009_GL0034153	SYEDFINKNL.....VKQENIDKFI-----NGNRLGY--	205
DLF014_GL0011914_	GYNKVINKAIN-----DEKIKDRIEDLKEFIE-----NGNRLGY--	250
CasR_R_...albus	TYKIFCDTSN.....LFDDTKQKVENQKREFFDKIIK-----SGRLGY--	251
Ga0187910_10040531	GYDAERNSTNQ.....KVQETIYBEFQEFFVR-----PKELLY--	244
tpg!DJXD01000002.1	DYQTLRGQIN.....KYDFEKNYIK-----PKELLY--	190
CasR_E_...siraeum	TFDDFEKKKES.....T.....NSREKADPDAPEKFIG-----NYRLAY--	258
FIG-016_GL0023397	TYDKFRGRSE-----RIVKSFNDFVS-----NERLAY--	244
FIG-025_GL0099734	TYDDYTSDSKN.....KEIEEDFYS-----KPSLGY--	263
Ga0129317_1008067_chimera	TFDNYDFNHKN.....NKDFFRFLK-----LPRLGY--	193
EYZ-362E_GL0088915	SFDVFMHPSQS-----SRADIYKKINDSRPEENLLR-----TKRLGY--	257
uncultured_Ru_sp	IYDVFIDPDN.....S.....SLSDDKKANVRKSLSKFNALLK-----TKRLGY--	256
R_...flav_...FD-i	TYDVFMDFSKN.....S.....SLSPKDRKNIDNSRAKPEKLLS-----TKRLGY--	260

FIG. 18P

tpg DBYI01000091.1	TYEVFMEPSKN.....P.....RLDSNARKNIENSREFEYLLD.....TORLGY...	259
037_...__emb OIZA01000315.1	YENIFYDHG.....	266
emb OCTW011587266.1	FPDILCVLEKVNKDR.....	216
k87_11092736	FGSDVFKVTKKRE.....	199
EMZ-11B__GL0037915	FGSAFKPIPKKPGDKTSGKYAGGLKER.....	222
BMZ-11B__GL0037771	FGGIFRTQKKEDKSDNSLSEEEKEKKK.....	209
BMZ-11B__GL0069617	LGAGFKQSERPHQKNSRDPDPROKIE.....	201
Ga0099364_10024192	FGFAFRPVGDDVKELIILSDEVKRNIEKILALEEQKRNFPSTPRFKQENINLEIENAMGKF	244
530373__GL0023589	YSRSYANKISTAKKICICLOEDHFHYA.....TVKKAADQFTLLKRIORS	258
emb ODAI011611274.1	FGKEFCLVRNPKQP.....	279
EMG_10003641	LGNALYSDSLNLTKDE.....	242
CasR_FIEG_metageno	LGAAFYDVIKSKSR.....	222
emb OIZX01000427.1	YSAAFYETLDNGK.....	253
160582958_gene49834	FGELFYKPKETGDAK.....S	293
gi 1198542314 gb NFIR01000008.1	FGSTFCIFNENERITS.....	270
emb OGNF01009141.1	FGSAFYNTPEDISAK.....	207
Ga0129306_1000735	YGNAPFRKLSKAFFLARG.....REIFDK.....ESPERRQEILGSRGKN	254
MH0288__GL0082219	YGEAFYRRLSNERSKLGMEYERMSDA.....QRDAFORKYPIDFCDG	301
PIG-022__GL0026351	FGKVIFYEKVIDPRNK.....	218
PIG-028__GL0185479	FGKVIFYEKSTDAKDK.....	220
250twins_35838_GL0110300	FDGLFVSPKREDDGKI.....	270
PIG-014__GL0226364	FGDVFKQVKLSKEQOE.....	211

FIG. 18Q

250twins_36050_GL0158985	YGTLEKVKAKSKEQ	269
emb!OJMM01002900.1!	FDGVFKQITDRSNK	248
emb!OGPN01002610.1!	FGDVFKSKKGGKDE	271
emb!OHCFO1000044.1!	FDDVFKKKNKPKNKE	267
PIG-046_GL0077813	FDNVFKKIDKKNRN	270
pig_chimera	FDNVFKKIDKKNKKS	272
emb!OIEN01002196.1!	FHNILNSDSDNNSKMN	211
O2_UC29-0_GL0096317	FDLGFEYAEPSPYKKS	204
CasR_Anaerobic_dig	YGVVFIRDVKKSLST	242
emb!OGDF01008514.1!	FSEAFFCETTVKPPDS	245
emb!OGZC01000639.1	FGKAFTHAESE	281
Ga0224415_10048792_chimera	WGKAPFTGQGN	214
emb!OCVV012889144.1!	YCKAFFEKNDLHNPN	205
Ga0187910_10006931	FGKAFCKEGD	223
R_flav_XFD	FGQAPFSKEGR	218
Ga0224415_10007274	LCKAFFYKDGK	200
Ga0187911_10069260	FGDTFAFLDKRI	204
Ga0187910_10015336	FGNAFFKDEGNK	282
ODAI_chimera	YGAAFAEKVQFT	282
31009_GL0034153	FGNVFYKNKNE	216
DLF014_GL0011914	FGDIFFKKEEVKNCK	266
CasR_R_albus	FGFAFMVNSGNS	263
Ga0187910_10040531	FGSAFYNGDI	254
tpg!DJXD01000002.1!	FGFAFYHENE	200

FIG. 18R

CasR_E_sitaeum	273
PIG-018_GL0023397	256
FIG-025_GL0099734	273
Ga0129317_1008067_chimera	205
EYZ-362B_GL0088915	267
uncultured_Ru_sp	268
R_flav_FD-1	273
tpg!DEYI01000091.1	271
037__emb!OIZA01000315.1	294
emb!OCTW011587266.1	251
k87__11092736	229
BMZ-11B_GL0037915	253
BMZ-11B_GL0037771	249
BMZ-11B_GL0069617	228
Ga0099364_10024192	286
530373_GL0023589	298
emb!ODAI011611274.1	312
EMG_10003641	275
CasR_PLE0_metageno	261
emb!OIZX01000427.1	289
160582958_gene49834	339
gi 1198542314 gb NFIR01000008.1	304

FIG. 18S

emb!OGNF01009141.1ITTKIKSNEEIIYYTFMLLSTARNSAHYLDNRNRAKS.....SDAEDFDGCT	253
Ga0129306...1000735	KSVDDAIRALAFWVFRERDVYSELVIMSELRQSCFHGQOK-----N	297
MH0288...GL0082219	ELHFEIEKLAPEYVRKSEKEIYYILAMLSELRCQCFVHDSAA-----S	344
PIG--022...GL0026351	-----SKKMRLISEKEMYYIFALLGGLRQFCIHDKNR-----A	252
PIG--028...GL0185479	-----NKKMRLISEKEMYYIFALLGGLRQFCIHDKNR-----A	254
250twins...35838...GL0110300	-----KESEKAKADQDFVLRLLSVGRQLTFHSDKSNE-----	305
PIG--014...GL0226364	-KSEEEKRRIWQELWKNKFDHNKNIIVNSLIKFRQMCFHQYENS-----KEVS	259
250twins...36050...GL0158985	-----RAKAEIDACTAHNYDVLRLLSLMRQLCMHVSACTAF-----KLA	309
emb!OJMM01002900.1!	-----DREIKNSYNALVKVLYLRLQFCMHGNTYTKR-----NEESFLS	287
emb!OGPN01002610.1!	-----NNEDEYKKNLRHNFNVRVLSFLRQICITHAVKCTGGAKNNGDSIKVEAES	321
emb!OCHF01000044.1!	-----GDNSKQYQENLRHNFNLRVLSFLRQICMHAHVHVS-----DEGCTRTQNYTDS	317
PIG--046...GL0077813	-----DDYEGRNEILRYNFNVLVLSFLRQICAHAKVKVSS-----DNDREKGGVFDVS	320
pig...chimera	-----DDCKRERNEILRYNFNVLVLSFLRQICAHAKVKVSN-----EHDREKGGGLVDS	322
emb!OIEN01002196.1!	YNKVNKGKEEKDNNNENIEKLEKALEVIKIIRVDSFHGVDGIKGD-----QKF	260
O2.UC29--0...GL0096317	EQ-----QIEQEREREKENLETLEKALDVISVMRTDSFHGYDMCNSTS-----AIVKF	253
CasR...Anaerobic...dig	-----GKGEKIEVMYRSDEEIFTFQILSYVROSIMHNDI-----GN	279
emb!OGDF01008514.1!	-----GRIVECLEQRSDDIYNILPILSIVRQTCMHDNA-----SM	281
emb!OGZC01000639.1	-----RKIVKTEKEVYHILTLIGSLRQWITHTSTEG-----GIS	315
Ga0224415...10048792...chimera	-----NAKVRQENQCFHIFALLISLRNWATHSNELDK-----HTK	249
emb!OCVV012889144.1!	-----KKKRDKNEFYLKYDNECYIIFALLSGLRHWNHSHAKDD-----LVS	247
Ga0187910...10006931	-----KVLNKQDNERYHILALLSGLRNWVWHNNEVES-----KID	258
R...flav...XPD...	-----NYIINYGNECYDILALLSGLRHWVWHNNEES-----RIS	253
Ga0224415...10007274	-----RNNRQKDPICYHLLALLCGLRNWVWHNNEEKD-----LIK	236
Ga0187911...10069260	-----ADADKKEQVYAMLAACLGSLRQACSHYRYSVNG-----KNVDADA	245

FIG. 18T

Ga0187910_10015336	317KVMRIEKEIYYILGMLNEVRNVSTHTEEDN.....RDWA
ODAI_chimera	313	-----LKYREAEIYYIICMLGEARQMFAGND-----DRI
31009_GL0034153	252	-----LKDKKDIIDMELLLGSLRQFCFHFKGAKFEGV-----EQIS
DLF014_GL0011914_	315	-----KPKKVEVLKSEKEIFDILALLGSLRQFCFHDEAVFENEDEV-----IDKEYND
CasR_R_...albus	298	-----TKLRPEKEIYHIFALMASLRSQYFHGVKDT-----DYQG
Ga0187910_10040531	290	-----RRDEKVIYHILSLAASVRQFCFHNDYTSDDGK.....GFIK
tpg!DUXD01000002.1	235	-----RRYEEDIFAILLLSALRQFCFHSLSDES-----DHVN
CasR_E_...siraenum	306	-----AKNVLREDKELYVLTILGKLRHWCVHSEE-----GRA
FIG-018_GL0023397	287	-----SKRRSEKEIYSMLAMLGKLRHWCVHSES-----GEA
FIG-025_GL0099734	301	-----KRSKREIYDILALAIATIKWCISEE-----DK
Ga0129317_1008067_chimera	237	-----FEKRSDEEVYNICALMQIROCCEFHGKQ-----EKYQ
EYZ-362B_GL0088915	295	-----NQKKRLYILMSMIGQLRQFAFHKDS-----PFN
uncultured_Ru_sp	299	-----VSQAYKKRVYHMLAIVQIROCQVPHKS-----GAK
R_...flav_...FD-1	304	-----KNEEIKKRLYHLTAFAGQLRQWSFHSAG-----NYP
tpg!DBYI01000091.1	302	-----KSEEIKKRLYHLVAFAGQLRQWSFHSVE-----GLP
		* * *
037_...__emb!OIZA01000315.1	328	DIELYSENYS--E-----EYVFINCLINEEFVKNKFNVNKN--F
emb!OCTW011587266.1	286	RIAVFKNQFDS-----FSHTFTVTKIYSAKIACVNNEN--F
k87_11092736	274	SLAIFGANAMPEI-----PFFQGAIGGRQLWNDVIAPLWKKRIEVPKRS--F
BMZ--11B_GL0037915	288	SLWASDLGVEK-----YFKNEM--SVVDANYKARAKKIDAN--F
BMZ--11B_GL0037771	287	YEFWKSSDIYET-----KIFKAGW--SVIEADYKKKIQDVNKQ--F
BMZ--11B_GL0069617	265	--SILFCDNLKG-----Q----FLGNGNW--ITIEHYDEMITFINES--F
Ga0099364_10024192	327	--ATLFMSDVELPK-----ILK--KEFSKADW--QIVEDYYAKLVDRINEG--F

FIG. 18U

530373_GL0023589	SAWMFNLEKVL	332
emb ODAI011611274.1:	FAWLYRLDRQLS	346
EMG_10003641	NSSIYTLSSYDSTMNTAELLGKGDSSLVALATDARVEARAILDEIYKKGVDSINN-SF	334
CasR_P1E0_metageno	TAALYHLQALP	294
emb OIZX01009427.1:	AAGLYNLRRLG	322
160582958_gene49834	KFPFYNLEKNLF	374
gi 1198542314 gb NFIR01000008.1:	RAEYKLGKFEFRL	346
emb CGNF01009141.1	SVIMYNLDNEELYK	300
Ga0129306_1000735	SARIFRLDNDLG	333
MH0288_GL00822219	RSWLYNLEKVL	377
PIG-022_GL00226351	RNWLYSLDTMLN	286
PIG-028_GL0185479	RNWLYSLDTMLN	288
250twins_35838_GL0110300	-AYLFDLSKLR	349
PIG-014_GL0226364	STALFNIEQFL	293
250twins_36050_GL0158985	ESALFNIEDVL	343
emb OJMM01002900.1:	DTALYNAKEFFAK	324
emb OGPN01002610.1:	LDALFNITEYFAK	358
emb OHCPO1000044.1:	LEALFNISKAFGK	354
PIG-046_GL0077813	LDALFNISSEFFDA	357
pig_chimera	LDALFNISRFFDA	359
emb OIE01002196.1:	PRSKYNLAVN	294
O2_UC29-0_GL0096317	PSKKNLDQ	284
CasR_Anaerobic_dig	KSSILAEK	312
emb OGDF01008514.1:	RTVMFTLGQNSVR	322

FIG. 18V

emb!OGZC01000639.1	RLWLYQLEDALSREYQETMNCYNSTIYGLQKD--F	349
Ga0224415_10048792_chimera	KRWLYKLDFTNILNAEVKTLNLYDIADDELTKS--F	285
emb!OCVV012889144.1	YRWLYNLDISILNREYFISTLNLYLDDIADDELTKS--F	281
Ga0187910_10006931	RKWLYNLDKNLDKEYITITLDYMSYDIADDELTKS--F	292
R_...flav_...XPD...	RWLYNLDKNLDNEYFISTLNLYLDRIITNELTNS--F	287
Ga0224415_10007274	YTWLYNLDKYLDAEYITITLNYMYNDIGDELTDTS--F	270
Ga0187911_10069260	DTWLFSSAQLDQTDPLFSEMNLNRYSHKIKVYVQNQ--F	281
Ga0187910_10015336	KANLYNLSNRLKVGSKEVLNQLYKEKIDKIDANGF	352
ODAI_chimera	KTFLYNIVGYRNGTEINSFTAVLDGIYSKKVEELNGN--F	352
31009_GL0034153	NSWLYNLSQLHNEFKSTLDSFYNEKIEDINGN--F	285
DLF014_GL0011914...	NSWLYNLEKLEPFKDTLDYFYNRKINDINKG--F	348
CasR_R_...albus	TTWAYTLEDKLKGPSHEFRETIDKIFDEGFSKISKD--F	335
Ga0187910_10040531	ADWMYRLEEALPAEYKDITLDALYLEGVEGLDQS--F	324
tpg!DJXD01000002.1	SFWLYQLEDQLSDEFKETLSILWEEVTERIDSE--F	269
CasR_E_...siraenum	EFWLYKLDELKDDFKNVLDVVYVNRVPEEINNR--F	339
PIG-018_GL0023397	EFWLYKIDSLNSEFIDVINTLYNRSVNEINRG--F	320
PIG-025_GL0099734	KTWIYNAENVLDSEYLDILDVYVGLVEKINKN--F	335
Ga0129317_1008067_chimera	LKWLYNFHNEFKSNKRFDTLTKHFDEMDRINKN--F	272
EYZ-362E_GL0088915	RWLYYQLDEKLSKSHKDTLNYFFDQRYDEIHEN--F	329
uncultured_Ru_sp	RFDLYSFINNIDPEYRDTLDYLVEERLKSINKD--F	333
R_...flav_...FD-i	RWLYKIDSLDKREYDUTLCHYFDKRRNDINDD--F	337
tpg!DBYI01000091.1	RTWIYQLDNPKLAQYRDTLDYFFNERFDINKD--F	337

;

FIG. 18W

037___emb|OIZA01000315.1| I--SNE--KNNLYIILKAYGEDTENVEVVKYYSKELYKLSVLKTNKNLGVNVKLRSALEY 386
emb|OCTW011587266.1| L--NNE--GNNLYIILKATNWD-----IKKIIVPQLYRFSVHKSDDKMGFNMRKLRFAVES 388
k87___11092736 L--SNS--AKNLWVLYQVFKDDTD--EKKARARQYHFSVLKEGKNLGFNLTKREYFLDK 330
BMZ--11B__GL0037915 M--ATS--RANLVLLFRILNAKTK----EKEKIAVDYKFAIRKEGKNLGVNLKAREILLDK 344
BMZ--11B__GL0037771 F--SKN--KVNLAIFDLDDVRDV--KQKRIISDEFYRFTIRKDKGNLGMNLVKIREIIDD 343
BMZ--11B__GL0069617 F--NNS--RNMLKIYLVLNAATL---EDKKIITEEYRFSILKEGKNLGVNMRKLRMVMVDM 321
Ga0099364__10024192 C--KNA--ATNVHFLTELLPE-----ESKQUTEDYFRFALLKEGKNLGVNMRKLRVVFAL 380
530373__GL0023589 I--ENS--NKNLNAIFRLEFPEKS-----QADLIVEYYDFSVKAKYKNLGFSVKLLREMLLKT 385
emb|ODAI011611274.1| L--TNS--TVNLEVFRALKTGTD--PERKTVTQYEQYQYTVRKEDGNLGFSLKILREILLSA 402
EMG__10003641 L--SNS--INDLENLEFKIYKCDSS---EKKTELIKQYDFCIRKFPQNMGFSITTTIREGMFTR 390
CasR_P1E0__metagenc L--DTN--KVNLLLEFYEGAETK--QARVALTQDFYRFVVLKEQKNMGFSLRRLRELLKL 350
emb|OIZX01000427.1| Q--DHN--TTNLTCLFDILNISDS---ARQQLAEFYRYVYVFKQKNLGFSVKLRREMLLL 378
160582958_gene49834 G--TAGKTQVSVLLKVFEKEQASQEQALLKEYYDFKQKTYKNMGFSIKKLRALMEI 433
gi|1198542314|gb|NFIR01000008.1| L--KNS--KSDILMLFRIYNAESK---EAKRKLAEYEFIMLSYKNTGFSIKHLRETWIDK 402
emb|OGNF01009141.1 I--KNS--AKDFVILCEVLGIKSR--DEKTKFVKDYDFVVRKNYKHLGFSVKLELLELFAN 356
Ga0129306__1000735 D--KTSASSNFRLLFNAYHADN----EKKKELAQEFYRFSVLKVSKNIGFSIRTLREKLIED 389
MH0288__GL0082219 S--KTSSKSNFRLLFNALHADTD---AQKQJAEDEFYKFSVSKTYKNIGFSIKLRESMLDT 434
FIG--022__GL0026351 V--DNSAKKDFSIIFNALNITDI---ELKKNIAKEYYEFVCRKNFKNIGFSIKKLRLEYLLEK 343
FIG--028__GL0185479 V--ENSAKKDFSIIFNALNITDI---ELKKNIAKEYYEFVCRKNFKNIGFSIKKLRLEYLLEK 345
250twins_35838__GL0110300 V--KHA--KNNLYVLNQLYPS-----LKANDLIGEYMFVYKKNRNI GIRLITVYRELIIEH 402
FIG--014__GL0226364 N--KNS--SNNLYILEQIYCN-----D----KNLQRYDFVYVYKSKQC NLGLDTKLIRELLIRK 344
250twins_36050__GL0158985 V--QHS--GNNLYVLQQLYPN-----ETIERIAEKYRYRLTVKEDLNMGVNIKLRRELIIVGQ 396
emb|OJMM01002900.1| M--AHA--KNNMYIICEVYKYN-----EAFDSLMEKEYYDFVYVYKSKGNNLGFNTROLREILDK 377
emb|OGPN01002610.1| V--TNG--KNNLYILSKVYYPD-----MQNELVKKYQYVVCQKGGNNVGINTRKLRKESIISQ 411

FIG. 18X

emb!OHCPO1000044.1!	V--KNG--KNNLYILSKVYPN-----EKREVLRLREYVNFVVCKEGSGNIGISTRKIKETMIAQ	407
PIG-046...GL0077813	V--KND--KNNFYILSKIYPE-----VARENLLKEYYYFVVSKEGNNIGISIKKLKEAIIAQ	410
pig...chimera	V--KNG--KNNFYILSKIYPE-----VAREDLLREYVNFVVSKEGNNIGISIKKLKEAIIVQ	412
emb!O1EN01002196.1!	Y--RNS--CVNIDFLKEIMYGSNYTDRGSDSLECSYFNFAILKONKKNMGFSITSIKRECLLDL	352
O2_UC29-0...GL0096317	A--QNS--KVINFLSYVLYGNGCTEEQKELSRFYFNFAILKQYKKNLGFSSITSIKREQLLSL	342
CasR_Anaerobic_dig	IDNNS--QTNEWVLSIFGL--QDHTSGADKICRNFYDFVFKADSKNLGFSILKRELMIDL	370
emb!OGDF01008514.1!	L--RNQ--KNNIELLSKIYGS--SADSPERDLVQNFYDFVLSQDKNLGFSIKKREKLLDS	379
emb!OGZC01000639.1	E--KTN--APNLNFAELILGK-N-----ASELAEPYFRFIITKEYKNLGFSSIKTLREMLDQ	401
Ga0224415_10048792_chimera	S--KNG--AVNVNVLAKKYNI-K-----DDLEGFSEQYFRFSIMKEQKNLGFNISKURENMLDF	339
emb!OCVV012889144.1!	S--KNS--SANVNYIAETLNI-D-----PSEFAQOYFRFSIMKEQKNMGFNVSKLREIMLDR	333
Ga0187910_10006931	S--KNS--AANVNYIAEILNI-D-----SKTFAEQYFRFSIMKEQKNLGFLLTKLRECMLDR	344
R...flav_XFP...	S--KNS--AANVNYIAETLGI-N-----FAEFAEQYFRFSIMKEQKNLGFNIITKUREVMLDR	339
Ga0224415_10007274	S--KNS--AANINYIAETLGI-D-----PKTFAEQYFRFSIMKEQKNLGFNLTKUREVMLDR	322
Ga0187911_10069260	FENNR--KANFPIKKNMYPE-----TTLKVLNNEYDYSFKGKKNFGFSIKSREALLSP	335
Ga0187910_10015336	VNKGC--KRDFSILPKSLNL-I-IDKDKGELVVGFDYDFSPKNYKNIGFSIKTLREYMLKI	409
ODAI...chimera	I--KNS--KKDLTILFSAFGC-Q--DEASKKKLIEKYYDFIVKTKNKNMGFSVTKLREKILIEG	408
31009_GL0034153	I--EKN--KVNIIQLCNDIKL-----DNKLVQDYVNFVLSKNHKNMGFSIKKIREYIIDI	337
D1F014_GL0011914...	I--KTN--QINLHIICDELGL-N-----WDKEQVVGDIYDFLLAKKYKNIGFSIKKIREHMFDI	402
CasR_R...albus	G--KMN--KVNLIQILEQMIQE--LYGSIERQNLTCDIYDFIQLKHKHYLGFSSIKRLRETMLET	392
Ga0187910_10040531	L--KNN--TVNIQILCSIFNH-----DDPNKIAEYYGFLMTKEYKNMGFSIKKURECMLEL	377
tpg!DJXD01000002.1!	L--KTN--TVNLHLCHVFPK-----ESKETIVRAYEFLIKKSPKNMGFSIKKUREIMLEQ	322
CasR_E...siraenum	I--ENN--KVNIIQLIGSVYKN-----IDIAELVRSYYEFLITKKYKNMGFSIKKURESMLEG	392
PIG-018_GL0023397	V--DTN--KVNIIQLIQAVCKN-----NDLISLIKAYEFLITKKYKNMGFSIKKUREMMLAE	373
PIG-025...GL00999734	I--KDN--KVNLIILSDILDM-K--SKDDFEELIRQYRFRFIITKEQKLLGFSITIKREAMELD	391

FIG. 18Y

Ga0129317_1009067_chimera	I-KNN-TPDLILLGLYPD-----MAKKELVRLFYDFIVVKEYKMGFSVKKLREKMLLES	325
EYZ-362B_GL0988915	N-KIN-KVNLELLAEAFPS-----EDYATLANQYDFVFKAHKNLGFSSIKKLRKLIEN	382
uncultured_Ru_sp	I-EDN-KVNISLLIDMMKG-----YEADDIIRLYDFIVLKSQKNLGFSSIKKLRKMLDE	386
R_flav._FD-1	V-TKN-ATNLYLLKEVFE-----ANPKDIADLYDFIVLKSQKNLGFSSIKKLRKMLDEC	390
tpg!DEYI0109091.1:	I-ETN-NINLHILKEVFA-----EDFQKLAALYDFIVKKTFFKNIGFSIKNLRQMLEC	390
	: . : : * : *	
037__emb!OIZA01000315.1:	GYCELP-----YDKEKEVAKLSSVHKLYKTYDFVITHILNSN-----	424
emb!OCTW011587266.1:	KNIDL-----SRLNDRKFLTNRRKLYKVIDFIIYYHLNKVL-----K	375
k87_11092736	FFPIFH-----SSAPDVKRKVDTFRSKYAYAILDFIIYEASVSV-----ANSQQMGKVAPW	380
BMZ-11B_GL0037915	HYPD-----AKNKQFDSYRCKLYITDFIIYRMAED-----	376
BMZ-11B_GL0037771	YASG-----LRDKKHDPHQKINVIADFLIFRALSON-----	375
BMZ-11B_GL0069617	FYSE-----IKDKQYDSYRCKLYITLDFLLFREIKDS-----	353
Ga0099364_10024192	FVPELT-----APETKKRYDSYRCKLYIGLDFLLFKHHT-----	416
530373_GL0023589	QQELS-----PLNDKAYDSIRSCLYTMDFVLYKLYVD-----N	419
emb!ODAI011611274.1:	YKH-----EVRDKEYDSIRHKLYQLFSFALYHYKFTG-----VG	436
EMG_10003641	CSEANTLLCDEGSTVKLNVDHDTMKSFKYKKNLDFMIYKYKYE-----N	434
CasR_P1E0_metageno	PDAA-----YLTQEYDSVRQKLYNMLDFLLCRLYAGE-----R	384
emb!OIZX01000427.1:	PDAA-----VIADKRYDTCRSKLYNLMDFLILRVYPTG-----R	412
160582958_gene49834	PDAA-----KFKDDLYSSLRHKLYGLDFILVKHFLDT-----SDSE	470
gi 1198542314 gb NFIR01000008.1	MDEDK-----KEKLDKKNPIRRKLYRIMDFMIYQYQEP-----EH	440
emb!OGNF01009141.1	HDSNKYIK-----EFDKISNKKFDSVRSRLNRLADYIIYDYNK-----N	396
Ga0129306_1000735	HAA-----QYRDKIYDSMRKKLFTDFDLWRFEYEE-----R	421
MH0288_GL00982219	FAP-----EYKEKSYDSVRCKLYQLVDFLIWNLYDK-----N	466

FIG. 18Z

FIG-022_GL0026351	PEFC-----NLKEDKYNVRSKLYRLIDFIIYLYNYKDAKF-----ENDN	382
FIG-028_GL0185479	PEFR-----NLKDDKYNVRSKLYRLIDFIIYLYNYKDAKF-----KNDN	384
250twins_35838_GL0110300	NYT-----NLKDSKYDYYRNKLYTVLNFILFRLEQEN-----	434
FIG-014_GL0226364	LSD-----EGLDINDSRDYTYRNKLYTVFGFILYVELTKN-----	379
250twins_36050_GL0158985	YFPE--VLDKEXDUSKNGDSVVTYRSKIYTYVMNYILLIYLEDH-----	437
emb!OJMM01002900.1!	YVG-----NLRGKKYNTFRNKLTYVLGFILVKEIKRN-----	409
emb!OGPN01002610.1!	HPWI---TTPQD-----NNKANDYESCRHKLYTIMCFILVVELDAH-----	449
emb!OHCFO1000044.1!	NMP-----SLKEENTYRNKLYTVMNFILVRELKNC-----	437
FIG-046_GL0077813	DMS-----YIKSKDYDYYRNKLYTVLFCFILVKELNER-----	442
pig_chimera	DMS-----YIKSEDYDYYRNKLYTVLFCFILVKELNER-----	444
emb!OIEN01002196.1!	YEL-----NFESMQNLRFRANSFCDFLIYDYCSNE-----	383
O2_UC29--0_GL0096317	YDI-----DFECKRNLRSRANSFCDYVLYRYAVQF-----	373
CasR_Anaerobic_dig	PNA-----NMLRDHQFDVTRSKFYLLDFIIYQHYLEE-----	403
emb!OGDF01008514.1!	PAL-----SVVRSKDYDYYRNKLYSLDFMIYRPFSEN-----	412
emb!OGZC01000639.1	PDLQ---E-----IRENHNVYDSIRSKLYKMDVFLVYAYSNERK-----	438
Ga0224415_10048792_chimera	KDMS---V-----IRDDHNRYDKDRSKLYTMMDFVIYRYEEDNNN-----	376
emb!OCVV012889144.1!	KELS---D-----IRDNRHVFDVIRSKLYTMMDFVIYRYEIEEAAKTEAENRNLPENEKK	385
Ga0187910_10006931	EELS---D-----IRDNAKVFDVIRSKLYTMMDFVIYRYEIEEAKKIEENKTLSDDKKK	396
R_...flav_XFD_...	KDMS---E-----IRKNHKVFDVIRSKLYTMMDFVIYRYEIEEAKVAAANKSLFDNEKS	391
Ga0224415_10007274	KDMS---E-----IRENHNFVDSIRAKVYTYMMDFVIYRYEIEEAAKVAANKSLFDNEKS	374
Ga0187911_10069260	QYES---LIGV---QIKDNKEYDYVRSKLYQLFDALTRYFNQH-----	373
Ga0187910_10015336	SNS-----I-----LCADTISNNAIRPKAYKLYDFIINHYYMKN-----	443
ODAI_chimera	YRFR---G-----NPVTGTDYDSVRHKLYRTFFDIICSYYDDH-----	443
31009_GL0034153	DYA-----SYIKEOGYDSVRSKLYTLIDFIIYDLYVNKE-----	371

FIG. 18AA

```

DLF014_GL0011914_    YEA-----FDIKRFRFSVRSIMYKIIDFVIYYSFIHYK----- 436
CasR_R._.albass      TPA-----ECYKAECYNSERQKLYKLIIDFLIYDLYNKK----- 426
Ga0187910_10040531  PEL-----SGYKEDQYNSVRSKLYKLFDFIIAHYFRKH----- 410
tpg!DJXD01000002.1| SDL-----KSFKEDKYNSVRAKLYKLFDFIITTYIDHH----- 355
CasR_E._.siraenum    KGY-----ADKEYDSVKNLYQMTDFILYTGYNED----- 423
FIG-018_GL0023397  TAYT---SNPD-----DRLYDKEINSVRNNLYQMTDFIILEYEDN----- 411
FIG-025_GL0099734  TV-----FKIDKKYDSIRSKLYKLIIDFMMFYATYTDSE----- 425
Ga0129317_1008067__chimera  EEA-----SDFRDKDYDSVRRKLYKLMDFCIYYLYYS-D----- 358
EYZ-362B_GL0088915  -----ALRDEKIDSFRSKLYRLIDFCLFRYLK-D----- 411
uncultured_Ru_sp    YG-----FRFKDKQYDSVRSKMYKLMDFLFCNYYRND----- 419
R._.flav._.FD-1    DGA-----DRIKEQDMDSVRSKLYKLIIDFCIFKYH-E-F----- 423
tpg!DBYI01000091.1| DEA-----EKIRSKDMNSVRSKLYKLFDFCFIFYQYFI-D----- 423

                                :      :
                                :      :

037_--__emb|OIZA01000315.1|  DKLILEIVETLRISK-----ND-DEKENVYKKA-----E-KLFKADDVINPIKAISKLFAE 474
emb|OCTW011587266.1|    DSFVDDFVAALRASQ-----SE-EEKKLYAQYS-----E-PLFADEGLKSAIKKAVDMISD 425
k87_11092736            KGALDNALVKLRAP-----DE-EAKEKIYNVLA-----A-SIRNDSLFL-----RLKSACDKFG 428
BMZ-11B_GL0037915      GAFLEESVQKLRSTP-----SE-EEKKVVYDVLA-----E-IWNATKDD---VKKVFDIRQQ 424
BMZ-11B_GL0037771      QGIIDKTVSSLRITK-----DE-EEKDHVYQNAE-----E-LVWGMVSNC---LTFYFNDPKN 423
BMZ-11B_GL0069617      -EELETWVADLRSE-----SD-EEKKELYHDYA-----E-LFWDGVKDL---IVPFFFAFDG 400
Ga0099364_10024192    -KQLEEWVAVLRFTS-----NE-DAAENLYDEFA-----R-TAWNTVGDSE---AKQLIENMQS 463
530373_GL0023589      EEQIDQFVRKLRAST-----SN-VEKEALYQCES-----E-PTWNEIRGQ---IIDNVIPFID 467
emb!ODAI011611274.1|  AERREAFVAKLRVM-----TA-EAKQRAYADEA-----A-EIWNDEGSG---IRAAFLEILE 484
EMG_100003641        PEKGEKLIEDLRKSKIKGKKED-EDKKQRYAES-----A-CILFKAKRDI---IKKDLTEAAN 467

```

FIG. 18BB

CasR_P1E0_metageno	432	ADRCEELVSAALCAL.....SD--EEKDFVYQAEA.....A--ALWQALGDT----LRRKLLPLLK
emb O1ZX01009427.1	460	ADKCKLFEALRAAL.....TD--EERAVVYHKEA.....L--SLWEMRTL---ILDGLLPQMI
160582958_gene49834	518	NLQNDIFRQLRACR.....CE--EEKDQVYRSIA---V--KVWEKVKKK--ELNFKQVVV
gil 1198542314 gb NFIR01000008.1	488	QEEAEELVRKLNAAE.....IE--AKKELAYRKEA.....E--KLKKELEKI----IENSVLPSCD
emb OGNF01009141.1	445	NAKVSDLVKYLRAAA.....DD--EQKKKIYLNES---INLVKSGILER--IKKILPKLNG
Ga0129306_1000735	469	EDEAEELRACLRAAR.....SD--EEREQIYAEAA.....A--SCWPSVKPF...VESVAATLCD
MH0288_GL0082219	516	EKWADALVAQLRAEVK....RDR--DEKSPVYDREA....A--RLWTMIREA----VAQTAKTLRA
FIG-022_GL0026351	429	DEKIERFVSKLRKCP.....DE--ASKEFVYQTES---D--IIKKSIYGS---INSIKRGMN-
FIG-028_GL0185479	431	DEKIERFVSKLRKCP.....DE--ASKEFVYQTEA----D--IIKKSIYGS---INSIKRDMN-
250twins_35838_GL0110300	481	SLAIKNFREKLRSTE.....K---AQPALYQAFN---N--KIYPMVQAK----FAKALDFEE
FIG-014_GL0226364	431	ETFCQEIYAKLRBFG.....GDEIERKNIYNEFLRKE--AVYNOIAEK---FKQSVCVWFKK
250twins_26950_GL0158985	486	DSSRESMVEALRQNR.....EGDEGHEEIIYRQFA.....K--KVWNGVSGL---FGVCLNLFKI
emb OJMM01002900.1	458	PKIQDSFIAKLRANQ.....NGDEGKLNLYNEFA---P--KIWSVSVSSK---LNSAITCFDE
emb OGPN01002610.1	498	ESIRDNMVAELRANM.....DGDDGRDAIYEKYA---K--DIYHIVKDK---LLAMQKVFDE
emb OHCPO1000044.1	486	ATIREQMIKELRANM.....DEEEGRDRIYSKYA.....K--EIIYLVKDK---LKLMLNVFKE
FIG-046_GL0077813	491	TTIREQMVAELRANM.....NGDIGREDIYSKYA---K--IYYAQVKPR---FDGMKNVFEE
piq_chimera	493	TTIREQMVAELRANM.....NGDIGREDIYSKYA---K--IYYAQVKPR---FDTKMSAFEE
emb O1EN01002196.1	430	--SERANLVDCLRGAA.....SE--EEKNNIYFQTA.....E--RVKKEFRNA---FNRI-SREFDAS
O2_UC29-0_GL0096317	421	PERIVDLVKLRCSA.....NE--KEKTNVYIEEA---N--RIKEDLRIA---FNRINNFDAK
CasR_Anaerobic_dig	451	KSRIIDNMVEKLRMTL.....KE--EKEEVLVYAAEA....K--IVWNAIGAK----VINKLVPMMN
emb OGDF01008514.1	461	HVAVDDFVEELRSGUL.....TE--DEKESAYSRSWA.....FTLINDGFAQE---ILVKLLPQTD
emb OGZC01000639.1	485	--SKADALASNLRSAI.....TE--DAKKRIYQNEA---D--QLWTSYQEL---FKRIKGFKGA
Ga0224415_10048792_chimera	422	--DSIDFINKLRSSI.....DE--KSKEKLYNEEA....N--RLWNKLYKEY----MLYIKKEFNGK
emb OCVV012889144.1	433	YSEKDFEVINLRGSP.....DE--NQNEKLYIEEA.....K--RLWKEKLIKDI---MLNKEEFRGE

FIG. 18CC

Ga0187910_10006931	LSEKDFIISLRGSF.....SE--EQKDKLYSDEA.....E--RLWAKLGLK.....MLEFKKFRGQ	444
R_...flav_...XPP...	LSEKDFIVINLRGSF.....ND--DQKDALYYDEA.....N--RLWKKLENI--MHNIKEFRGN	439
Ga0224415_10007274	LSEKDFIVISLRGSF.....NE--DQKDRLYYDEA.....Q--RLWSKVGLK--MLKIKKFRGK	422
Ga0187911_10069260	PDMVDAFVVELSLA.....KDEDAKNAVYEKYA.....K--AVWNDVKOP----IAVMLSYMNG	422
Ga0187910_10015336	PKINDFVEKLRQN.....KN--DEAIKLYYDEA---V--CLLSELGRE--IHTMFSVHN	491
ODAI_chimera	AKKDTLVSSLRASL.....ND--TKKENLYAGEA.....K--TIAESIKDR--MDTVCRMYNK	491
31009_GL0034153	KDRIDDIIVSKLRGAI.....YD--YKDDIYKAEA.....D--WVWKKLNQO----IKALTKLLSN	419
DLF014_GL0011914...	QDEVDEIVNYLRSTL.....SE--EQKKEKYYIKLA---G--NINAEYKTO---FNKLDILLA	484
CasR_R_...albus	PARIEIVDKLRSEV.....ND--EERESIYSVEA.....K--YVESLSKV---LRSLSKNSVS	474
Ga0187910_10040531	PEKGEEMVDCLELGM.....TE--DEKDSHYEGTA---K--KLVRRELAYD---MQEAAEQANG	458
tpg!DJXD01000002.1	AEEKEALVSSLRSSL.....TE--ENKEEIIYIKTA---R--TLASALGAD---FKXAAADVNA	403
CasR_E_...siraenum	SDRADDLVNTRSSL.....KE--DDKTFVYCKEA.....D--YLWKKYRES---IREVADALDG	471
FIG-018_GL0023397	KSIIDITLVSKLRSSV.....TD--EAKAVYREEA---A--NLGRLLDA---VALIAKSLNG	459
FIG-025_GL0099734	KEHADSIVEKLRATL.....SG--ESKEEIIYKAEA---N--FIWNLNLYEDI---IMNKIEPALD	473
Ga0129317_1008067_chimera	SERNENLVSRLRESL.....ID--ENKDIYSKEA.....K--IVWNELRKK---FSTILDNVKG	466
EYZ--362E_GL0088915	PDRAEKNVRFRLRSCI.....NE--QEKEQFYTOEA---N--ALKSENPGI---VNDFCMQLKD	459
uncultured_Ru_sp	IAAGESLVRKLRFSM.....TD--DEKEGIYADEA.....A--KLWGKFRND---FENIADHMNG	467
R_...flav_...FD-i	PELSEKNVDILRAAV.....SD--TKKDNLYSDEA.....A--RLWSIFKEK---FLGFCDKIVV	471
tpg!DBYI01000091.1	EERSRENVNLYLRSTL.....ND--EQKDAFYEEEG---K--PLMSENRRK---FIYFCDNINK	471
037_...emb OIZA01000315.1	KGNKLFKEK.....IIKKEYIEDVSDKNIIDFTFVIFVFFMTCF	513
emb OCTW011587266.1	TKSNIFKMK.....TPLDKALIEIKVNSDASDFCKLIYVFTRE	464
k87_11092736	AEQNRPFVFPNELR.....NNRDIRNVRSEWLEATQVDVAAAFAVQLIAFLCNF	475

* * ; *

FIG. 18DD

BMZ-11B_GL0037915	DIKNAAAQSLDDWLDSTVTEKEKEFFVVKLLAFLCNF	462
BMZ-11B_GL0037771	KYILKYKDA	-----KTPGDFEDWITSKISEDDEGPFVVKVLSFLCNF	464
BMZ-11B_GL0069617	DFTVFTTEK	-----VPADTSKEFFLSSANTEFLVKLLAFLCNF	438
Ga0099364_10024192	YFTKKEKEIT	-----IAQPVLSSTSSIAHTSKKITQFSSFAKLLAFLCNF	508
530373_GL0023589	SIQKQENEDKLTKR	-----IDQSIIDKAVEHFVNSRDLSYFAKVVVVVAVSF	513
emb ODAI011611274.1	AVDFCSAVKGIKAPSS	-----VAGDKRFAEWLEEVIRPEGVSCFTKLMYLLTRF	534
EMG_10003641	KDLFADLVKSNKNEKQK	-----FKNEYELLEPFMIPVKVDYFSELIYLVTRF	535
CasR_P1E0_metagenc	GKKLQDKDK	-----KRSDELGLSRDVLGVLEF-----PAQQGSRANADYFCRLMHLSTWF	483
emb OIZX01000427.1	PENLSRLSGQKRCGELSLDDAMLKECLYE	----PGPVPEDAAPEANAEYFCRMTYLATLF	517
160582958_gene49834	IPSLSKDE	-----LQKQEMTKNTELLSSLETISTQASLFSSEMIFMMTYL	562
gi 1198542314 gb NFIR01000008.1	RILSEMDERRNKVNQESSDTRKEEPL	-----DSEIAEGITFIKETAHSFSEMILLTVF	543
emb OGNF01009141.1	KLIIGNMQPDSI	-----TASMLHNTGKDWHPISENAHYFFIKWIYTLTLF	489
Ga0129306_1000735	VVKGR	-----TKLNKLSADESTLVKRNAIDGVRI SPRASYFTKLIYLMTLF	516
MH0288_GL0082219	VTLNNTPGSIKELALSKYDRDLVENAIDSAVDERGRLKITDEASYFSKLLYLLTLF		576
FIG-022_GL0026351	-----SVKSKYKLELDKQTI	-----NQLKRAVDVFSINTKGDYFFIKMLYLLISLL	474
FIG-028_GL0185479	-----LVKSKYKLELDIQT	-----VQNKRAVDVFSINTKGDYFFTKMLYLLISLL	476
250twins_35838_GL0110300	QYKTKFKSEFKGGI	-----SIENMQQQNILLQ	525
FIG-014_GL0226364	EKKKKKEFPSS	-----KTKIDLRVQKLEDQNASVFAKMWYLLISRF	471
250twins_36050_GL0158985	EKRNRKRSKV	-----ALPDVSGAAYMLSSENIDYFVKNLEFFVCKF	526
emb OJMM01002900.1	ESLSKFKGYK	-----DIDESLISRYGITVANIDTLVKLIYFLCKF	498
emb OGPN01002610.1	ELVPPKVEGKNBPQ	-----Q-----FTHGKLGKKEIEFCLSDKNTSDIAKVVYFLCNF	547
emb OHCP01000044.1	EAEGIIPGKEDPV	-----K-----FSHGKLDKKEIEFCLTTKNTEDITTKVIYFLCKF	535
FIG-046_GL0077813	EAKGVIVPDKKEEAV	-----K-----FSNGALDKSEIERFCITSANTDSVAKIYFLCKF	540
pig_chimera	EAKDVIVPDKKKPV	-----K-----FSHGKLDKNEIERFCITSANTDSVAKIYFLCKF	542

FIG. 18EE

emb!OIEN01002196.1!	YLNRSREKNLSGGS---SLPK-----YSFIEGFTYKRSKINDENDEKNADLFCNMELYLAQF	483
O2_UC29--O_GL0096317	SIKLYSTKNNVDQ---FMLP-----FEQVSK-----DIKGGDAAREKADAFCDMTYYLITQF	479
CasR_Anaerobic...dig	GDALKEIKRKN-----RDRKLPQSVIATVQVNSDANVFSGLIYFLTILF	494
emb!OGDF01008514.1!	PAVIGKIKG-----KLLNDSIACIKLKKDASFFTKINVLICMF	500
emb!OGZC01000639.1	QVKEYSSKN-M-----P-----IPIQKQIQNLIKPAEQVTFYFKLMYLLTMF	526
Ga0224415_10048792_chimera	LASRTPDRD-G-----NISEFVESLPKRIHRLPRGQKISNFSKMLYLLTMF	467
emb!OCVV012889144.1!	KVKEYK-----KEVPRIERILPQGKDISAFSKMLMYLMSMF	468
Ga0187910_10006931	MTRDYK-----KSDTFTLNRILPESEDEVSTFSKMLYALTMF	480
R_flav_XFD_	KTREYK-----KKDAPRLPRILPAGRDVSAFSKMLYALTMF	475
Ga0224415_10007274	DTRKYK-----NMGTPRIRRLIPEGRDIDSTFSKMLYALTMF	458
Ga0187911_10069260	SAKNIKAFEL-----KPDQKELINGINNSNALDVPHFCKLVYFLTRF	464
Ga0187910_10015336	IENISYEITDK-----KQKEYYKMQINSLSADKVSDFSKVIYLVILF	534
ODAI_chimera	CECIKHITVDR-----FLAGACAGKLIDKCSISTECDYFCFKIYLMTLF	535
31009_GL0034153	RRIFSQDND-----ITNWNQVEFEKIGEQLKLEDNADYFCMLYFVTLF	462
DLF014_GL0011914_	NIKEPSDAKEDDV-----YEEFKNFKFNEVQKELGENAEYFCMLYLLILF	532
CasR_R...albus	GETIKDLQKRY-----DDEETANRIWDISQHSISGNVNCFCFKLIYIMTLM	518
Ga0187910_10040531	SNITQMQKNEQQG-----KT--KGMFAIRDEIRVSRKPVSYFSKVIYVMTLL	503
tpg!DJXD01000002.1!	KNIRDYQKKAN-----DYRISFEDIKIGNTGIGYFSELLYMLTLL	443
CasR_E...siraenum	DNIKKLSKS-----NIEIQEDKLRKCFISYADSVSEFTKLIYLLTRF	513
PIG--018_GL0023397	VNIKLLQON-----ILEIFDSKLENGCFIKSAEKVSFTKLNLYLLTRF	501
PIG--025_GL0099734	EGVISGLRK-----NKSVDNMSIEGIVAEETANVSYFSKIIYLLAQF	515
Ga0129317_1008067_chimera	SNIKKLEN-----VKEKFISEDEFDDIKLIDIDISYFSKLMYVMCYF	447
EYZ--362E_GL0088915	RYSSGNSSTENAE-----KKEDFPVFPHPCHKQKQVSNVSLFAKLMYAFCLF	505
uncultured_Ru_sp	DVIKELGK-----ADMDFDEKILIDSEKKNASDLLYFSKMIYMLTYF	508

FIG. 18FF


```

R_...flav_...FD-1          509 WVTGEHEK-----DITSVIDKDAYRNRSNVSYFSKLMYAMCFF
tpg|DBYI01000091.1:      509 WVKNDYSY-----EVAKCIDLNEFRVNSVSYFSKLLYAMSFF
                                *
037_..._emb|OIZA01000315.1: 548 LDGKEINDLLTNIISKLQVIEDHNNVIFH-----NKD
emb|OCTW011587266.1:      499 LDGKEINILLNSLKKFKQDIHSFNTVYKLLSE-----NNL
k87_...11092736          508 LECKEINELVTALIKKFKEGIQALIDLRLNLEGV-----
BMZ--11B__GL0037915      495 LDGKEINELLSTFVNRLENVQALIDLAKNLGET-----
BMZ--11B__GL0037771      497 LEGKEINELLTAYIHKFECIQDFLNVISSLGEN-----
BMZ--11B__GL0069617      474 WDGKEVNEVLSAYIHKFESIQAFIDVLESKEIN-----EKV
Ga0099364_10024192      541 WEGKEINELLSAYIHKFFENIQEFINLEKLECK-----
530373__GL0023589      545 LDSKEINMFFDALINELENIASFNDVLEQLDM-----
emb|ODAI011611274.1:      556 LDGKEINELLTGLINKLENIQSFLDVMOQEHRA-----
EMG__100003641          567 LSGKEINDLLTQLINKFENIAAFIRMVQNDQG-----
CasR_P1E0_metageno      515 MDGKEINTLLTTLISKLENIDSRSVLESML-----
emb|OIZX01000427.1:      549 MDGKEINTLLTTLISKFENIAAFIQIMEQLNI-----
160582958_gene49834      595 LDGKEINLLCTSLIEKFENIASFNEVLKSPQIG-----
gi|1198542314|gb|NFIR01000008.1: 575 LDGKEINILLTQLIHCFCFNISFMDTMEENL-----
emb|OGNF01009141.1      521 MDGKEINDLVTTLINKFDNIASFIEVLKQSV-----
Ga0129306_1000735      548 LDGKEINDLLTTLIHAFENIDSFVLGGERL-----
MH0288__GL0082219      608 LDGKEINDLITTLVNSFDEIDSFCCVLQKAKL-----
PIG--022__GL0026351      506 IDGKEINDLLTTLIHQFENIASFIDVMKKENI-----
PIG--028__GL0185479      508 IDGKEINDLLTTLIHQFENIASFIDVMKKENI-----
250twins_35838__GL0110300 557 LDGKEINELLCALINKFDNIADLLDISKQIGT-----

```

FIG. 18GG

FIG-014_GL0226364	LDGKEINELCTALLNKFKNINELVEIYAKRCGE-----	503
250twins...36950_GL0158985	LDGKEINELLCALINKFDNIADILDAAAQCGS-----	558
emb!OGMM01002900.1!	LDGKEINELCCAMNKFDNINDLIKIAAQCCE-----	530
emb!OGPN01002610.1!	LDGKEINELCCAMNKFDGIGDLIDIAKQCCE-----	579
emb!OHCFO1000044.1!	LDGKEINELCCAMNKLDGISDLEIYAKQCCE-----	567
FIG-046_GL0077813	LDGKEINEFCCAMNKLDGINDLMEIYAEQCGA-----	572
pig_chimera	LDGKEINELCCAMNKLDGINDLIEIYAEQCGA-----	574
emb!OIEN01002196.1!	LDGKEINIFLTSIHNIFQNIIDSFLVYKKEKGM-----	515
O2_UC29--0_GL0096317	LDGKEINVFLTTLHNLEFENIASFLDVYHKQNI-----	502
CasR_Anaerobic_dig	LDGKEINEMVSNLITKFNIDSLLVHVDREIYKSDERKDLLEIEKIALFFKGVVYRPNKATD	554
emb!OGDF01008514.1!	QDGKEINELVSSLVYKFNANIQSFVDVYKRSQGI-----	532
emb!OGZC01000639.1	LDGKEINDLLTTLINKFDNISSLLKIMEQLEL-----	558
Ga0224415_10048792_chimera	LDGKEINDLLTTLINKFENIQGFLDINPEINV-----	499
emb!OCVV012889144.1!	LDGKEINDLFTTLINKFDNIQCFLKIMPLIGV-----	500
Ga0187910_10006931	LDGKEINELLTTLINKFDNIQSMLKIMPLIGV-----	512
R...flav_XPD...	LDGKEINDLLTTLINKFDNIQSFLKVMPLIGV-----	507
Ga0224415_10007274	LDGKEINDLLTTLINKFDNIQSFLKVMPLIGV-----	490
Ga0187910_10069260	LDGKEINDLLTTLINKFDNIHSPVQVLTALGL-----	496
Ga0187910_10015336	LDGKEINDLLTTLINKFDNIASLLSVLEKQSG-----	566
ODAI_chimera	LDGKEINDLLTTLINKFDNIDSPSVKAVNNQ-----	567
31009_GL0034153	LDGKEINDLLTTLINKFDNIRSFIEVMEIEMSL-----	494
DLF014_GL0011914...	LDGKEINDLLTTLINKFDNIRSFVELMKEENI-----	564
CasR_R...aibus	LDGKEINDLLTTLINKFDNIASFIDVMDLGL-----	550
Ga0187910_10040531	LDGKEINDLLTTLINKFENIVSFDVLRQLNV-----	535

FIG. 18HH

tpg!DJXD01000002.1	LDGKEINLLTTLINKFDNIISFIDILKKNL-----	475
CasR_E...siraem	LSGKEINDLVYTHLNKFDNIRSFLEIMDELGL-----	545
FIG-018_GL0023397	INGKEINLLTTLINKFDNIRSFLEIMDELGL-----	533
FIG-025_GL0099734	IDGKEVNDLTTTLINKFDNIRSFIDIANQIGL-----	547
Ga0129317...chimera	LDGKEINLLTTLVSKFDNIGSIIIAATQIGI-----	479
EYZ-362B_GL0088915	LDGKEINELLSALLNKFDTIACWKCTFDLGI-----	537
uncultured_Ru_sp	LDGKEINLLTTLISKFDNIKEFLKMKSSAVD-----	541
R...flav...FD-1	LDGKEINLLTTLINKFDNIAQIKTAKELGI-----	541
tpg!DBYI01000091.1	LDGKEINLLTTLINKFDNIRSFIDIANFLNI-----	541
	. . * * : * : *	
037...__emb!OIZA01000315.1	AVYKDYSKYAIFR-----NA-----GKIAELEFAIKSI	577
emb!OCTW011587266.1	IINADYVDDYSLEF-----QS-----GTVARELMIKSI	528
k87...11092736	--DSIRFENEFALEN-----DD-----KGNMAGRIARQLRLIASV	541
BMZ--11B_GL0037915	-----VNFSKNYAVFN-----DC-----GSGAAGRIAEERLLIASI	526
BMZ--11B_GL0037771	-----VQFQPRFALEN-----NA-----SFAQNVAVQLRILIASI	526
BMZ--11B_GL0069617	VFSAKKGRNFSAPN-----EH-----GFAGRVADQLPVLIASI	506
Ga0099364...10024192	---KPQFTENYALFN-----EA-----AGQRAGEIAQNLRLIASI	573
530373_GL0023589	--PLDYQEKYRFFE-----RS-----REEAAGLRFIKSI	572
emb!ODAI011611274.1	---ETGLSDAFSEFE-----YS-----GEIAAELEPMTRSF	593
EMG...10003641	---KLEFTANYKMEIDPQKDPKDGHRVLSGS-----AKIAYYLRTINYI	610
CasR_P1E0...metageno	--AYSFVPAYAMFD-----HS-----RYIAGQLRVVNNI	542
emb!OIZX01000427.1	---EAELGPEYAMFT-----RS-----RAVAEQURVINSE	576
160582958...gene49834	--YETKYTEGYAATFK-----NA-----DKIARELQRQVNNM	623

FIG. 18II

```

gi|1198542314|gb|NFIR01000008.1|      602
emb|OGNF01009141.1                    548
Ga0129306_1000735                      575
MH0288_GL00822219                      635
FIG-022_GL0026351                      533
FIG-028_GL0185479                      535
250twins_35838_GL0110300              584
FIG-014_GL0226364                      530
250twins_36050_GL0158985              585
emb|OJMM01002900.1|                   557
emb|OGFN01002610.1|                   606
emb|OHCFO1000044.1|                   594
FIG-046_GL0077813                      599
pig_chimera                             601
emb|OIE01002196.1|                     542
O2_UC29-0_GL0096317                    529
CasR_Anaerobic_dig                      583
emb|OGDF01008514.1|                   559
emb|OGZC01000639.1                    585
Ga0224415_10048792_chimera            526
emb|OCVV012889144.1|                  527
Ga0187910_10006931                     539
R_flav_XPD_                              534
Ga0224415_10007274                    517

```

FIG. 18J

Ga0187911_10069260	-----SASYEADYKIFE-----DS-----GRVVEYLREINSF	523
Ga0187910_10015336	--KKVEFVENYSFFN-----SSNLLKEKTYLNKSENFTCKIVIELREINSF	609
ODAI_chimera	-NRVEFGDKYMFN-----KA-----GEICESLRSINSF	595
31009_GL0034153	---KCEFMEEYKFFN-----ES-----KDICLRKKEIHSF	521
DLF014_GL0011914...	--ECIFDKKFAFFD-----ES-----KNVCSLREINSF	591
CasR_R._albus	---EHSFTDNYKMFN-----DS-----KAICLQLQFINSF	577
Ga0187910_10040531	---DCTFKPEFAFFG-----Y.D-----RCRNISGELRLINSF	564
tpg!DJXD01000002.1	--EFKFKPEYADFF-----NMT-----NCRYTLEELRVINSI	505
CasR_E._siraenum	---DRFTTAEYSFFE-----GS-----TKYLAEELVELNSF	572
PIG-018_GL0023397	---ERTFTENYRFFE-----DS-----QKYRDELEIELNSF	560
PIG-025_GL0099734	--ECNLLDEYQFFN-----TS-----DTVKKELHVIKNL	574
Ga0129317_1008067_chimera	---NIEFIDDFKFFD-----RS-----KDISVELNIIRNF	506
EYZ-362B_GL0088915	---YAPFSKDFNFFE-----DS-----EQRAEEMNLVKSI	554
uncultured_Rn_sp	--VECLTAGYKLEN-----DS-----QRITNELFIVKNI	569
R._flav._FD-1	---NIAFVKNYDFFN-----HS-----EKYVDELNIVKNI	568
tpg!DBYI01000091.1	---DVKETKDYDFFN-----I.I-----CDYAGELNLIKNI	558
	;	
	;	
037_..._emb O1ZA01000315.1	ARMENKIE-----NAPKEFLKDALLAGVSSNDFDEKYEYFKTUVD---	620
emb OCTW011587266.1	SKMDFGLD-----NINLSFWYDDALRILGVSDENLPEVKREYF-----	566
k87_11092736	GKMKPDMT-----DARVLYKSALEILGAPPD-----EVS-----	571
BMZ-11B_GL0037915	GKMDAGLD-----SLKPLFIRFAIQILGGAPIS-----DEDV-----	558
BMZ-11E_GL0037771	GKMKPDLI-----EAKRPLYKAAIRMLCPPEK-----WEKYTS-----	559
BMZ-11B_GL0069617	GKMKPDLE-----GAKRALFRDAVETLGVSNWLNEDGIIIS-----	544

FIG. 18KK

Ga0099364__10024192	GKMKPDLG-----DAKRLYKAAIEMLGIDTE-----EYIS-----	604
530373__GL0023589	AKMKTRVA-----VKNDIKYVALKQYCDAGAVLGEINAEKIKAAAFGLG-----	617
emb!ODAI011611274.1!	ARMAAADP-----EAKRFMVVDGAKLLGNPKDTESEDEGIIRAIYGDAC	638
EMG__10003641	ARMESFEI-----KSDKTAINDAISLLGINSNEHRDEFITYTMAKHVVDK	655
CasR_P1E0_metageno	ARMKFPAI-----GAKREMYRAAVVLLGVDSPEAAAAITDBLLQI-----	582
emb!OIZX01000427.1!	ALMKKPOV-----NAKQQLYRAAVTLLGTEDPDGVTDEMLCIDP-----	615
160582958__gene49834	ARMTKPLG-----GVNYKCVMYNEAAKILGAKFMSKAELESVFNLDNHD-----	667
gi 1198542314 gb NFIR01000008.1!	ARMTFVP-----KTERIMFIEAAQILGYSNGEKELEGYVDALLD-----	642
emb!OGNF01009141.1	ARMEAPGA-----SSKRAMFVEAARILGDMRSKEELEEEYFDIL-----	586
Ga0129306__1000735	ARMTTEPF-----NSKLVMFEDAQAQLFGMSGGLVEHAEELREYLDNK-----	617
MH0288__GL0082219	ARMTTETV-----QKPCMFEDAADLLGMETDLEEVREALEAFYNKE-----	677
FIG--022__GL0026351	ARMIGFMEN-----DESTKIFMFEAAADILGTSYTDDELTDHLNL-----	573
FIG--028__GL0185479	ARMTGFMEN-----DESYKIFMFEAAADILGTSYTDDELTDHLNL-----	575
250twins__35838__GL0110300	AHLRPAIQ---EAQSSKDNADAAGTATLLIDAYNMLNIDIQL-----	624
FIG--014__GL0226364	SKMKKLTG---KE-----DKSINFPQVDDALELLEGLKIEKYEKKDDQGNFVLDQ-----	576
250twins__36050__GL0158985	ASKDFKKS---KKD-----SDESYEPQLYLDALALLGDVFSKYKQNRDGSVVIDDQ-----	633
emb!OJMM01002900.1!	SKMKPELS---KI-----GEALJLDAIDILGYKINKYKYDAAGNRLVDSN-----	599
emb!OGPN01002610.1!	SKMKNKVN---ID-----NDIYYLDAIEELLGRKIEKYQKDNKILLGTFD-----	648
emb!OHCP01000044.1!	SRMKKEMT---ID-----NUTIFLDALELLGRKIEKYQKDKNGDYVKDEK-----	636
FIG--046__GL0077813	SKMKKEID---VD-----NEIIFLDAFELLGKIEKYKDDSGNVLNRDD-----	641
pig__chimera	SKMKKEIA---ID-----NDTIFLDALELLGRKIDKYKDKKDATGKYLKDN-----	643
emb!OIE01002196.1!	AKMKKTL-----DFYNAQALKDAVTILGVSKKHQYLDMNSYLDIFYMF-----	584
O2_OC29-0__GL0096317	SRMKKAL-----DFYNGQALRDAIRVLGTSERHVDMDMKEYYVDKYL-----	571
CasR_Anaerobic__dig	TRMDNEI-----FPSEGVFLDAANVLGVKGDGDFDSNEFVGGDLHSD-----	625

FIG. 181L

emb!OGDF010089514.1	ARMCHSIAGLDVYKIYGSDDKFGHVSRRVYTDAAAYILSGEGERSEDNDGY-----VDD-----	611
emb!OGZC01000639.1	ARMGNPI-----SNLKEVMNVDAIQILGTEKSEQELQSM--ACF-----	622
Ga0224415_10048792...chimera	AQMGEPA-----ATLKLEMTADAIAIKILGTEKEDAELIKL--AES-----	563
emb!OCVV012889144.1	ARMGEPV-----ANTKRVWVIDAVKILGIDLSDDDELVM---ADS-----	564
Ga0187910_10006931	ARMGEPV-----ANAKRDMMIDAIAIKILGIDLDNNEIKKL--ADS-----	576
R...flav_XFD_	ARMGEPI-----ADARRAMYIDAIRILGINLSYDELKAL...ADT-----	571
Ga0224415_10007274	ARMGEPV-----ADARRAMYIDAIRILGIDLSDDDELKAL---ADS-----	554
Ga0187911_10069260	ARMTVDM-----EKIKRSAYKKALLILGSSKYSDEDLDAKVDEMLGV-----	565
Ga0187910_10015336	ARMTGDC-----KIKRSAFEDASQLLGYHDKTVNNLFEVLRKELESK-	652
ODAI...chimera	ARMESPA-----PAKRIMFVDAIRMFGFADVSDNDAEKFFYKENIEP--f--	638
31009_GL0034153	ARMKRPLD-----NKNVEKEMKYDAISILMKDSSSIENSIDVIFNTYIY-----	565
DLF014_GL0011914_	ARMQKPLN-----NVSVKREMYKDAIKILLKIDTRAEQCIDRILNEYIP-----	635
CasR...R...albus	ARMSKID-----DEKSKQLFRDALVILDIGNKDETWNNYLDSDIFK-----	620
Ga0187910_10040531	ARMQKPS-----AKKHVWYRDALRILGLDNGMSEALDQEVRRILQ-----	606
tpg!DJXD01000002.1	ARMQKPS-----ADARRIMYRDALRILGMNRPDEEIDRELEETM-----	545
CasR...E...siraenum	VKSCSEF-----INAKRIMYRDALDILGIESDKTEEDIEKMDNLIQ-----	614
PIG--018_GL0023397	AKSCSEF-----ISAKRIMYRDALDILGIAELSEDEIGRMIDNLIQ-----	602
PIG--025_GL0099734	TGMGYD-----VSIKHQMYKDAIDILGLHDMLSDEEINQLIDTNIL--F--	617
Ga0129317_1008067...chimera	ARMQAPV-----PNAKRAMQEDAIRILGSEEDIFSILDDMTG-----	544
EYZ--362B_GL0088915	ARMTKPD-----IOAKEQLFRDALDILGYEVGVSEALQKEIH-----	602
uncultured_Ru_sp	ASMRKPA-----ASAKLTFRDALTILGIDDKITDDRISGILK-----	607
R...flav...FD-1	ARMKKPS-----SNAKKAMYHDALTILGIPEDMDEKALDEELDLILE-----	610
tpg!DBYI01000091.1	ARMKKPS-----PSAKKNMYRDALTILGIFTEMSDEQLDAEIDKILE-----	610

FIG. 18MM

037__emb OIZA01000315.1	-----	620
emb OCTW011587266.1	-----	566
k87__11092736	-----	592
BMZ-11B_GL0037915	-----	576
BMZ-11B_GL0037771	-----	580
BMZ-11B_GL0069617	-----	563
Ga0099364_10024192	-----	654
530373_GL0023589	-----	617
emb ODAI011611274.1	-----	649
EMG_10003641	-----	711
CasR_P1E0__metageno	-----	582
emb OIZX01000427.1	-----	615
160582958__gene49834	-----	667
g1 1198542314 gb NFIR01000008.1	-----	642
emb OGNF01009141.1	-----	586
Ga0129306__1000735	-----	617
MH0288_GL0082219	-----	677
PIG-022_GL0026351	-----	573
PIG-028_GL0185479	-----	575
250twins__35838_GL0110300	-----	644
PIG-014_GL0226364	-----	598
250twins__36050_GL0158585	-----	655
emb OJMM01002900.1	-----	621

FIG. 18NN

emb!OGFN01002610.1!GKRLYTOEKYFNDFNFNA.....	667
emb!OHCFO1000044.1!GKVVYTKDYNFQDMFFEG.....	655
FIG-046_GL0077813GKVLRTKEYNNIRKMLCKK.....	660
pig_chimeraGKLYSKEYDDFQYMFYKD.....	662
emb!OIEN01002196.1!D.....D.....	585
O2_UC29--0_GL0096317D.....D.....	572
CasR_Anaerobic_digAN.....AN.....	627
emb!OGDF01008514.1!F.....F.....	611
emb!OGZC01000639.1F.....F.....	623
Ga0224415_10048792_chimeraL.....L.....	564
emb!OCVV012889144.1!F.....F.....	565
Ga0187910_10006931F.....F.....	577
R...flav_XPD...F.....F.....	572
Ga0224415_10007274F.....F.....	555
Ga0187911_10069260D.....D.....	566
Ga0187910_10015336DWRKRTDDEQQEYDRLN.....	670
ODAI_chimeraIYDNGDIED.....EW.....	650
31009_GL0034153EW.....EW.....	565
DLF014_GL0011914...EW.....EW.....	635
CasR_R_albusEW.....EW.....	620
Ga0187910_10040531EW.....EW.....	606
tpg!DJXD01000002.1!EW.....EW.....	545
CasR_F_siraenumI.....I.....	615
PIG-018_GL0023397L.....L.....	603

FIG. 1800

FIG-025_GL0099734I	618
Ga0129317_1008067_chimera	-----	544
EYZ-362E_GL0088915	-----	602
uncultured_Ru_sp	-----	607
R_...flav_...FD-1	-----K-----	611
tpg DBYI01000091.1	-----K-----	611
037_..._emb CIZA01000315.1	-----ADKHQKVSFFLNNVINNSRFKY	644
emb OCYW011587266.1	-----GKTKNLSAYIRNNVLENSRFKY	588
k87_11092736	-----KTVNFFRNYIAKNVITERSFY	614
BMZ-11B_GL0037915	-----PSNTERNFITNNVTKSRRFVY	597
BMZ-11B_GL0037771	-----KQVNPFRNFIAGNVIESRRFMY	602
BMZ-11B_GL0069617	-----RAINPFRNFIAANNVIESRRFMY	585
Ga0099364_10024192	-----RSLIDYEYLMPEINPFRNFVAKQVIESRRFMY	687
530373_GL0023589	-----KFNTTEEARVDHIFRNFIIINNVIINSRRFNY	646
emb ODAI011611274.1	-----EAFYVQEGLYGKEEKEFSPAYAFHDTSLRNFIAKNVIESARFRY	694
EMG_10003641	-----NLPAYFSEEDKEKLARYIVHSDGTYKFKFKESFYAIEELPNEGFRNFISNNVINNSRRFNY	771
CasR_P1E0_metageno	-----DPETGKVRPRSDSAPDTGLRNFIAANNVIESRRFTY	617
emb CIZX01000427.1	-----VTGKMLPPNQRRHGDGLRNFIAANNVIESRRFFQY	649
160582958_gene49834	-----YIYSPSGKKIPKNFRNFIIINNVIINSRRFLY	698
gi 1198542314 gb NFIR01000008.1	-----TKNKINDAKKKGFVRYIWNVVKISIRFRY	671
emb OGNF01009141.1	-----FDKSASAKKGFRRNFIRNNVYDSNRFKY	614
Ga0129306_1000735	-----MLDKTKLRLLPDKGVDTGFRNFIIISNVIESRRFRY	652

FIG. 18PP

MH0288_GL00822219K..MSLKWTKDQGEADTGRNFIIISNVLSKSSRFY	710
FIG-022_GL0026351	-----KKTRLNARSKADTGLRNFIIISNVLSKSSRFKY	604
FIG-028_GL0185479	-----KKTRLIARSKADMGLRNFIIISNVLSKSSRFKY	606
250twins_35838_GL0110300KY.....NKKGKKVDYDHKFRNFIIINNVLSKSKWFFY	677
FIG-014_GL0226364	-----SM---NDAEKSFVAKGHQKRFIIISNVVLNKRFLY	631
250twins_36050_GL0158985	-----ESGGIKYKSKPEYNNHQRRNFIIINNVLSKSKWFFY	691
emb OJMM01002900.1	-----EFGRVKYNKKKGFVNHRRNFIIINNVLSKSKWFFY	657
emb OGPN01002610.1	-----GNHKVRNFIANNVMSKQWFFY	688
emb OHCPO1000044.1	-----KNHRVRNFEVSNVLSKSKWFFS	676
FIG-046_GL0077813	-----DRRLNFIANSIKNNKWFYS	679
pig_chimera	-----SHRVNFIISNEVLSKSKWFFS	682
emb OIEN01002196.1	-----NRSGATGKNAGDHNLRNLFVSNVLSKSKWFFY	617
O2_UC29-0_GL0096317	-----DC---KDVHADHNLNRFIVITNVLSKSKWFFY	600
CasR_Anaerobic_dig	-----KKIINKINGTIEDRNLRNFIIINNVLSKSKWFFY	659
emb OGDF01008514.1	-----YVSSKLLGGADKNLRNFIIINNVLSKSKWFFY	641
emb OGZC01000639.1	-----FRDKNGKLLTGEHGMRFIGNNVLSKSKWFFY	655
Ga0224415_10048792_chimera	-----FKDENGKLLGNKQHGMRNFIGNNVLSKSKWFFY	596
emb OCVV012889144.1	-----FNDSEGNLLKCKGKGMRFIIINNVLSKSKWFFY	597
Ga0187910_10006931	-----FKDSNGKLLHKGKGMRFIIINNVLSKSKWFFY	609
R...flav_XPD...	-----SLDENGKLLKFKGKGMRFIIINNVLSKSKWFFY	604
Ga0224415_10007274	-----SLDENGKLLKFKGKGMRFIIINNVLSKSKWFFY	587
Ga0187911_100069260	-----YNQNGEKIKVRVDTGFRNFIIANNVLSKSKWFFY	598
Ga0187910_10015336	-----KHHYFKSGKKLPDTGLRNFIIINNVLSKSKWFFY	702
ODAI_chimera	-----KKGKLVHKKDEKKGKGVNFIIATNVLSKSKWFFY	682

FIG. 180Q

31009_GL0034153QKDNKNAKDFRNFIKKNVKSRRFIY	592
DLF014_GL0011914...	-----SKEDKGIKKDFRNFIIKKNIIKSNRFMY	662
CasR_R._.albus	-----LDKEGNKLGARHDFRNFIANNVKSRRFKY	651
Ga0187910_10040531	-----IGADGKPIKANKGFRNFIAASNVISSRRFY	637
tpg!DjXD01000002.1	-----PVGADGKFIKKGQGRNFIAASNVISSRRFYH	576
CasR_E._.siraenum	-----DANGD.....KLLKNNGLRNFIASNVIDSNRFKY	645
PIG--018_GL0023397	-----DANGERISKNGENGLRNFIASNVIDSNRFKY	635
PIG--025_GL0099734	-----GADGKPVKMAKKGGRNFIIISNVLKSSRRFY	650
Ga0129317_1008067_chimera	-----YDKSGKLLAQSKKGRNFIIINNVVSSRRFKY	575
EYZ--362E_GL0088915	-----QYFQKGAHDFRNFIINNVIINSTRFIY	629
uncultured_Ru_sp	-----LKEKGIHGLRNFIITNNVIESSRFVY	634
R._.flav._.FD-1	-----KTDPTGKPLKGNPLRNFIANNVIENSRRFIY	643
tpg!DBYI01000091.1	-----KINPVIKTEKGNPFRNFIAANNVIENKRRFIY	643
	* * *	
037_...emb OLZA01000315.1	VVKYINPADINGLAKNRYLVKFLSKLPE---E-----QIDSYYKLFSEEE-----	688
emb OCTW011587266.1	VIKYIHPSDVQKACNKAIAGFVLRMPD---T-----QIKRYYDSLINKGA-----	632
k87_11092736	LVRyakftAVRKLMSNFKIVRYVLRLEPE-----KQVASYYSAIWTQSES-----	659
BMZ--11B_GL0037915	LVRyTKADAVRAVMKNGKIVRYVLRLEPE-----TQVDSYYKSVVDSSE-----	641
BMZ--11B_GL0037771	LVRYSKPKAVRALMQNRSIVNYVLRLEPS-----EQIKTYSRVFEEDF-----	645
BMZ--11B_GL0069617	LARYAKPKSVRELMKHEEIRYVLRLEPE-----KQIDTYFSNISEGDE-----	629
Ga0099364_10024192	LVRyTKPKTVRALMSNFAIVHYVLSRIAD--I--QDHHMTESQIDRYYQNLPOYNEQ----	740
530373_GL0023589	VMRFMNPKSARRIMQCESLVAFLKIDIP---T-----QIERYCKSVNHPFDS-----	691
emb ODAI011611274.1	VIRYVSPEIARKYARQEAIVRFALHRVPL--L-----QLRRYYQSCCGPKKD----	739

FIG. 18RR

EMG_10003641	IMRFCNPEKIANIGKKNKVLISFALSGLAEKTD.....MIAKYRVVFCDFRID.....	817
CasR_PLEO_metageno	LLRYMTFEQARVLAQNEKLI AFVLSVTPD--T-----QLERYCRICGREDI-----	661
emb:OIZX01000427.1:	LIRYSDPAQLHQLASNKKLVRFVLSVIPD--T-----QINRYYETCGQTRL-----	693
160582958_gene49834	LIRYGNPEKIRKIAINPSIISFVLKQIPD---E-----QIKRYYFFCIGKRI-----	742
gi 1198542314 gb NFIR01000008.1:	LVRVADFKKVRFAANKKVVAVFLKIPD--D-----QIRAYYNSCFQNSDSSS	719
emb:OGNF01009141.1	LTRYIDTSSVKAFSNKKALVKFAIKDIFQ---E-----QILRYNSCFGASER-----	659
Ga0129306_1000735	LVRVYCEPRAVRDYMSCRPLIRLTLRDMFD---T-----ILRRYEQSVGAAT-----	696
MH0288_GL0082219	VVRVYGNFKKLRVVMQNEAVVKFVLDLPE---E-----QINRYSENILGLDA-----	754
FIG-022_GL0026351	LIRYSNPKKVKVKAIAENKKLVRFVLDKIPD---P-----QIIRYCNTCGF-KV-----	647
FIG-028_GL0185479	LIRYSNPKKVKVKAIAENKKLVSFVLDKIPD---T-----QIIRYCNTCGF-KV-----	649
250twins_35838_GL0110300	IARYVKPADCARMSNKKMIEFALRDIPE--T-----QIKRYYTYITGNEA-----	721
FIG-014_GL0226364	VAKYGSPTACNQMMKNQIVRMALESISE---D-----QLRKYARRVYDNKEH-----	676
250twins_36050_GL0158985	VVRVYNRPSSCRRLMKNKEILRFVLDKIPD---S-----QVRRYFKAVQGEA-----	735
emb:OJMM01002900.1:	VAKYNRPSECQEMSKKLIALLVKDVPE--T-----QIARYYQSVTGGRT-----	701
emb:OGPN01002610.1:	VVRYNKPAEQIIMPNKTLVKFILDLELFD---M-----QIQRYYSVFGDNN-----	732
emb:OHCP01000044.1:	VVRYNKPAEQALMPNSKLVKFDLDELFD---S-----QIEKYYISVFGKKS-----	720
FIG-046_GL0077813	IVRYNQPSECQAIMKNKILVKFALDELFD---L-----QIQRYFVALYGDDE-----	723
pig_chimera	IVRYNQPSECRAIMKNKTLVKFALDELFD---L-----QIQRYFVALYGDDE-----	726
emb:OIE01002196.1:	LSRYSNLAEVKKLAQNPQLVQFVLSRIEF--S-----LICRYYESSQGIS-----	660
O2_UC29-0_GL0096317	LARYSDMATVSKLSKNKEVVKFVLRVYDP---K-----QIERYYRSVCDVR-----	643
CasR_Anaerobic_dig	IARHMNTHYVKQLANNETLNRFLVLMGDI---A-----KIINRYYESISGNT-----	703
emb:OGDF01008514.1:	TVRYMNPKRAKLVQNDALVVALSGIPE--T-----QIDRYYKSCIEKR-----	684
emb:OGZC01000639.1	LIRYGNPQKHLHLSQNETVRFVLSRIAK---NORVQGMNGKNQIDRYYETCGGHN-----	708
Ga0224415_10048792_chimera	LIRYGDPAHLHKIATINKNVRFVLRGRIAD---MQKKQGGKGNQIDRYYEVCGVGNK-----	649

FIG. 18SS

emb OCVV0128899144.1	LIRYGDPAHLHEIAKNEAVVRFVLGRAD----IQKKQKDGKNQIDRYYEICVGKD-----	650
Ga0187910_10006931	IIRYGDPAHLHEIAKNEVVVRFVLGRAD--IQKKQKGGKNQIDRYYEICIGNG-----	662
R_...flav_...XFD_...	LIRYGDPAHLHEIAKNEAVVRFVLGRAD--IQKKQKGGKNQIDRYYETCIGKD-----	657
Ga0224415_10007274	LIRYGNPVLHEIAKNEAVVRFVLGRAD----IQKKQKGGKNQIDRYYETCIGKD-----	640
Ga0187911_10069260	LIRYCHPRKIRNLAGNAALIEYQLRLPE--L-----QILRYYEACTEPIK-----	642
Ga0187910_10015336	IVRYADFKKIRKCTENNELLKFAFKVFD---S-----QVDRYYNICVTNK-----	745
ODAI_...chimera	LVRVYANTEKIALAKNETIVRFVIKKLPA---A-----QIVRYYIACNREIN-----	727
31009_GL0034153	LIKYSNFTVNRKFASTNVIKFVLGLPD---K-----QIERYYKSCGLSD-----	635
DLF014_GL0011914_...	LVKYSNFTDVRNLASNRNVKFLNALIED---T-----QIDRYYKSCIQHY-----	705
CasR_R_...aibus	LVKYSSADGMILKLTNEKLIQFVLKLEPE---T-----QIDRYYESCGLD-----	693
Ga0187910_10040531	LVRVYNNPHKTRMTAQAQNEAIVRFVLSLIPD--E-----QIRRYDVCVCRDPK-----	680
tpg DJXD01000002.1	LVRVYNNPHKTRILVKNFNVKFLVLEGIPE---I-----QIKRYFDVCKGQE-----	619
CasR_E_...siraenum	LVRVYGNPKKIRETAKCKPAVRFVLEIPD---A-----QIERYYEACCPKN-----	688
PIG-018_GL0023397	LVRVYGNPKKIRETAKCPPAVKFLDGLIPD--S-----QIERYYNSCFED-----	678
PIG-025_GL0099734	LVKYCNPKKIRKIANNEKIVKFLGRIFE---I-----QIERYYYSCPELK-----	694
Ga0129317_1008067_...chimera	IVRYSNPQKIRKLLANNSVVVGFVLGKLPD---A-----QIESYFNWSCLPN-----	617
EYZ--362B_GL0088915	VIKFCNPKIAGQIIRNENILRFVLQRMPO---E-----QIERYYHACTKN-----	671
uncultured_Ru_...sp	LIKYANAQKIREVAKNEKVVVMFVLGGIIPD---I-----QIERYYKSCVEFP-----	677
R_...flav_...FD-1	LIKFCNPNVVRKIYVNTKVTFFVLKRIED--A-----QIERYYKSCIDS-----	685
tpg DBYI010000091.1	VIKFCNPKVVRKLVNNTKVTFFVLKRMPE---T-----QIDRYFESCIEG-----	685
	: . : : : *	
037_...emb OIZA01000315.1	-----PSCEEKIKLLTKKLSKLNFOLENNK-----	715
emb OCTW011587266.1	-----TDIQQAQAKALLDCITGISFDAQKDKHLHKS-----KE	665

FIG. 18TT

k87_11092736	-----NSNEMVKLIEMIDRLTTEIAGFSFAVLKDKKDSIVSA-----SR	698
BMZ-11B_GL0037915	-----AGNGKPKIKKIKALAEKLGESFDLIAENRDIIDNA-----KL	699
BMZ-11B_GL0037771	-----SDTEAEIDFLVNLSEFSFEFFISNQQTILNNS-----KRGFSP	684
BMZ-11B_GL0069617	-----SEISRKIDITLIKALTGFSFNSLFINRDKKIVENT-----KL	664
Ga0099364_10924192	-----QHKNSLETKIDALADYLCKYTFEKNVLKQKNGIV-L-----NT	778
530373_GL0023589	-----ANPNYAHMKEQLTYKLLQVQFCNFSKVSNSSHQ-----	724
emb!ODAI011511274.1;	-----PDAECVDFLAGVVRVDFANFIDVPTGDSK-----	771
EMG_10003641	-----DQKTMEDYLVNKLTKISYTEFLNVNCKANAE-----	848
CasR_P1E0_metageno	-----TGRFAQIRYLTAQIMGVRYESFTDVEQGRG-----	692
emb!OIZX01000427.1;	-----AGRAAKVEFLDMIAAIRFDQFRDYNQKERG-----	724
160582958_gene49834	-----DDVTLMRDELCKMLQSVNFEQFSRVNKKQNAK-----	774
gi 1198542314 gb NFIR01000008.1	NNSNASWDADSNKRDISVDMRKALTEKITGLNFGDFEESKKGIR-----	765
emb!OGNF01009141.1	-----YYNDGMSDKLVEAICKINLMQFNQVIGQADRN-----MLPE	695
Ga0129306_1000735	-----VDRERILDITLADKLLSLRFTDFENVNQRANAE-----	728
MH0288_GL0082219	-----AAKEKKVSELAPFLRALSLDFRDFVEQSANAA-----	786
PIG-022_GL0026351	-----GERQENINTLAKAITNFSFSAFEKVKQSANVE-----N-----	680
PIG-028_GL0185479	-----GKRQENINTLAFAITYFSESFNFEKVKQSANVE-----N-----	682
250twins_35838_GL0110300	-----LGDASLKGVIIEQLHAFSIKNTLLSIKNNM---G-----EGEY	756
PIG-014_GL0226364	-----FQQTADIRENLAKKLNFNIGAEIKTVGNL-----SEKD	712
250twins_36050_GL0158985	-----YASAEAMRTRLVDAISQFSVTACLDEYVGM---I-----DKEF	770
emb!OJMM01002900.1;	-----QANSEAMRMTLTKLLHEFSIKNVLSDVGTM---T-----ASEN	736
emb!OGPN01002610.1;	-----MPAVDEMRRKLLDKINQFSVRGFLDELDELIVLM-----SDEE	769
emb!OHCP01000044.1;	-----SSSNEEMRRELKKLKCDFSVRGFLDELIVLL---S-----EDEM	755
PIG-046_GL0077813	-----LPSYGENMKILLKLLKLNDFSIIRDFDFEIGSL---F-----D-EM	757

FIG. 18UU

```

pig_chimera          761 .....LPSYGE MRKILKKLHDFSIKFLDEIVLL--S-----DLDM
emb!OIEN01092196.1  699 .....SEGITIDEQIKKLTGIVDMNIDSFNINNGEIGMR-----Y----SK
O2.UC29-0_GL0096317  683 .....NISADEIEKLVDIINYFNIDELEQVNAEIGTG-----HSGRKV
CasR_Anaerobic_dig   737 .....PNIEVRSOIDLVKRLRSFSFEDLNDVKQKVRPG-----
emb!OGDF01098514.1  723 .....SFNFDLNEKIAALSEMITLTKIDDFEDVKQNFKNA-----NYE
emb!OGZC01090639.1  743 .....SWSVSEEEKINFCLKILLNMSYDQFQVQKQSGAEI-----
Ga0224415_10048792_chimera  694 .....PIKKTIEEKIDALIDIIIVNMNYDQFEAKKAVIENQ-----NRGKTFEKN
emb!OCVV012889144.1  697 .....RGKTVSQKLDDELTKIIVNMNYDQFEQKKEIIENTRTKTYMRYDEEQ
Ga0187910_10006931   696 .....YGKSVSEKIDALTKVIIINMNYDQFEAKRVIENT-----
R_flav_XPD_          691 .....KKSVSSEKVDALTKIITGMNYDQFKRSVIEDI-----
Ga0224415_10007274   674 .....SSKSVAEKVNALTKIITGMNYDQFDSRNVIENT-----FSDAEKK
Ga0187911_10069260   685 .....PIARTMDEKIGILIDLIVKMDFSQFEDVQQNDRVRV-----
Ga0187910_10015336   779 .....ITNATREKIERLVPIKSMNLSKVAIVKQDKQ-----
ODAI_chimera        762 .....INSVLTSDDMAESLVKLVKMGDMDFGKIKGVNDDR-----
31009_GL0034153      667 .....GIGKTEWVRELSEIITGIDYSDFLNVKQSAKS-----
DLF014_GL0011914...  738 .....SNEVDKETQIQELIDIIITGIDYLDFLDVRQSYR-----
CasR_R_ailbus       727 .....NAVVDKVRIEKLSGLIRDMMKFDDFSVKTSNKA-----
Ga0187910_10040531   721 .....LPRSSSREAQVDILITGIIITDVNYRIFEDVPQSKKINK-----DRPD----
tpg!DJXD01000002.1  660 .....IPPTSDKSAQIDVLAARISSVDYKIFEDVPQSAKINK-----EDFS----
CasR_E_siraeum      726 .....TALCSANKRREKRLADMIAEIKFEMFSDAGNYQKANV-----IS-----
PIG-018_GL0023397   716 .....TAICSADKRRRELAEAMISNISFDWFKDAGNVQRVNA-----ED-----
PIG-025_GL0099734   731 .....RAYYPGREAIDDLTKIILINMKFDDFNNKQSANV-----ED-----
Ga0129317_1008067_chimera  652 .....RVYSTPDKARESLRDLHNI SFNDFADVQKQDDR-----A-----
EYZ-362B_GL0088915  706 .....PSLCTHDEKIQCLSNIMEVRYKKFCNVQAQKSRD-----T-----

```

FIG. 18V


```

uncultured_Ru_sp          711
.....DMNSSLGVKRSELARMIKKNISFDDFNKVKQAKG.....
R_...flav_...FD-1       718
-----EMNFPTEKKITIELAGKCLKDMNFGNFRNVRQSAK-----
tpg!DBYI01000091.1|    720
-----NLNFTTEKKIEKLAEMIKNIKIFEEFRNVKQKVRD-----N-----

                                :
                                :
                                :
037_..._emb|OIZA01000315.1| 772
-----IPNVEKERKKAIIITLYFTIIVYLLVKNLVNINGLYTTIALYFVERDGFYKQICEKKLI
emb!OCTW011587266.1|    724
KSPQRSADRRERKAMITLYYTIIVYFVKQMLHINSLYTIGFFYLERDQRFIYSEAKKEN-
k87_11092736           758
ESRAVNLEVERLKNLITLIMSIAIYIAVKS LVKNARYIAYSALERDLYFFNEKYGEEFR
BMZ--11B_GL0037915     740
DEEKNVEIERLKAVTGLYLTVAFIAIKNLVKANARYIAYSAFERDSELLKEKLGEEFF
BMZ--11B_GL0037771     741
NRRETAEIEIERLKAITGLYLSVAYIAIKNLVKANARYIYAFVFERDKELVKAADAR.....
BMZ--11B_GL0069617     722
DTSNKNVKIERLKAITGLYLTVAFVAVKNLVKANARYIYAFSAFERDYGFMKSHDPD----V
Ga0099364_10024192    838
KSAIKNVEIEIHLKALITGLYLTVAIYAVKNLVKANARYIYAFSIFERDYALFEKKGKIDIL
530373_GL0023589      779
-----AVEKERLKAIVGLYLSVLYRIVKSVIRINTSYITAFAAAYERDCVILNKKMGKNLY
emb!ODAI011611274.1|  828
-----SEQEKKQKQIAIVGLYLTIVVYIVKLVNNSRYVMAFHILERTVLLLEGKPLFVGG
EMG_10003641          903
-----KNKEKORSQKLIIGLYITLLYEIVKNLVNIINSRYNIYAFQECNDNSIMILQCGYDER--
CasR_P1E0_metageno    748
-----DNPKKERYKALIGLYLTVLVLAVKNMWCNARYIYAFYCRDRDIALYQKEVVCWYDL
emb!OIZX01000427.1|  780
-----ANTQKERYKAMGLYQTVLVLAVKNLVNINARYVMAFHCVERDMFLYDGEIIDPKG
160582958_gene49834   830
-----QNPNGEKARLQACVRLYLITVPLYLETKNMVNIINARYVLAFAHCLERDHALCENSEKLN---
gi|1198542314|gb|NFIR01000008.1| 818
-----KEESDKIIRLTLTVLVLVQKNLIYVNSRYTAFHCAERDEMLYNGETIDNNK
emb!CGNF01009141.1    755
EKKKANAQKEKYSIIRLTLTVCYLFFKNLVVYNSRYYSAFYNLEKDRSLFEINGELKPT
Ga0129306_1000735     784
-----RNREKQKMMGIISLYLNVAIQIVKNLVYVNWARYTMAHYCAERDETELLNAAGEGNL
MH0288_GL0082219      842
-----RNARKARYQGAMSLYLNVIYQIKNMVYINSRYTLAFAKYEHDA SLLLDNWNVDYDPY
PIG--022_GL0026351    737
-DTFD AIDKAQKQGLISLYLTVLVLFFKNMVYVNSRYVLAFAHCLERDSQLLKITSGYN----
PIG--028_GL0185479    739
-DTFD AIDKAQKQGLISLYLTVLVLFFKNMVYVNSRYVLAFAHCLERDSQLLKITSGYN---

```

FIG. 18WW

250twins__35838__GL0110300	KIQQIGSSKEKKAIVNLYLTVAYLLFKSLVKVNIRFSAFGCLERDLVLQKXS-----	810
PIG-014__GL0226364	FRSKNALKDKSKSLIGLYLTVAYLLFKNNIKINTRFSAFAYERDMFLLQQ-----	766
250twins__36050__GL0158985	ASQRAVDSKEKRAIRLYLTVAYLLFKSMVKVNTRFSAFVSLERDYLLIDG-----	824
emb!OJMM01002900.1!	KRQIENSRKERMAIVKLYLTVVYLLAKSLVKVNIRFSAFSAYERDVSLADENELIA--	795
emb!OGFN01002610.1!	SKRNKSSEKEQKSLIRLYLTIAYLLFKSMVKINTRFSAFACAMYERYALLCQS-----	823
emb!OICP01000044.1!	KQKDKFSEKEKKSLIRLYLTIIVYLLFKSMVKINTRFSAFACATYERDYILLCQS-----	809
PIG--046__GL0077813	KRKDKFCESDRKCLFRLLYLTAYLLFKSMVKINTRFSAFACATYERDYALLCAS-----	811
pig__chimera	ESQDKYCEKEQKSLFRLLYLTAYLLFKSMVKINTRFSAFACATYERDYALLCAS-----	815
emb!OIEF01002196.1!	ATPQSIERRNQRFVCGLYLNVLYQIENLNMVNAVRYLAFAFERDALMNFLEECK--	758
O2. UC29-0__GL0096317	ASKRSVQERNQACVSLYLVNLYQIVKNLNVNNSRYSIAFAMTERDMIVEGKIEAIEF	743
CasR__Anaerobic__dig	--TNESIEKEKKALVGLCLTIQYLVYKLVNINARYYTAFYCLERDSKLGFGVDVWR--	794
emb!OGDF01008514.1!	AKKNQRI SKERYKACIGLYLTVLVLICKNLVKINARYSIAIGCLERDTQLHGVDKFAA--	782
emb!OGZC01000639.1	--TAEKRRKERYKALISLYLTVLYQLKKNLVNINARYYIAFHCLEERDAILYSSKFNISIN	802
Ga0224415__10048792__chimera	KYKRDNAEREKFKKILISLYLTVIYHILKLVNNSRYILGFHCLERDKQLYIEKYKND---	752
emb!OCVV012889144.1!	KLRKDNAEREKFKKILISLYLTVIYVYKLVNINSRYVIGFHCLEERDAQLYSEKYNKE---	755
Ga0187910__10006931	---GRDNAEREKFKKILISLYLTVIYQILKKNLVNNSRYVIGFHCVERDAQLYKEKGYDINT	754
R__flav__XPD__	---GRENAEREKFKKILISLYLTVIYHILKLVNINARYYVIGFHCVERDAQLYKEKGYDINL	749
Ga0224415__10007274	---GAGNAEREKFKKILISLYLTVIYHILKLVNINSRYVIGFHCVERDAQLYKEKGYDINL	732
Ga0187911__10069260	EKIRKMKRQRYQSIISLYLTMLYLVKNLVNINARYYVMAFQAWERDNYLLQLSGKE---	743
Ga0187910__10015336	-----DNVEKQKQLAIMSLYLNILYQIAKNLVVNSRYVMAFHSLERDSQMLEDAYYDV----	833
ODAI__chimera	-----SGDKEMKKAIVTLYLTIYHILTKNLVYVNSRYVMAFHCLEERDMNLFSGQKQALR	817
31009__GL0034153	--KEEKEDKAKHQAVTLYLTIYHILTKNLVNVNSRYVIAVHCIERDSELYKTLYEKLK	725
DUF014__GL0011914__	-----NEDKSKQAVVTLTYLTIYHILTKMLVNVNSRYVIALHCLERDSKLYGTYKLSKKN	793
CasR__R__aibus	--GDNDKQDKAKYQAIISLYLVMVLYQIVKNMIVVNSRYVIAFHCLEERDFGMYGKDFGKY---	764

FIG. 18XX

Ga0187910_10040531	777	ANDRMTLKKQRYQAIIVSLYLTVVNYLVTKNLLVYVNSRYVMAFHAFHALERDAYLYGIINI-----
tpg!DJXD01000002.1!	718	RNFSDAKKQRYQAIIVSLYLTVVXLLTKNLLVYVNSRYVIAFHCLERDAFLHGVTLPKM--
CasR_E_...sraeum	784	-RTSEAEIKRKNQAIIRLYLTVMYIMLKNLVVNNARYVIAFHCVERTKLIYAESGLEVG-
PIG--018_GL0023397	774	--KSRNAEIKRKNQAIIRLYLTVMYIMLKNLVVNSRYVIAFHCVERTKLIYKLSGLTIE--
PIG--025_GL0099734	790	-SSKEAQEKMKYQTIISLYLTVCYHLVKNLVNINARYAMAFHSLERDARLYGIGSEKDY
Ga0129317_1008067_chimera	705	--TPEKVEKERKAIIGLYLTVMYHLVKNLVVNSRYVMAFHCLERDAMHYDVS-----
EYZ--362E_GL0088915	763	-----TKQMEKERKAIIRLYLTAIYLVVKNLVVNNARYVIAFHCLERDADLFGIQLKQTLIP
uncultured_Eu_sp	767	----RENVAKERKAVIGLYLTVMYLLVKNLVVNNARYVIAHCLERDFGLYKELIPEL-A
R_...flav_...FD-1	768	-----ENMEKERKAVIGLYLTVVYRVVKNLVVNSRYVMAFHSLERDSQLYVVS-----
tpg!DBYI01000091.1!	772	----SQAVEKERKAIIGLYLTVIYLLVKNLVVNSRYVMAFHCLERDAKLYGVQ-----
037_...emb O1ZAO1000315.1	803	-----ETLKKKKKKQLYNDDDDYLLPELPSGSKYRE-----
emb OCTW011587266.1	756	-----KNPSKNSYLNDERSVTAIPIFSEIMKRIEKNE-----
k87_11092736	803	-----LHFIP-----YELNGKTCQFEYLAALKYVLARDEETLKRKCEICEEIKVG
BMZ--11B_GL0037915	787	-----YKLCIEFKN-----KKGETGYNNAFALKYIYDRDEDIWRKKEATHRGRDEN
BMZ--11B_GL0037771	782	-----IQTTIENIKFTNYFCLIQYVYLDREDEKFFPGDPRDKEAFFE
BMZ--11B_GL0069617	772	-----AQCLAYEEIMDGKSKTKYNEIYALTEYVYKKEEENDYHPEPCQFFDKEA
Ga0099364_10024192	888	-----EKVVKPFKYIDKGEEREKKNFFALTEYLLDKDNSLRYQWNNLSDENK
530373_GL0023589	804	GL-----KIDEMGIINDFMHPMTVAPSNPH-----
emb ODAI011611274.1	851	MK-----AEDPFLITDGYVSRQDAYVRK-----
EMG_10003641	919	-----AVQESKLTKKPIISNQK-----
CasR_P1E0_metageno	775	EE-----DKKSGKQKQVEDYTAITRYEVSQGY-----
emb O1ZX01000427.1	811	ES-----VSAFLAVNGKKGVPQYLLLTQLFERRDY-----

FIG. 18YY

160582956_gene49834DDSYEMANKFQMVRRKAKKEQYKEXKCK.....	859
gi 1198542314 gb NFIR01000008.1	EK-----GSEKDRKFAKQFIMEHS-----	838
emb OGNF01009141.1	GK-----FDEGHTGLVKLFLDNGW-----	775
Ga0129306_1000735	LRDRSWPARLHLFRALARRRDRVEVMERDVARGPEAVNRDENLG-----	830
MH0288_GL0082219	-----KIVFSDLTFEDSLQKCK-----	858
FIG-022_GL0026351	-----NSSYNNLITNYFIDNKY-----	753
FIG-028_GL0185479	-----NLSYNNLITNHFLEKYY-----	755
250twins_35838_GL0110300	-----EKKFDAIINEILLEDDDKIRKCEDKERAQAKTL	842
FIG-014_GL0226364	-----KKEKGRFDWLALTQKFLSEDKNYYFEHDLRKEINKI	804
250twins_36050_GL0158985	-----KKSSTYTGEDMLALTRKFGVEDAGLYREWKEKNAEAKDK	864
emb OJMM01002900.1	-----LANNEDDWKKGNYVFALITKHEWENDEPIFDKYNMALQQIRSI	838
emb OGPN01002610.1	-----EMKGGPWGGGAQALAVTRKFLNHDRVDFDRYCAEAEIARL	864
emb OHCPO1000044.1	-----EKAERAWKCATAFALTRKFLNHDKPTFEQYYTREFEISAM	850
FIG-046_GL0077813	-----NKQERAWSSGATALALTRKFLNQDKLIFEKHYAREGEISKL	852
pig_chimera	-----NKQERAWSSGATALALTRRFLNQDKLIFEKHYAREGEISKL	856
emb OIEO1002196.1	-----KNKKRSGGFSFIEMIQFFLDKKLKFVAT-----	787
O2_UC29-0_GL0096317	FKDDP-----DAGKKSKKQIAYFTKYIALITEEYLDKGYIKCAT-----	781
CasR_Anaerobic_dig	-----DFESYALTNHFIKEGY-----	811
emb OGDF01008514.1	-----YMRDVFIAKGI-----	795
emb OGZC01000639.1	-----LKKRYTALIEMLIGYETDEKAR-----	824
Ga0224415_10048792_chimera	-----KLDGFVALTKFCUGDEERFEDL-----KAK	777
emb OCVV012889144.1	-----IDKKISSLTKFCUGSEDEKLLK-----AL	779
Ga0187910_10006931	-----NNLESKGFSTVTKLCVGIADDDPVKYNVEIELKER	790
R_...flav_XPD...	-----KKLEKGFSSVTKLCAGIDETAPDKRKRKDYKEMAER	785

FIG. 18ZZ

Ga0224415_10007274KKLKKGFTSVIKLCAGIDECK.....DVEKEMTER	764
Ga0187911_10069260AAEAYLNLTRHFIEPLDGAKP.....	764
Ga0187910_10015336KRGYCDLSIVLLFGVDDLQNRNRGSSXYLRDN	865
ODAI_chimera	NTGDY.....PADN.....ADLRRRDIKYCVLTKAHKCFEYKSRND.....	855
31009_GL0034153	PEQPE.....KLYKIKPKLQPAQYFLFTKLLQNDFFGE.....	761
DLF014_GL0011914_NKKLERIKPKYHELTKEYFDNRYFDRK.....	821
CasR_R_albusYQGCCKLTDHFEEEKYMK.....	803
Ga0187910_10040531KGDYRKLTDNLLADENYKFF.....	797
tpg!DJXD01000002.1NKKIVYSQLTTHLLTDKNYTTY.....	740
CasR_F_siraeumNIEKKNTNLIMAYMGVYKLENG.....IYKT.....	809
PIG-018_GL0023397GNIDKNRLLTATILGITSDRAPSGEIVA.....	804
PIG-025_GL0099734	V.....EKLKKEQAVLVEKLLSDCPETAG.....	814
Ga0129317_1008067_chimeraLDNYRDLIRHLISEGDSSCN.....	725
EYZ-362B_GL0088915	D.....RKKKSDYKLVADQCGDNFENKP.....	787
uncultured_Ru_sp	S.....KNLKNDYRILSQILCELCDKSPN.....	791
R_flav_PD-1VDNDYLALIDILYKEGDNRSR.....	789
tpg!DEYI01000091.1NIGGDYALALIAKLCAEGDDYGKKLSEAKQ.....	801
037_>_emb OIZA01000315.1	803
emb OCTW011587266.1	756
k87_11092736	CEKHKKNANPPYEQEWIDKKKALNSE--RKACERRLHF-----STH	844
BMZ-11E_GL0037915	GKLDYDPPD.....	796
BMZ-11B_GL0037771HL.....	784

FIG. 18AAA

BMZ_11B_GL0069617	CRKHLD	778
Ga0099364_10924192	QALRKHLDKKE	899
530373_GL0023589		804
emb:ODAI011611274.1		851
EMG_10003641		919
CasR_PIE0_metageno		775
emb:OIZX01000427.1		811
160582958_gene49834		859
qi 1198542314 gb NFIR01000008.1		838
emb:OGNF01009141.1		775
Ga0129306_1000735		830
MH0288_GL0082219		858
FIG-022_GL0026351		753
FIG-028_GL0185479		755
250twins_35838_GL0110300	P.....RELAQE	849
FIG-014_GL0226364	SD.....RDERKK	812
250twins_36050_GL0158985	YF.....DKAERK	872
emb:OJMM01002900.1	V.....DPNERR	845
emb:OGPN01002610.1	P.....S-EERK	870
emb:OHCP01000044.1	P.....Q-EKRR	856
FIG-046_GL0077813	P.....K-EERK	858
pig_chimera	P.....K-EERK	852
emb:OIEN01002196.1EA	789
O2_UC29-0_GL0096317ER	783

FIG. 18BBB

CasR_Anaerobic_dig	811
emb!OGDF01008514.1	795
emb!OGZC01000639.1	829
Ga0224415_10048792_chimera	811
emb!OCVV012889144.1	813
Ga0187910_10006931	824
R_...flav_XPD_...	819
Ga0224415_10007274	798
Ga0187911_10069260	764
Ga0187910_10015336	925
ODAI_chimera	855
31009_GL0034153	761
DLF014_GL0011914_...	821
CasR_R_...albus	803
Ga0187910_10040531	797
tpg!DUXD01000002.1	740
CasR_E_...siraenum	811
PIG-018_GL0023397	806
PIG-025_GL0099734	814
Ga0129317_1008067_chimera	725
EYZ-362B_GL0088915	787
uncultured_Ru_sp	791
R_...flav_PD-1	789
tpg!DBYI01000091.1	803

FIG. 18CC

037_--_emb OIZA01000315.1	-----ET-----KNLKLPEKDKRDIEMKKYLPENDKDKREYNKFFKQYRN	841
emb OC1W011587266.1	-----NKGELEDFEALWNSC---GKTSRLKEKEDVLLIYARYISFDHALKNYKMILNSYRN	807
k87_1i092736	WAQYATKRDENMAKHPQKWYDILASHYDELLAL-----QATGWLAIQARN	889
EMZ-11B__GL0037915	-----RRKHYPKRRWLNILKDNLAEMERI-----DPTGMLAAQMRN	832
EMZ-11B__GL0037771	-----PKTRRHFSKQWRWLNELKADAKSS-----QATGLLLREARN	821
EMZ-11B__GL0069617	-----SIRRHFTKKWRDIFRKEIDEAVKI-----HFTGYLGVWARN	814
Ga0099364_10024192	-----IRSQRHFSQYWUDIFARQIENAKKT-----SESGYLLTAARN	936
530373_GL0023589	-----NDFINEFKLSKEVCAVKNKFTL-----CTNPKYQVYRN	838
emb ODAI011611274.1	-----RIGENKRNHRGLNCVLENRNALGS--D-----PASTDAAASLIWSYRN	893
EMC_10003641	-----LNSYSCRYLTHNISQLD-----RCNDFIRQYRN	947
CasR_P1E0_metageno	-----LNKHACGYLRSNMNG-----ISNSLLIAYRN	801
emb OIZX01000427.1	-----LKRSAEQIQHNMEN-----ISDRLLREYRN	837
160582958_gene49834	-----KQETGTAHTKIEKLNQCIAYIDKDIKN-----MHSYTCRNYRN	898
gf:1198542314 gb NFI01000006.1	-----PKKKVKDYLAKNFEY-----SNKWSLKEFRN	864
emb CGNF01009141.1	-----INPRASAYLTVNLAN-----SDEFATIRFRN	801
Ga0129306_1000735	-----LVRTLRRREKRVCDNLHNNYAYLCGAD-----AEPGDASLILLFVYRN	872
MH0288_GL0082219	-----LNPRAAKYLQINLEH-----SDNTLIRQYRN	884
FIG-022__GL0026351	-----LNFHASRYLKVNISN-----ADEWSIVKFRN	779
FIG-028__GL0185479	-----LNSHARNYLIRVNI SN-----SDEWSIVKFRN	781
250twins_35838_GL0110300	REFAQIKRRESGCYFKSYHYDYLSKNSNEFKQ-----NHIDFAVTSYRN	893
FIG-014__GL0226364	QHRENDKYDSCMHFSRHSFEYISKNFAEVQSH-----PDPLKLFIKYRN	857
250twins_36050_GL0158985	KVLRQNDKMIKRMHFTPHSLNVVQKNLESVQS-----NGLAAAVIKEYRN	916

FIG. 18DDD

emb!OJMM01002900.1!	LAYRANDKVVXHTHENLHSHYKYVKNHYEEISK.....ASKHITAYRN	887
emb!OGPN01002610.1!	PLRKANDKLLKQHTYHNHSYIYVNNLNSFTDIDYCAKDVGLPAPFNKNDNASLLGEMRN	930
emb!OHCPO1000044.1!	ELRKENDQLLKKTHYSKHAICYIVDNNLITGAVANDNGRGLPCLSEKNDNANLFLMRN	916
PIG--046_GL0077813	AMRKVNDQLLKQTHFSKHSYCYIVDNNRLTGGECDTRKRVLPVINEKNDNAGILSDFRK	918
pig...chimera	AMRKVNDQLLKRTHFSKHSYCYIVDNNRLTGGECDTRKRVLPVINEKNDNAGILLDFRK	922
emb!O1EN01002196.1!	IKKNVLKYNQNEFSLNHPGEYICKNMEGYHEN.....TVRNFNR	829
O2_UC29-0_GL0096317	IKENIKKYGDFPSIDHIPGEYIKKNLECYCAG.....SINTFRN	823
CasR_Anaerobic_digLPVRKAEILRANLKHLDCE-----GFKYFRN	838
emb!OGDF01008514.1!NPKKFTVKSIIKEQYAFITPY.....IFTTYRN	822
emb!OGZC01000639.1	TVYEKAEAAKRNRLKNVKNCKTRENLENA.....DKNAIVAFRN	869
Ga0224415_10048792...chimera	LNRERVKNARNALKNIKNYIMIRLQIRDQDTS-----SGYLCGEFRD	854
emb!OCVV012889144.1!	INREKAKTAMNAHLRNTKWDIMYGQLKDLADS.....KSRICSEFRN	856
Ga0187910_10006931	INREKAKTALNHLRNTKWNVIIRENIENT.....EKDACKQFRN	854
R...flav_XPD...	INREKAKTALNHLRNTKWNVIIREDLRLI.....DNKTCTLFRN	859
Ga0224415_10007274	INREKAKTALNHLRNTKWNVIIREDLRR.....DNKACKIFRN	838
Ga0187910_10069260YLKRRPVEYLKKNMAMVGN-----SIRHFRN	791
Ga0187910_10015336	LDESASSYLSHKLVNKHVHSYKYLKQNLDCADNT.....MINQFRN	965
ODAI...chimeraTDSKSQQKAKSRPRWCILLKEDIDKINGC.....ESLRILYRN	894
31009_GL0034153YTSKKNQYIQKNMKIYSKI.....NKNFNVYVYRN	792
DLF014_GL0011914...KKNKKYGEYTSKISTYIETNKNFICE.....QERTPERYRDTQIDMFVKYRN	871
CasR_R...albusGKLGCKNKVGRYLKNNISCCIDG.....LINTYRN	833
Ga0187910_10040531GH.....FKNKKWRGIAEQNLNNSDVP.....VIKSFNR	826
tpg!DJXD01000002.1!GHLKNGKGRKRWYVLVKNNLQMSDIT.....AVSSFRN	773
CasR_E...siraenum	DKSF--AENAANKYLRNARWYKLIIDLNLKKSERA.....VVNEFRN	850

FIG. 18EE

FIG--018_GL0023397	DLGR--AETAAERHLKRRWYVQIFNNLKQADRN-----VINEFRN	845
FIG--025_GL0099734	-----NLHRSKKWNRLMRVNLNYSAA-----ASQNRN	844
Ga0129317_1008067_chimera	-----HFISHNRRMRDCIEENVKNSQLI-----FGKEDAVIRFN	761
EYZ--362E_GL0088915	-----KGYLARNKRLCQCVRDIDNAKHL-----NGDITDYN	820
uncultured_Ro_sp	-----LELKNNERLRKCVVDINNADSS-----MFRKYRN	821
R...flav._FD-1	-----RYLAGNKRRLRDCVKQDIDNAKWF-----VSDKYSITKYRN	826
tpg DRYI01000091.1	NQDKVQMPKNYFLANKRNRWEAIEQDIDNAKWF-----IGEKENVKNYRN	850
	* ,	
037__emb OIZA01000315.1	NIVHLKIIA--KLSEL---TKNIDK-----DINSYFDIYHYCTQRYVMFDYCKKNN	886
emb OCTW011587266.1	KIAHINVIM--SAGKY---TGGIK-----RMDSYFSVFQHLVQCDDILSNPNNKG	851
k87__11092736	DAEHLNPNV--EFDVY-----IEDLRPEGTPKNKDYHIGSYFEIYHYIRQRAYLEEVLAQR	945
BMZ--11B_GL0037915	DAMHLNAMF--KIAPF---VDFRK---G-----SSHEMRSFEELFHFLEQKTLDSKEDFV	881
BMZ--11B_GL0037771	DVEHLNVLR--AIPDY-----IQDFRH---GEGK--ETAMNSYFELYHYLMQRMLKNTLDEL	872
BMZ--11B_GL0069617	QSAHLNVLTLQALPQF-----IGQFAR-----NSKIGKTARMTSYFQLYHYLLQRLLCEDLNFDV	868
Ga0099364_10024192	CALHLNVLTL--ALEEF---VGEFRK---I-----GDKMTSFEELYHFLLQKLMLEAAGLNL	984
530373_GL0023589	IIMHLNAVS--AEFQY---LEGLK-----VVHSYEDLYHYTYVMMLLHPQNEKE---	881
emb ODA1011611274.1	AAAHLTAVA--AAQFY-----VSELR-----EIHSYFEVYHYAMQRYLKSQAIFA---	936
EMG...10003641	KVAHLEVVS--NIDEY---LSGIK-----HIESYVALYHYLMQKCLKLNRYRIE---	990
CasR_F1E0_metageno	AVDHLNAIP--PLGSL---CRDIG-----RVDSYFALYHYAVQQYLNQFYRKYR---	844
emb OIZX01000427.1	AVAHLNVAIA--HLADY-----SADMR-----EITSYGLYHYLMQRHLFKRHWQI	881
160582958_gene49834	LVAHLNVS--KLQNVVSELPNDY-----QITSYFSFYHYCMQLGLMEKVSCK---	944
gi 1198542314 gb NFIR01000008.1	SVQHLNVIR--DAHKYIKYINDNK-----DVQSYFALYHYLVQRYISERAANRT	911
emb OGNF01009141.1	TAEHLEALR--NAUKY---LNDLK-----QFDSYFEIYHYITQRNIKEKCEML---	844

FIG. 18FF

Ga0129306__1000735	KAHLSVLN--KGRL-----SGDLK-----EAKSWFYVYHFLMQRVLEEEFRNI--	915
MH0288__GL0082219	SVARLNAV--NMNLY----IGDLR-----EITSYFGIYHYIMQRDLIITRCQKV--	927
PIG-022__GL0026351	GTAHLSVR--NIDTC----IEGLK-----DFKSYFEIYAYTVQKSIKKYFDAEF	823
FIG-028__GL0185479	GVAHLSVR--NIDYS----IKGLR-----EFKSYFEIYAYTVQKSIKNYDAES	825
250twins__35838__GL0110300	NVERLNVH--CMTKY---FSEVK-----DVKSYGVVYCYIMQMLCDELIIGN	937
PIG-014__GL0226364	NVMRLNIR--DLNDY----LPDFK-----G----KAFKIDTFDYDLYCYLQKMLLSEVD-----	904
250twins__36050__GL0158985	AVAHLNIN--RLDEY-----IGSAR-----ADSYSLCYCLQMYLSKNFSVG--	958
emb OJMM01002900.1	NVQHLNVMN--SITKY---LGDIS-----EVTSSYSLCYTLQRLLLDDNNND--	930
emb OGFN01002610.1	DIHNLNVH--DMVKY---IEELK-----DISSYAFYCYVLQRRLLVCKDPNC--	973
emb OHCPO1000044.1	KIVHLNVVH--DMVKY---INEIK-----NITSYAFFCYVLQRMIIIGNNSNE--	959
PIG-046__GL0077813	TIHNLNVH--KMVDY---VAEIK-----GITSYAFFCYVLQMPVGNKN-E-	960
pig__chimera	TIHNLNVH--KMVDY---VDEIK-----GITSYAFFCYVLQRMVLGNLNE--	965
emb CIEN01002196.1	MVAHLAVA--RVPLY---ISEVT-----QIDSYALYHYCMQNNILQGIQSG	873
O2_UC29-0__GL0096317	AVAHLALMA--RVSEY---ASDIK-----NVNSYFSLYHYCMQKHILYCARRFN	867
CasR_Anaerobic_dig	QVTHLNAIR--VAYKY---INEIK-----SVHSYFALYHYIMQRHLYDSLQAKA	882
emb OGDF01008514.1	MIAHLAAVT--NAYKY---IPQMD-----RFKSWFHLHYHYIYQHSLIQQYEYDR	866
emb OGZC01000639.1	IVAHLWIR--DADRF-----ITGMG-----AMKRYFDCYHYLLQRELGYILEKSN	913
Ga0224415__10048792__chimera	KVARLEVAR--HAHEY---IGNIK-----EVNSYFQLYHYIMQCRLYDVLKNNT	898
emb OCVV012889144.1	KAHLEVAR--YAHMY---INDIS-----EVKSYFRLYHYIMQRKIIDVIENNP	900
Ga0187910__10006931	KADHLEVAR--YAYKY---INDIS-----EVNSYFQLYHYIMQRIIIDSSGN--	906
R__flav__XPD__	KAVHLEVAR--YVHAY---INDIA-----EVNSYFQLYHYIMQRIIMNERYE-----	901
Ga0224415__10007274	KVAHLEAIR--YAHLY---INDIA-----EVNSYFQFYHYIMQRRIIMAERYD--	880
Ga0187911__100069260	ATVHLNVIM--EAHRY---TKDIK-----YIGSYALYHYIYQHRHLLDKIEEDS	835
Ga0187910__10015336	NVARLNTIK--NMDG-----IENVT-----GITSYFQIYHYLMQKALYKEFKKCR	1008

FIG. 18GGG

ODAL_chimera	AVAHLNIPR--NAIDY-----ICGFK-----KADSYFAIYHYNMQQYLSKKHGLP--	937
31009_GL0034153	NIAHLNSIK--NATKY----VNDIG-----FTISYFQLYHYLMQCDLKDWWKKEY	836
DLF014_GL0011914...	SHAHNAVR--KASKY----INDIR-----GFDTYFELYHYIMQDLDKILEKIA	915
CasR_R...albus	QVDHEAVVR--KIGNY-----AAYIK-----SIGSWFELYHYIQRIVFDEYRFAL	877
Ga0187910...10040531	MAAHISVIR--NIDLY----IGDIQ-----KVDSYFALYHFLMOKLIIQRVVPENT	870
tpgIDJXD01000002.1	IVAHISVVR--NSNEY----ISGIG-----ELHSYFELYHYLVQSMIAKNWYDT	817
CasR_E...siraeum	TVCHLNAIR--NININ-----IYEIK-----EVENYFALYHYLIQKHLENRFADKK	894
PIG-018_GL0023397	SVCHLNAIR--NINTN----IEGIG-----FVKNYFALYHYLIQRHISKRIS---	887
PIG-025_GL0099734	AVAHLNPIR--NADVL----IEGIE-----NITSYFDIYHYIVQRSVLNRINKT--	887
Ga0129317...1008067_chimera	NVAHLSAIR--NANEY----IGDIR-----EITSYFALYHYLMQRKLIDDCKVN--	804
EYZ-362E_GL0088915	CVAHLNVR--NCDQY----LENID-----TVHSYFSLYQYLVQHYLLKRFEXDK	864
uncultured_Ru_sp	CIAHLTVR--ELKEY----IGDIC-----IVDSYFSIYHYVMQRCITKRENDTK	865
R...flav...FD-1	NVAHLTAVR--NCAEF----IGDIT-----KIDSYFALYHYLIQRQLAKGLDHER	870
tpgIDBYI01000091.1	NVAHLTAIR--NCAEF----IGEIT-----KIDSYFALYHYLIQRQLAGRLDPNH	894
	* :	
	* :	
037..._emb OIZA01000315.1	NVVLVLA-----KMKDLAHIKSDCDEFSSKHTYPYSSAVLRFM	922
emb OCTW011587266.1	KCFES-----ESL---KPLL---DMKFDGDEKLYSKRLTRAL	884
k67...11092736	KEYRD-----SGSFTDEQLDK--LQKILDDIPRARGSYDKNLLKLE	983
BMZ-11B_GL0037915	QGMRE-----S-----YGDRIRVEPCVDLIKYL	904
BMZ-11B_GL0037771	S--HWS-----G-----WIMFSGRPDRDLIQIA	893
BMZ-11B_GL0069617	PEEWK-----N-----KIQETGLPCNDLIHIC	890
Ga0099364...10024192	D--EYR-----E-----RIDTYQIACKDLINI	1005
530373...GL0023589	-----KRRNEKEKKQNEKENNMILSIA--ALEKYDVMVKYQSANKDFIYAI	925

FIG. 18HHH

emb ODAI011611274.1ELVSKNGPASGK--IAAWANAVDRCHSFCKDWLWLL	970
EMG_10003641	-----DHSQNEYKN--LNDFSSKLDKKGTYVKDFVKAL	1021
CasR_P1E0_metagenc	-----TPR--EQELFAAMAQHRITWCSDDLVKAL	869
emb OIZX01000427.1	-----TEEEOKLIEQEQQLAW--EKALFDKILQYHSYNKDLVKAL	926
160582958...gene49834	-----NIPLVESLKNEANDAQSISAKK--TLEYFDLIEKNRTYCKDFLAKAL	988
gi 1198542314 gb NFIR01000006.1	-----DKESLSEGR--LOYYSLQVKEYRNTYCKDFVKAL	942
emb CGNF01009141.1	-----KEQ--TVKYNNDLKXHGYSKDFVKAL	869
Ga0129306_1000735	-----QALPER--LRELLMBAERYRGCSKDLIKVL	943
MH0288_GL0082219	-----KMNPV--VVGYFDKVPYRTCTCKDLVKAL	955
FIG-022_GL0026351	-----SEDRNLEDGKISLCNGVYIHERQKNGKFKIYVKINPK--TIEYMESVDKYGSYTKDFVKAL	882
FIG-028_GL0185479	-----VQINPK--TIEYMESVDKYGSYTKDFVKAL	854
250twins_35838_GL0110300	-----VRQI--PEEYNRLKDHGTYSKNLMWLL	968
FIG-014_GL0226364	-----G--FSAINENIDKYCTASKDFMNL	927
250twins_36050_GL0158985	-----Y--LINVQKQLEEHHTYMKDLMWLL	981
emb OJMM01002900.1	-----K--FASIKGNLRKPCIYNKDFMNL	953
emb OGFN01002610.1	-----QNKF--KAKYAKELNDYGTYNKNLMMML	999
emb OHCPO1000044.1	-----QNKF--KAKYSKTLQEFFGTYSKDLMWVL	985
FIG-046_GL0077813	-----KNAL--KEYSEYAKSYCTYSKDFMWLI	986
pig_chimera	-----KNAI--KEYSATYKSFGTYSKDFMWLI	991
emb OIEH01002196.1	-----DNIK--LKNALENARVHRHTYSKDAVKYL	902
O2_UC29--0_GL0096317	-----GNEK--MAAALNDABJHRSYSKDAVKFL	895
CasR_Anaerobic_dig	-----SGF--VIDALKKSEHKKIYSKDLLHVL	910
emb OGDF01008514.1	-----GAPVYSER--VLQLLEQCREHSNYSRDLHL	901
emb OGZC01000639.1	-----GSEY--TKKSLKQVQYHSYCKDFLHML	940

FIG. 18H

Ga0224415_10048792_chimera	KAEAMVKGN--AKEYFEALEKEGTYNDKLLKIA	929
emb10CVV012889144.1	K--AKYEGK--VKVYFEDVKNNKKYKNLLKLM	929
Ga0187910_10006931	---NANGM--IKKYYESVISDKKYNDRLKLL	933
R_flav_XFD_	---KSSGK--VSEYFDVNDKKKYNDRLKLL	928
Ga0224415_10007274	---KSSGK--VREYFDVNNKYYNDRLKLL	907
Ga0187911_10069260	YAEKTVS--EKLWESQISQYGYTSKDFVKAL	864
Ga0187910_10015336	ENAVRWIPIYITENAEFKYVWNKKEQQEVEVSNFKIFGYMENIKNHSNTYCKDFVKAL	1068
ODAI_chimeraEGLGCTHGVYCKDLVKAL	955
31009_GL0034153	EKKISDKY--YNLEKYKNNNIYKDFVKAI	867
DLF014_GL0011914_	EGAAKETEEGQKQDEL--LKITFDNLDPKYGYTVKDFVKAL	953
CasR_R_albus	N--NT--ESNYKNSIKHHTYCKDMVKAL	902
Ga0187910_10040531	KGLSDQ--TKKYYDALEQYNTYCKDFVKAY	898
tpg1DJXD01000002.1	S-HQPK--TAEYLNLLKHHHTYCKDFVKAY	844
CasR_E_siraeum	VERD--TGDFISKLEEHKTYCKDFVKAY	920
PIG-018_GL0023397	FNSF--SAEYISNLEKYHTPSEYFIKAY	913
PIG-025_GL0099734PK--MKEYEELINKYHIYKDFVKAL	911
Ga0129317_10068067_chimeraDT--AHKYFEQLTKYKIYVMDMVKAL	828
EYZ-362E_GL0088915	YRCKDDL--AIHYYSQVKQYRPSYVKDLVKAL	893
uncultured_Ru_sp	QEEKIKYEDDLLKNHGYTKDFVKAL	890
R_flav_FD-1	SGFDR--NYPQYAPLEPKWHIYVKDVVKAL	897
tpg1DBYI01000091.1	PGFEK--NYPQYAPLEPKWNTYVKDMVKAL	921
037_...emb OIZA01000315.1	NLPFAYNVPRFKNLSYKTFDKQ	945

FIG. 18J

emb OCTW011587266.1.1	NIPFGYVPRYKNLTFEKIYILKSSINE.....	911
k87...1.1092736	YLFFAYNLPRYKNLTFTEALFDDDDSVGKKRVAEWREKREKTREAEERQRRQR-----	1034
BMZ-11B_GL0037915	YITLGSLSLPRYKINTVSGLFDPDSEDAKERAKGKRA-----	940
BMZ-11B_GL0037771	FVSLAYNLPRYRNLTKEHFDDIVLOKIREKESLD-----	928
BMZ-11B_GL0069617	FVPLGYNLPRYKNLTFIECLFDEDSVSAKEGKKA-----	923
Ga0099364_10024192	YVSLGYNLPRYKNLTCEPLFDEESATGKERQTRLDEKSKEKKQKGGQK.....	1054
530373_GLU0023589	NSPLAYNAARYINLSCRTFEEGFGK.....	951
emb ODAI011611274.1.1	NVFFAYNPARYKNLSIANLFDKNEAPVTEDESEQKEDF-----	1009
EMG_10003641	NVFFGYNLPRYKNLSIDELFDRENKLTGGTIEKKE-----	1057
CasR_P1E0_metageno	NTPFGYNLARYKNLSIDGLFDREGDHHVVREDGEPAAE-----	906
emb OIZX01000427.1.1	NAPFGYNLARYKNLSIEPLFSKEAAPAAEIKATHA-----	961
160582958_gene49834	NAPFSYNLPRFKNLSIEALFDKNIVYEQADLKE-----	1022
gi 1198542314 gb NFIR01000008.1.1	NVFFAYNLPRYKNLSIDELFDNNYLPNKAKKWLPEKKEGEYVMECCNKDAGQVANA-----	1001
emb OGNF01009141.1	CVFFGYNLPRFKNLSIDALEFDKNDKREKLKKGCFD-----	904
Ga0129306_1000735	NLTFAYNLPRYKNLSIDGRFDKNHPDPSDE-----	973
MH0288_GL0082219	NITFFGYNLPRYKNLSIEGLEFDKNRPGDKKESALLAE-----	991
FIG-022_GL0026351	NTPFGYCLARYKNLSIEPLFDPRNIVYNKEDVIEEFVSNKSEQ.....	924
FIG-028_GL0185479	NTPFGYCLARYKNLSIEPLFDPRNTVYNKKDVKEEFVAVNSK.....	894
250twins_35838_GL0110300	NFPFAYNLARYKNLSNEDLFNKAKNNDQKSK-----	998
FIG-014_GL0226364	NIIFAYNVPRYKNLSNKELFYRLDSEQ.....	954
250twins_36050_GL0158985	NIPFAYNLARYKNLSNEKLFYDEEAAAEEKADKAENERGE-----	1020
emb OJMM01002900.1.1	NIPFAYNLPRYKNLSNEEIFYDELQK-----	979
emb OGPN01002610.1.1	NLFPFAYNLPRYKNLSSEFLFYDMEYNNKKDDE-----	1030
emb OHCPO1000044.1.1	NLFPFAYNLPRYKNLSNEQLFYDEEERMEKIVGRKNDUSR-----	1023

FIG. 18KK

PIG-046_GL0077813	NLPFAYNLPRYKNLSNEQLFYDEERNETEGQIDRL-----	1022
pig_chimera	NLPFAYNLPRYKNLSNEQLFYDEERNETEEQIDRL-----	1027
emb102196.1	CLPFAYNISRYKALTIKIDLEFDWTEYSCKKDE-----	933
O2_UC29--0_GL0096317	CLPFAYNLPRYKALITVEGLEFDWNEYGODDD-----	925
CasR_Anaerobic_dig	HSPFGYNTARYKNLSIEALFDKNEISRPVNFSLTND-----	946
emb1008514.1	NLPFGYNLPRYLNLSSEKYPDANAI-----	926
emb1000639.1	CLPFAYCIPRYKNLSIAELFDREHEPEAEFKEEASVNSQFIIT-----	984
Ga0224415_10048792_chimera	CVPFYGCIPRYKNLSMEELEFDNNEKKFKKKKAPENT-----	965
emb10289144.1	CVPFYGCIPRFKNLSIEQMFDMNETDNSDKKKEK-----	963
Ga0187910_100006931	CVPFYGCIPRFKNLSIEALFDKNEAKYDKIKKKVAVR-----	971
R_flav_XPD	CVPFYGCIPRFKNLSIEALFDREARAKFDKKEKKVSGNS-----	967
Ga0224415_10007274	CVPFYGCIPRFKNLSIEALFDNNEAVKFDKKEK-----	940
Ga0187911_10069260	CCPFYXNLPRFKNLSIEQLEFDRENSKEITDAIAPRQ-----	900
Ga0187910_10015336	CAPFAYNLPRFKNLSIEELEFDHHELSEEPKESMKLTD-----	1105
ODAL_chimera	NAPFGYNLPRFKNLSVEALFDNNEEPAPEGKERSKNADGCN-----	996
31009_GL0034153	NVPFAYNYSRFKNLSIDKLEFDKNDPRKKKWIVTFNF-----	903
DLF014_GL0011914	NIPFAYNRYRYKNLSIDELEFDKNNTKTIKIQDKKLEERT-----	993
CasR_R_albus	NTFFGYDLPRYKNLSIGDLEFDNNYLNKTKESIDANSSIDSQ-----	944
Ga0187910_10040531	CIPFAYVTPRYKNLTIIDGLEFDKNRPEGDK-----	927
tpg1DJXD01000002.1	CIPFGYVVERKYKNLTIINELEFDNNPNPEPKKEEV-----	877
CasR_F_siraeum	CIPFGYNLRYRYKNLTIIDGLEFDKNYPGKDDSDDEQK-----	954
PIG-018_GL0023397	CIPFAYNLVRYKNLTIIDGLEFDKRNAPSEK-----	941
PIG-025_GL0099734	CAPFGYVNVVRFKALTIYENFDNRYPEEKRAEEAKFSC-----	948
Ga0129317_1008067_chimera	CSPFGYNLPRFKNLSIEGKFPDMHESK-----	854

FIG. 18LU

EYZ...362E...GL0088915	NAPFGYNIPRFKNLITVEQLFDRNADPEITIA.....	924
uncultured_Ko...sp	NSPFGYNIPRFKNLSIEQLFDRNEYLTEK-----	919
R...flav...FD-1	NAPFGYNIPRFKNLSIDALFDRNEIKKNDGEKKSDD--	933
tpg:DEYI01000091.1:	NSFFGYNIPRFKLSIDALFDRNEMKEETDDEKKIQI.....	958

:. * * : : :

FIG. 18MMIM

RNA TARGETING METHODS AND COMPOSITIONS**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a continuation of U.S. application Ser. No. 16/257,493, filed Jan. 25, 2019, which is a divisional of U.S. application Ser. No. 15/937,699, filed Mar. 27, 2018, now U.S. Pat. No. 10,476,825, which claims priority to U.S. Provisional Application No. 62/548,846 filed Aug. 22, 2017, U.S. Provisional Application No. 62/572,963 filed Oct. 16, 2017, and U.S. Provisional Application No. 62/639,178, filed Mar. 6, 2018, all of which are herein incorporated by reference in their entireties.

ACKNOWLEDGMENT OF GOVERNMENT SUPPORT

This invention was made with government support under 5 DP5 OD021369-02 and 5 R21 AG056811-02 awarded by The National Institutes of Health. The government has certain rights in the invention.

FIELD

This disclosure relates to a CRISPR/Cas system for modifying (including detecting) RNA, which utilizes novel Cas13d proteins (also referred to as CasR and nCas1) and guide RNAs.

BACKGROUND

Mapping of transcriptome changes in cellular function and disease has been transformed by technological advances over the last two decades, from microarrays (Schena et al., 1995) to next-generation sequencing and single cell studies (Shendure et al., 2017). However, interrogating the function of individual transcript dynamics and establishing causal linkages between observed transcriptional changes and cellular phenotype requires the ability to actively control or modulate desired transcripts.

DNA engineering technologies such as CRISPR-Cas9 (Doudna and Charpentier, 2014; Hsu et al., 2014) enable researchers to dissect the function of specific genetic elements or correct disease-causing mutations. However, simple and scalable tools to study and manipulate RNA lag significantly behind their DNA counterparts. Existing RNA interference technologies, which enable cleavage or inhibition of desired transcripts, have significant off-target effects and remain challenging engineering targets due to their key role in endogenous processes (Birmingham et al., 2006; Jackson et al., 2003). As a result, methods for studying the functional role of RNAs directly have remained limited.

One of the key restrictions in RNA engineering has been the lack of RNA-binding domains that can be easily retargeted and introduced into target cells. The MS2 RNA-binding domain, for example, recognizes an invariant 21-nucleotide (nt) RNA sequence (Peabody, 1993), therefore requiring genomic modification to tag a desired transcript. Pumilio homology domains possess modular repeats with each protein module recognizing a separate RNA base, but they can only be targeted to short 8 nt RNA sequences (Cheong and Hall, 2006). While previously characterized type II (Batra et al., 2017; O'Connell et al., 2014) and VI (Abudayyeh et al., 2016; East-Seletsky et al., 2016) CRISPR-Cas systems can be reprogrammed to recognize

20-30 nt RNAs, their large size (~1200 amino acids, aa) makes it difficult to package into AAV for primary cell and in vivo delivery.

SUMMARY

This application provides bioinformatic analysis of prokaryotic genomes to identify sequence signatures of CRISPR-Cas repeat arrays and mine previously uncharacterized, compact Cas ribonucleases that can be used for RNA targeting tools. Engineered Type VI-D CRISPR effectors can be used to efficiently knockdown endogenous RNAs in human cells and manipulate alternative splicing, paving the way for RNA targeting applications and further effector domain fusions as part of a transcriptome engineering toolbox.

Provided herein are methods of modifying one or more target RNA molecules, such as a clustered regularly interspaced short palindromic repeats (CRISPR)-CRISPR associated (Cas) system-mediated RNA editing method. Such methods can include contacting one or more target RNA molecules with a non-naturally occurring (e.g., does not naturally occur in the cell or system into which it is introduced) or engineered CRISPR-Cas system. Such a CRISPR-Cas system can include (1) at least one Cas13d protein or at least one Cas13d nucleic acid coding sequence (such as a mRNA or a vector encoding the at least one Cas13d protein) and (2) at least one CRISPR-Cas system guide nucleic acid molecule (such as a guide RNA, gRNA) that hybridizes with the one or more target RNA molecules, or at least one nucleic acid molecule encoding the gRNA. The Cas13d protein forms a complex with the gRNA, and the gRNA directs the complex to the one or more target RNA molecules and modifies (e.g., cuts, detects) the one or more target RNA molecules. In some examples, the one or more target RNA molecules (or a cell containing the one or more target RNA molecules) are contacted with a complex including the at least one Cas13d protein and the at least one gRNA. In some examples, the system includes Mg²⁺. However, in some example, the system does not include Mg²⁺, such as if cleavage of the target RNA is not desired.

In some examples, contacting the one or more target RNA molecules with the non-naturally occurring or engineered CRISPR-Cas system includes introducing into a cell (such as a eukaryotic or prokaryotic cell) containing the one or more target RNA molecules the non-naturally occurring or engineered CRISPR-Cas system, for example using endocytosis, a liposome, a particle, an exosome, a microvesicle, a gene gun, electroporation, a virus, or combinations thereof. In some examples, contacting the one or more target RNA molecules with the non-naturally occurring or engineered CRISPR-Cas system includes contacting a cell-free system (such as a biological or environmental sample, or a cell lysate) containing the one or more target RNA molecules with the non-naturally occurring or engineered CRISPR-Cas system (for example in a diagnostic method to detect a target RNA).

In some examples, the least one Cas13d protein includes one or more HEPN domains, is no more than 150 kD, no more than 140 kD, no more than 130 kD, no more than 120 kD, such as about 90 to 120 kD, about 100 to 120 kD or about 110 kD; includes one or more mutated HEPN domains, and can process the guide RNA, but cannot cleave or cut the one or more target RNA molecules, includes an Cas13d ortholog from a prokaryotic genome or metagenome, gut metagenome, an activated sludge metagenome, an anaerobic digester metagenome, a chicken gut metag-

enome, a human gut metagenome, a pig gut metagenome, a bovine gut metagenome, a sheep gut metagenome, a goat gut metagenome, a capybara gut metagenome, a primate gut metagenome, a termite gut metagenome, a fecal metagenome, a genome from the Order Clostridiales, or the Family Ruminococcaceae; includes an Cas13d ortholog from *Ruminococcus albus*, *Eubacterium siraeum*, a *flavefaciens* strain XPD3002, *Ruminococcus flavefaciens* FD-1, uncultured *Eubacterium* sp TS28-c4095, uncultured *Ruminococcus* sp., *Ruminococcus bicirculans*, or *Ruminococcus* sp CAG57; includes at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253, or combinations thereof. In some examples, the least one Cas13d protein has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253, and includes at least one motif shown in SEQ ID NO: 195, 196 or 197. In some examples, the least one Cas13d protein further includes one or more other agents (e.g., is a fusion protein), such as one or more subcellular localization signals, one or more effector domains, or combinations thereof. In some examples, the least one Cas13d protein that includes one or more HEPN domains, is no more than 1500 aa, no more than 1200 aa, no more than 1100 aa, no more than 1000 aa, such as about 800 to 1500 aa, about 800 to 1250 aa or about 850 to 950 aa.

Also provided are isolated nucleic acid molecules encoding such Cas13d proteins, such as a cDNA, genomic DNA, RNA, or mRNA. Such isolated nucleic acid molecules can be part of a vector (such as a plasmid or viral vector), and can be operably linked to a promoter. In some examples, an isolated nucleic acid molecule encoding a Cas13d protein has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 124, 125, 126, 127, 128, 139, 140 or 141. In some examples, an isolated nucleic acid molecule encoding at least one Cas13d protein (which can be part of a vector) includes at least one Cas13d protein coding sequence codon optimized for expression in a eukaryotic cell, such as human cell, for example a Cas13d coding sequence having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%

sequence identity to SEQ ID NO: 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 142, 143, 144, or 145.

In some examples, the gRNA that hybridizes with the one or more target RNA molecules in an Cas13d-mediated manner includes one or more direct repeat (DR) sequences, one or more spacer sequences, such as one or more sequences comprising an array of DR-spacer-DR-spacer. In some examples, the one or more DR sequences have at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 129, 130, 131, 132, 133, 134, 135, 136, 137, 148, 150, 151, 152, 154, 156, 157, 159, 161, 163, 165, 167, 169, 176, 178, 180, 182, 184, 186, 188, 190, 191, 192, 193, 199, 201, 203, 205, 207, 209, 211, 213, 215, 217, 219, 221, 223, 225, 227, 228, 230, 232, 234, 236, 238, 240, 242, 244, 246, 248, 250, 252, or 254. In one example, the gRNA includes additional sequences, such as an aptamer sequence.

In some examples, a plurality of gRNAs are generated from a single array, wherein each gRNA can be different, for example target different RNAs or target multiple regions of a single RNA, or combinations thereof.

Methods of targeting one or more target RNA molecules are provided. In some examples, an entire RNA is targeted. In some examples, a portion of an RNA is targeted. Targeting an RNA molecule can include one or more of cutting or nicking one or more target RNA molecules, activating one or more target RNA molecules, deactivating the one or more target RNA molecules, visualizing or detecting the one or more target RNA molecules, labeling the one or more target RNA molecules, binding the one or more target RNA molecules, editing the one or more target RNA molecules, trafficking the one or more target RNA molecules, and masking the one or more target RNA molecules. In some example, modifying one or more target RNA molecules includes one or more of an RNA base substitution, an RNA base deletion, an RNA base insertion, a break in the target RNA, methylating RNA, and demethylating RNA.

In some examples, such methods are used to treat a disease, such as a disease in a human. In such examples, the one or more target RNA molecules is associated with the disease.

Also provided are isolated proteins, including non-naturally occurring proteins. In some examples, a protein has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, or 113. In some examples, an isolated protein is an Cas13d ortholog from a prokaryotic genome or metagenome, gut metagenome, an activated sludge metagenome, an anaerobic digester metagenome, a chicken gut metagenome, a human gut metagenome, a pig gut metagenome, a bovine gut metagenome, a sheep gut metagenome, a goat gut metagenome, a capybara gut metagenome, a primate gut metagenome, a termite gut metagenome, a fecal metagenome, a genome from the Order Clostridiales, or the Family Ruminococcaceae. In some examples, an Cas13d ortholog includes an Cas13d ortholog from *Ruminococcus albus*, *Eubacterium siraeum*, a *Ruminococcus flavefaciens* strain XPD3002, *Ruminococcus flavefaciens* FD-1, uncultured

Eubacterium sp TS28-c4095, uncultured *Ruminococcus* sp., *Ruminococcus bicirculans*, or *Ruminococcus* sp CAG57. The protein is an Cas13d protein that further includes one or more other agents or domains (e.g., is a fusion protein), such as one or more subcellular localization signals, one or more effector domains, or combinations thereof.

Also provided are isolated guide RNA (gRNA) molecules. In some examples, an isolated gRNA includes one or more direct repeat (DR) sequences, such as an unprocessed (e.g., about 36 nt) or processed DR (e.g., about 30 nt). In some examples a DR has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 129, 130, 131, 132, 133, 134, 135, 136, 137, 148, 150, 151, 152, 154, 156, 157, 159, 161, 163, 165, 167, 169, 176, 178, 180, 182, 184, 186, 188, 190, 191, 192, 193, 199, 201, 203, 205, 207, 209, 211, 213, 215, 217, 219, 221, 223, 225, 227, 228, 230, 232, 234, 236, 238, 240, 242, 244, 246, 248, 250, 252, or 254. Such a gRNA can further include one or more spacer sequences specific for (e.g., is complementary to) the target RNA.

Also provided are ribonucleoprotein (RNP) complexes, which include an Cas13d protein provided herein and a gRNA provided herein.

Also provided are recombinant cells that include any Cas13d protein (or nucleic acid molecule encoding Cas13d), any gRNA, any RNP complex, or any vector, provided herein. In one example, the cell is not a bacterial cell. In one example, the cell is a bacterial cell.

Also provided are compositions that include one or more of any Cas13d protein (or nucleic acid molecule encoding Cas13d), any gRNA, any RNP complex, any isolated nucleic acid molecule, any vector, or any cell, provided herein. Such compositions can include a pharmaceutically acceptable carrier.

Also provided are kits. Such kits can include one or more of any Cas13d protein (or nucleic acid molecule encoding Cas13d), any gRNA, any RNP complex, any isolated nucleic acid molecule, any vector, any cell, or any composition provided herein. Such reagents can be combined or in separate containers.

The foregoing and other objects and features of the disclosure will become more apparent from the following detailed description, which proceeds with reference to the accompanying figures.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B: Bioinformatic pipeline for the identification of the RNA-targeting class 2 CRISPR system Cas13d. (A) Schematic describing a computational pipeline for CRISPR system identification. A minimal definition for a putative class 2 CRISPR locus was used, requiring only a CRISPR repeat array and a nearby protein >750 aa in length. The initial search was performed on prokaryotic genome assemblies derived from NCBI Genome, and later expanded via TBLASTN of predicted Cas13d proteins against public metagenome sequences without predicted open reading frames. DR, direct repeat. (B) Phylogenetic classification and alignment of full-length Cas13d effectors and metagenomic fragments. Cas13d effectors and metagenomic Cas13d protein fragments cluster into several distinct branches, which are colored for ease of interpretation. Shading indicates residue conservation using the Blosom62 matrix. Full-length Cas13d effectors used in this study were sampled from distinct branches of the Cas13d family. Alignment of Cas13d proteins and protein fragments was per-

formed using ClustalOmega 1.2.4 and maximum-likelihood tree building was performed with PhyML 3.2.

FIGS. 2A-2C: Type VI CRISPR-Cas13d is a family of single effector CRISPR ribonucleases. (A) Maximum-likelihood phylogenetic tree of Cas13d effectors used herein, with the full Cas13d CRISPR locus depicted along with conserved HEPN RNase domains. Grey rectangles denote CRISPR direct repeats (DRs) and blue diamonds indicate spacer sequences. (B) RNA sequencing of a heterologously expressed Cas13d locus from an uncultured *Ruminococcus* sp. sample. Mature gRNAs mapping to the CRISPR array indicate a processed 30 nt DR and a variable spacer length from 14-26 nt (SEQ ID NO: 278). Co-fold analysis of direct repeat truncation indicates a strong hairpin structure. (C) Purified *E. siraeum* Cas13d and catalytically dead Cas13d (dCas13d) protein are each sufficient to process a guide array into its two component gRNAs. Addition of EDTA does not impair gRNA processing. 'd', dCas13d(R295A, H300A, R849A, H854A).

FIGS. 3A-3D: Phylogenetic classification of RNA-targeting class 2 CRISPR effectors and sequence conservation within the Cas13d family. (A) HEPN motif conservation in Cas13d effectors used herein with conserved residues shaded according to Blosom62. Consensus sequence of HEPN1 domain region (SEQ ID NO: 279) and alignment of HEPN1 domain region of seven Cas13d proteins (SEQ ID NOS: 280 to 286). Consensus sequence of HEPN2 domain region (SEQ ID NO: 287) and alignment of HEPN2 domain region of seven Cas13d proteins (SEQ ID NOS: 288 to 294). The RxxxxH HEPN motif is highlighted. (B) Maximum-likelihood tree of type VI CRISPR-Cas families. Average amino acid lengths of Type VI Cas13 superfamily effectors are indicated in red. Alignment of previously described class 2 CRISPR RNA-targeting proteins (Abudayyeh et al., 2017; Cox et al., 2017; East-Seletsky et al., 2017; East-Seletsky et al., 2016; Smargon et al., 2017) and Cas13d effectors was performed using MAFFT 7.38 and maximum-likelihood tree building was performed with PhyML 3.2. Branch labels and scale bar indicate substitutions per site. (C) Predicted Cas13d direct repeat RNA secondary structure. (D) Sequence logo of full length 36 nt Cas13d direct repeats.

FIGS. 4A-4C: Purification of recombinant Cas13d protein. EsCas13d was expressed as N-terminal His-MBP fusion and purified by successive affinity, cation exchange, and size exclusion chromatography. The His-tag was removed by TEV protease cleavage. (A) Chromatogram from Superdex 200 column for EsCas13d. (B) SDS-PAGE gel of size exclusion chromatography fractions for *E. siraeum* Cas13d. (C) SDS-PAGE gel of purified *E. siraeum* Cas13d and dCas13d (R295A, H300A, R849A, H854A mutations of predicted catalytic residues in both HEPN motifs).

FIGS. 5A-5D: Programmable RNA targeting by Cas13d in vitro. (A) *E. siraeum* Cas13d requires a matching guide array or mature gRNA to efficiently cleave complementary ssRNA targets. Denaturing gel depicts cleavage reactions incubated at 37° C. for 1 hour. NT, non-targeting. (B) Substitution with dCas13d or addition of EDTA abrogate Cas13d-mediated RNA targeting with both the guide and array. 'd', dead Cas13d. (C) Denaturing gel depicting guide-target match dependent activation of Cas13d cleavage activity. Scrambled target RNA ('A') is fluorescently labeled, while guide-complementary activator target RNA ('B') is unlabeled. RNA cleavage activity is abolished by the individual removal of guide RNA or complementary target RNA, as well as the addition of EDTA or the catalytic inactivation of Cas13d (indicated as 'd'). (D) A model for

guide and target-dependent activation of Cas13d RNase activity. The ternary Cas13d:gRNA:target RNA complex is capable of cleaving the complementary target RNA or bystander RNAs.

FIGS. 6A-6H. In vitro characterization of Cas13d properties. (A) Schematic showing the length and sequence of gRNA spacer truncations and spacer position relative to the complementary ssRNA target (from top to bottom, SEQ ID NOS: 295 to 309). (B) Denaturing gel depicting EsCas13d cleavage activity of target RNA with different spacer lengths. (C) Denaturing gel depicting EsCas13d cleavage reactions paired with 12 guides from FIG. 3A tiling a complementary ssDNA version of the ssRNA target. (D) Denaturing gel depicting cleavage reactions using EsCas13d paired with the same 12 guides tiling a dsDNA version of the complementary target. (E) Quantification of cleavage efficiency from FIG. 3A. Each PFS base is the average of 3 different spacer sequences tiling a complementary target RNA. Cleavage percentage is determined by the ratio of cleaved band intensity divided by total lane intensity. Mean is depicted \pm SD with each data point representing an independent replicate. (F) Cas13d-mediated cleavage of target RNA carrying different PFS bases given an invariant spacer sequence. Quantification of Cas13d cleavage efficiency and a representative denaturing gel depicting EsCas13d cleavage activity are shown. Differences are not significant (one-way ANOVA, $P=0.768$). Cleavage percentage is determined as above, and mean is depicted \pm SD with $n=3$. (G) and (H) Optimal temperature range for Cas13d activity. Denaturing gels depicting EsCas13d cleavage activity at temperatures ranging from 16-62° C. for two different target RNAs.

FIGS. 7A-7B: Characterization of Cas13d target substrate preference. (A) Cas13d can be generalizably reprogrammed with multiple guides and does not exhibit a protospacer flanking sequence (PFS) requirement. RNA cleavage by EsCas13d and 12 guides tiling the target RNA is shown. Control lanes are from a separate gel run in parallel. (B) Cas13d preferentially cleaves uracil bases in the loop of a hairpin or a linear homopolymer repeat, which is interrupted every 5 nt by a transition mutation (X) to enable synthesis.

FIGS. 8A-8D: RNA knockdown activity screen of engineered Cas13d orthologs in human cells. (A) Schematic for mammalian expression constructs encoding for engineered Cas13d effectors and guides. NLS, nuclear localization signal. pre-gRNA, artificial unprocessed guide RNA containing a single 30 nt spacer sequence flanked by 2 full length 36 nt DRs. gRNA, predicted mature guide RNA with a single 30 nt processed DR and 22 nt spacer sequence (B) Heatmap of mCherry protein knockdown in a Cas13d ortholog activity screen in human HEK 293FT cells using pools of 4 pre-gRNAs or gRNAs. Normalized MFI, median fluorescent intensity relative to non-targeting condition. Positions in gray were not tested, with $n=3$. (C) Immunocytochemistry of Cas13d showing localization and expression of engineered constructs. Scale bar, 10 μ m. Blue pseudocolor, DAPI staining of nuclei. (D) Comparison of Adm and Rfx Cas13d ortholog constructs for knockdown of endogenous B4GALNT1 mRNA reveals RfxCas13d-NLS (CasRx) to be most effective for both guide RNA architectures. Pools of 4 guides were used for targeting. NT, non-targeting. Values are mean \pm SEM with $n=3$.

FIGS. 9A-9H: CasRx mediates efficient and specific knockdown of diverse human coding and noncoding transcripts. (A) Multiple guide RNAs tiling a target transcript can be expressed as a single array and processed by RfxCas13d-NLS (CasRx) into individual gRNAs within the same cell. (B) Arrays of 4 guides each mediate target

knockdown by CasRx in 293FT cells via transient transfection. Knockdown relative to GFP vehicle control was determined by qPCR, with $n=3$. (C) Schematic of CasRx target sequences and spacer position-matched shRNAs. (D) Relative target RNA knockdown by individual position-matched shRNAs and CasRx gRNAs. NT, non-targeting. CRISPRi, dCas9-mediated CRISPR interference for transcriptional repression ($n=3$) (E) Volcano plot of differential transcript levels between B4GALNT1 targeting and non-targeting (NT) shRNAs as determined by RNA sequencing ($n=3$). 542 non-specific transcript changes were identified. (F) Volcano plot of differential transcript levels between B4GALNT1-targeting CasRx and non-targeting (NT) guide. Targeting guide position is matched to the shRNA shown in (E). B4GALNT1 was the only transcript exhibiting a significant change, with $n=3$. (G) Summary of significant off-target transcript perturbations by matched shRNAs and CasRx guides. (H) CasRx targeting of 11 endogenous transcripts, each with 3 guides and a non-targeting (NT) guide in 293FT cells. Transcript levels are relative to GFP vehicle control, mean \pm SEM with $n=3$.

FIGS. 10A-10D. Comparison of engineered Cas13 superfamily effectors for targeted knockdown and splicing. (A) Relative target RNA knockdown by individual position-matched gRNAs for CasRx, NLS-LwaCas13a-msfGFP (Abudayyeh et al., 2017) and PspCas13b-NES (Cox et al., 2017) in HEK 293FT cells. NT, non-targeting. Values are mean \pm SEM with $n=3$. (B) Comparison of Cas13 median knockdown efficiencies. $n=3$ per guide RNA. **** indicates $P<0.0001$ according to Friedman's test. (C) Exon exclusion by catalytically inactive NLS-dCas13a-msfGFP on the bichromatic splicing reporter. Guides are position-matched to those reported in FIG. 6B for CasRx. Values are mean \pm SEM with $n=3$. (D) Comparison of splicing modulation by NLS-dCas13a-msfGFP and CasRx. Fold change in targeted exon exclusion relative to non-targeting guide is shown. **** indicates $P<0.0001$ according to two-way ANOVA.

FIGS. 11A-11B. RNA sequencing from CasRx and shRNA targeting of ANXA4 in human cells. (A) Volcano plots of differential transcript levels between ANXA4 targeting and non-targeting (NT) shRNAs as determined by RNA sequencing ($n=3$). 915 non-specific transcript changes were identified. (B) Volcano plot of differential transcript levels for an ANXA4 targeting CasRx array used in FIG. 9B containing a guide position matched to the shRNA shown in (A) and a non-targeting (NT) array. ANXA4 was the only transcript exhibiting significant downregulation with $n=3$. HIST2HBE was the only transcript identified to exhibit significant upregulation. H2B is a dimer partner of H2AX (Du et al., 2006) which has been shown to interact with ANXA4 (Yang et al., 2010).

FIGS. 12A-12F: AAV delivery of catalytically inactive dCasRx splice effectors to manipulate alternative splicing. (A) Schematic of bichromatic exon skipping reporter. +1 and +3, reading frame. BP, intronic branch point-targeting guide. SA, splice acceptor site-overlapping guide. EX, exonic guide. SD, splice donor site-overlapping guide. AUG, start codon. UGA, stop codon. Inclusion of the second exon leads to an out-of-frame (+3), non-fluorescent translation of dsRed followed by in-frame mTagBFP2. Exclusion of the targeted exon leads to an in frame translation of dsRed (+1) followed by a stop codon. (B) Induced exon exclusion by dCasRx and an N-terminal hnRNPa1-dCasRx fusion protein targeted to pre-mRNA. The Gly-rich C-terminal domain of hnRNPa1 is used as the effector domain Exon skipping efficiency is depicted as a relative percentage of cells carrying primarily the dsRed or BFP isoform, determined through flow cytometry.

etry. NLS, nuclear localization signal. 'A', CRISPR array carrying all 4 guides. Values are mean \pm SEM with $n=3$. (C) AAV design carrying dCasRx and a three-guide array with total transgene size <4.3 kb, including AAV inverted terminal repeats (ITRs). (D) Schematic of frontotemporal dementia (FTD) disease modeling. Neurons are generated via Neurogenin-2 (Ngn2) directed differentiation of patient-derived and control iPSCs followed by transduction with dCasRx or vehicle control AAV (EFS-mTagBFP2). (E) FTD is associated with SNPs in a putative intronic splice enhancer following exon 10 of the MAPT transcript encoding for tau. Alternative splicing of MAPT exon 10 results in 4R tau (by inclusion) and 3R tau (by exclusion). SNPs in the intronic splice enhancer including the indicated IVS 10+16 mutation result in increased exon inclusion and higher levels of 4R tau. To facilitate reduction of 4R tau levels, gRNAs contained in a dCasRx array were targeted to the exon 10 splice acceptor (g1) as well as two putative exonic splice enhancers indicated in purple (g2, g3). (F) Relative 4R/3R tau transcript ratios in differentiated neurons were assayed via qPCR at 14 days following transduction with AAV. FTD, frontotemporal dementia cells carrying IVS 10+16. Values are mean \pm S.D. with $n=3$. **** indicates $P<0.0001$.

FIG. 13 is a bar graph showing RNA targeting in human cells using the disclosed methods.

FIG. 14 is a schematic drawing showing how the disclosed Cas13d and DRs can be used to achieve alternative splicing.

FIGS. 15A-15D are a series of panels showing: (A) Cas13d is converted into an active RNase complex upon binding a target matching the spacer sequence of the guide RNA. It is capable of cleaving gRNA-complementary target RNA or non-complementary bystander RNAs. (B) Cas13d target-dependent RNase activity can be converted into a detectable signal, for example through cleavage of a labeled detector RNA that is cleaved only in the presence of a target matching the spacer of the Cas13d guide RNA. In this example, the detector RNA contains a fluorophore, 'F', and a quencher 'Q', that abolishes fluorescence. Only upon bystander RNA cleavage is the fluorophore liberated from the quencher and fluorescence is generated. (C) Cas13d from *E. siraeum* produces a visible signal only in the presence of a perfectly matched target and not in the presence of different mismatched targets. (D) Cas13d from *R. flavofaciens* strain XPD3002 produces a visible signal only in the presence of a perfectly matched target and not in the presence of different mismatched targets.

FIGS. 16A-16B: (A) Alignment of 7 orthologs shows regions with high (green bars) and low (red bars) conservation. Regions selected for deletion are marked 1-10. (B) knock-down (top) and splicing (bottom) activity of full-length CasRx and CasRx deletion variants. Deletion variant 5 is shown to retain full activity, demonstrating the feasibility of deleting areas of low conservation while retaining full activity.

FIGS. 17A and 17B show targeting of *ccdB* in bacterial cells. (A) construct introduced into bacterial cells expressing *ccdB*, and (B) relative expression of *ccdB* under various conditions.

FIGS. 18A-18MMM show an alignment of 53 different Cas13d proteins as follows: SEQ ID NOs: 310 (037_-embloIOZA01000315.1), 183 (embloCTW011587266.1), 189 (k87_11092736), 220 (BMZ-11B_GL0037915), 218 (BMZ-11B_GL0037771), 222 (BMZ-11B_GL0069617), 229 (Ga0099364_10024192), 216 (530373_GL0023589), 177 (embIODAI011611274.1), 200 (EMG_10003641), 139 (CasR_P1E0_metageno), 179 (embloIXX01000427.1), 208

(160582958_gene49834), 166 (gil1_198542314|gbIN-FIR01000008.1), 185 (embIOGNF01009141.1), 202 (Ga0129306_1000735), 239 (MH0288_GL0082219), 311 (PIG-022_GL0026351), 249 (PIG-028_GL0185479), 210 (250twins_35838_GL0110300), 243 (PIG-014_GL0226364), 212 (250twins_36050_GL0158985), 175 (embIOJMM01002900.1), 164 (embIOGPN01002610.1), 160 (embIOHCP01000044.1), 312 (PIG-046_GL0077813), 313 (pig_chimera), 187 (embIOIEN01002196.1), 241 (02.UC29-0_GL0096317), 140 (CasR_Anaerobic_dig), 162 (embIOGDF01008514.1), 155 (embIOGZC01000639.1), 206 (Ga0224415_10048792_chimera), 181 (embIOCVV012889144.1), 231 (Ga0187910_10006931), 92 (R._flav_XPD_), 198 (Ga0224415_10007274), 237 (Ga0187911_10069260), 233 (Ga0187910_10015336), 253 (ODAI_chimera), 214 (31009_GL0034153), 224 (DLF014_GL0011914_), 127 (CasR_R._albus), 235 (Ga0187910_10040531), 153 (tpgIDJXD01000002.1), 125 (CasR_E._siraeum), 245 (PIG-018_GL0023397), 247 (PIG-025_GL0099734), 204 (Ga0129317_1008067_chimera), 226 (EYZ-362B_GL0088915), 3 (uncultured_Ru_sp), 126 (R._flav_FD-1), and 149 (tpgIDBYI01000091.1), from top to bottom.

SEQUENCE LISTING

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. The Sequence Listing is submitted as an ASCII text file, created on Jun. 18, 2021, 924 KB, which is incorporated by reference herein.

SEQ ID NO: 1 is an exemplary Cas13d sequence from *Eubacterium siraeum* containing a HEPN site.

SEQ ID NO: 2 is an exemplary Cas13d sequence from *Eubacterium siraeum* containing a mutated HEPN site.

SEQ ID NO: 3 is an exemplary Cas13d sequence from uncultured *Ruminococcus* sp. containing a HEPN site.

SEQ ID NO: 4 is an exemplary Cas13d sequence from uncultured *Ruminococcus* sp. containing a mutated HEPN site.

SEQ ID NO: 5 is an exemplary Cas13d sequence from Gut_metagenome_contig2791000549.

SEQ ID NO: 6 is an exemplary Cas13d sequence from Gut_metagenome_contig855000317.

SEQ ID NO: 7 is an exemplary Cas13d sequence from Gut_metagenome_contig3389000027.

SEQ ID NO: 8 is an exemplary Cas13d sequence from Gut_metagenome_contig8061000170.

SEQ ID NO: 9 is an exemplary Cas13d sequence from Gut_metagenome_contig1509000299.

SEQ ID NO: 10 is an exemplary Cas13d sequence from Gut_metagenome_contig9549000591.

SEQ ID NO: 11 is an exemplary Cas13d sequence from Gut_metagenome_contig71000500.

SEQ ID NO: 12 is an exemplary Cas13d sequence from human gut metagenome.

SEQ ID NO: 13 is an exemplary Cas13d sequence from Gut_metagenome_contig3915000357.

SEQ ID NO: 14 is an exemplary Cas13d sequence from Gut_metagenome_contig4719000173.

SEQ ID NO: 15 is an exemplary Cas13d sequence from Gut_metagenome_contig6929000468.

SEQ ID NO: 16 is an exemplary Cas13d sequence from Gut_metagenome_contig7367000486.

SEQ ID NO: 17 is an exemplary Cas13d sequence from Gut_metagenome_contig7930000403.

SEQ ID NO: 18 is an exemplary Cas13d sequence from Gut_metagenome_contig993000527.

SEQ ID NO: 19 is an exemplary Cas13d sequence from Gut_metagenome_contig6552000639.

SEQ ID NO: 20 is an exemplary Cas13d sequence from Gut_metagenome_contig11932000246.

SEQ ID NO: 21 is an exemplary Cas13d sequence from Gut_metagenome_contig12963000286.

SEQ ID NO: 22 is an exemplary Cas13d sequence from Gut_metagenome_contig2952000470.

SEQ ID NO: 23 is an exemplary Cas13d sequence from Gut_metagenome_contig451000394.

SEQ ID NO: 24 is an exemplary Cas13d sequence from *Eubacterium_siraeum_DSM_15702*.

SEQ ID NO: 25 is an exemplary Cas13d sequence from gut_metagenome_P19E0k2120140920,c369000003.

SEQ ID NO: 26 is an exemplary Cas13d sequence from Gut_metagenome_contig7593000362.

SEQ ID NO: 27 is an exemplary Cas13d sequence from Gut_metagenome_contig12619000055.

SEQ ID NO: 28 is an exemplary Cas13d sequence from Gut_metagenome_contig1405000151.

SEQ ID NO: 29 is an exemplary Cas13d sequence from Chicken_gut_metagenome_c298474.

SEQ ID NO: 30 is an exemplary Cas13d sequence from Gut_metagenome_contig1516000227.

SEQ ID NO: 31 is an exemplary Cas13d sequence from Gut_metagenome_contig1838000319.

SEQ ID NO: 32 is an exemplary Cas13d sequence from Gut_metagenome_contig13123000268.

SEQ ID NO: 33 is an exemplary Cas13d sequence from Gut_metagenome_contig5294000434.

SEQ ID NO: 34 is an exemplary Cas13d sequence from Gut_metagenome_contig6415000192.

SEQ ID NO: 35 is an exemplary Cas13d sequence from Gut_metagenome_contig6144000300.

SEQ ID NO: 36 is an exemplary Cas13d sequence from Gut_metagenome_contig9118000041.

SEQ ID NO: 37 is an exemplary Cas13d sequence from Activated_sludge_metagenome_transcript_124486.

SEQ ID NO: 38 is an exemplary Cas13d sequence from Gut_metagenome_contig1322000437.

SEQ ID NO: 39 is an exemplary Cas13d sequence from Gut_metagenome_contig4582000531.

SEQ ID NO: 40 is an exemplary Cas13d sequence from Gut_metagenome_contig9190000283.

SEQ ID NO: 41 is an exemplary Cas13d sequence from Gut_metagenome_contig1709000510.

SEQ ID NO: 42 is an exemplary Cas13d sequence from M24_(LSQX01212483_Anaerobic_digester_metagenome) with a HEPN domain

SEQ ID NO: 43 is an exemplary Cas13d sequence from Gut_metagenome_contig3833000494.

SEQ ID NO: 44 is an exemplary Cas13d sequence from Activated_sludge_metagenome_transcript_117355.

SEQ ID NO: 45 is an exemplary Cas13d sequence from Gut_metagenome_contig11061000330.

SEQ ID NO: 46 is an exemplary Cas13d sequence from Gut_metagenome_contig338000322 from sheep gut metagenome.

SEQ ID NO: 47 is an exemplary Cas13d sequence from human gut metagenome.

SEQ ID NO: 48 is an exemplary Cas13d sequence from Gut_metagenome_contig9530000097.

SEQ ID NO: 49 is an exemplary Cas13d sequence from Gut_metagenome_contig1750000258.

SEQ ID NO: 50 is an exemplary Cas13d sequence from Gut_metagenome_contig5377000274.

SEQ ID NO: 51 is an exemplary Cas13d sequence from gut_metagenome_P19E0k2120140920_c248000089.

SEQ ID NO: 52 is an exemplary Cas13d sequence from Gut_metagenome_contig11400000031.

SEQ ID NO: 53 is an exemplary Cas13d sequence from Gut_metagenome_contig7940000191.

SEQ ID NO: 54 is an exemplary Cas13d sequence from Gut_metagenome_contig6049000251.

SEQ ID NO: 55 is an exemplary Cas13d sequence from Gut_metagenome_contig1137000500.

SEQ ID NO: 56 is an exemplary Cas13d sequence from Gut_metagenome_contig9368000105.

SEQ ID NO: 57 is an exemplary Cas13d sequence from Gut_metagenome_contig546000275.

SEQ ID NO: 58 is an exemplary Cas13d sequence from Gut_metagenome_contig7216000573.

SEQ ID NO: 59 is an exemplary Cas13d sequence from Gut_metagenome_contig4806000409.

SEQ ID NO: 60 is an exemplary Cas13d sequence from Gut_metagenome_contig10762000480.

SEQ ID NO: 61 is an exemplary Cas13d sequence from Gut_metagenome_contig4114000374.

SEQ ID NO: 62 is an exemplary Cas13d sequence from *Ruminococcus_flavifaciens_FDI*.

SEQ ID NO: 63 is an exemplary Cas13d sequence from Gut_metagenome_contig7093000170.

SEQ ID NO: 64 is an exemplary Cas13d sequence from Gut_metagenome_contig11113000384.

SEQ ID NO: 65 is an exemplary Cas13d sequence from Gut_metagenome_contig6403000259.

SEQ ID NO: 66 is an exemplary Cas13d sequence from Gut_metagenome_contig6193000124.

SEQ ID NO: 67 is an exemplary Cas13d sequence from Gut_metagenome_contig721000619.

SEQ ID NO: 68 is an exemplary Cas13d sequence from Gut_metagenome_contig1666000270.

SEQ ID NO: 69 is an exemplary Cas13d sequence from Gut_metagenome_contig2002000411.

SEQ ID NO: 70 is an exemplary Cas13d sequence from *Ruminococcus_albus*.

SEQ ID NO: 71 is an exemplary Cas13d sequence from Gut_metagenome_contig13552000311.

SEQ ID NO: 72 is an exemplary Cas13d sequence from Gut_metagenome_contig10037000527.

SEQ ID NO: 73 is an exemplary Cas13d sequence from Gut_metagenome_contig23_8000329.

SEQ ID NO: 74 is an exemplary Cas13d sequence from Gut_metagenome_contig2643000492.

SEQ ID NO: 75 is an exemplary Cas13d sequence from Gut_metagenome_contig874000057.

SEQ ID NO: 76 is an exemplary Cas13d sequence from Gut_metagenome_contig4781000489.

SEQ ID NO: 77 is an exemplary Cas13d sequence from Gut_metagenome_contig12144000352.

SEQ ID NO: 78 is an exemplary Cas13d sequence from Gut_metagenome_contig5590000448.

SEQ ID NO: 79 is an exemplary Cas13d sequence from Gut_metagenome_contig9269000031.

SEQ ID NO: 80 is an exemplary Cas13d sequence from Gut_metagenome_contig8537000520.

SEQ ID NO: 81 is an exemplary Cas13d sequence from Gut_metagenome_contig1845000130.

SEQ ID NO: 82 is an exemplary Cas13d sequence from gut_metagenome_P13E0k2120140920_c3000072.

SEQ ID NO: 83 is an exemplary Cas13d sequence from gut_metagenome_P1E0k2120140920_c1000078.

SEQ ID NO: 84 is an exemplary Cas13d sequence from Gut_metagenome_contig12990000099.

SEQ ID NO: 85 is an exemplary Cas13d sequence from Gut_metagenome_contig525000349.

SEQ ID NO: 86 is an exemplary Cas13d sequence from Gut_metagenome_contig7229000302.

SEQ ID NO: 87 is an exemplary Cas13d sequence from Gut_metagenome_contig3227000343.

SEQ ID NO: 88 is an exemplary Cas13d sequence from Gut_metagenome_contig7030000469.

SEQ ID NO: 89 is an exemplary Cas13d sequence from Gut_metagenome_contig5149000068.

SEQ ID NO: 90 is an exemplary Cas13d sequence from Gut_metagenome_contig400200045.

SEQ ID NO: 91 is an exemplary Cas13d sequence from Gut_metagenome_contig10420000446.

SEQ ID NO: 92 is an exemplary Cas13d sequence from new_flavefaciens_strain_XPD3002.

SEQ ID NO: 93 is an exemplary Cas13d sequence from M26_Gut_metagenome_contig698000307.

SEQ ID NO: 94 is an exemplary Cas13d sequence from M36_Uncultured_Eubacterium_sp_TS28_c40956.

SEQ ID NO: 95 is an exemplary Cas13d sequence from M12_gut_metagenome_P25C0k2120140920_c134000066.

SEQ ID NO: 96 is an exemplary Cas13d sequence from human gut metagenome.

SEQ ID NO: 97 is an exemplary Cas13d sequence from M10_gut_metagenome_P25C90k2120140920_c28000041.

SEQ ID NO: 98 is an exemplary Cas13d sequence from M11_gut_metagenome_P25C7k2120140920_c4078000105.

SEQ ID NO: 99 is an exemplary Cas13d sequence from gut_metagenome_P25C0k2120140920_c32000045.

SEQ ID NO: 100 is an exemplary Cas13d sequence from M13_gut_metagenome_P23C7k2120140920_c3000067.

SEQ ID NO: 101 is an exemplary Cas13d sequence from M5_gut_metagenome_P18E90k2120140920.

SEQ ID NO: 102 is an exemplary Cas13d sequence from M21_gut_metagenome_P18E0k2120140920.

SEQ ID NO: 103 is an exemplary Cas13d sequence from M7_gut_metagenome_P38C7k2120140920_c4841000003.

SEQ ID NO: 104 is an exemplary Cas13d sequence from *Ruminococcus bicirculans*.

SEQ ID NO: 105 is an exemplary Cas13d sequence.

SEQ ID NO: 106 is an exemplary Cas13d consensus sequence.

SEQ ID NO: 107 is an exemplary Cas13d sequence from M18_gut_metagenome_P22E0k2120140920_c3395000078.

SEQ ID NO: 108 is an exemplary Cas13d sequence from M17_gut_metagenome_P22E90k2120140920_c114.

SEQ ID NO: 109 is an exemplary Cas13d sequence from *Ruminococcus_sp_CAG57*.

SEQ ID NO: 110 is an exemplary Cas13d sequence from gut_metagenome_P1E90k2120140920_c43000123.

SEQ ID NO: 111 is an exemplary Cas13d sequence from M6_gut_metagenome_P13E90k2120140920_c7000009.

SEQ ID NO: 112 is an exemplary Cas13d sequence from M19_gut_metagenome_P17E90k2120140920.

SEQ ID NO: 113 is an exemplary Cas13d sequence from gut_metagenome_P17E0k2120140920_c87000043.

SEQ ID NO: 114 is an exemplary human codon optimized *Eubacterium siraeum* Cas13d nucleic acid sequence.

SEQ ID NO: 115 is an exemplary human codon optimized *Eubacterium siraeum* Cas13d nucleic acid sequence with a mutant HEPN domain.

SEQ ID NO: 116 is an exemplary human codon-optimized *Eubacterium siraeum* Cas13d nucleic acid sequence with N-terminal NLS.

SEQ ID NO: 117 is an exemplary human codon-optimized *Eubacterium siraeum* Cas13d nucleic acid sequence with N- and C-terminal NLS tags.

SEQ ID NO: 118 is an exemplary human codon-optimized uncultured *Ruminococcus* sp. Cas13d nucleic acid sequence.

SEQ ID NO: 119 is an exemplary human codon-optimized uncultured *Ruminococcus* sp. Cas13d nucleic acid sequence with a mutant HEPN domain

SEQ ID NO: 120 is an exemplary human codon-optimized uncultured *Ruminococcus* sp. Cas13d nucleic acid sequence with N-terminal NLS.

SEQ ID NO: 121 is an exemplary human codon-optimized uncultured *Ruminococcus* sp. Cas13d nucleic acid sequence with N- and C-terminal NLS tags.

SEQ ID NO: 122 is an exemplary human codon-optimized uncultured *Ruminococcus flavefaciens* FD1 Cas13d nucleic acid sequence.

SEQ ID NO: 123 is an exemplary human codon-optimized uncultured *Ruminococcus flavefaciens* FD1 Cas13d nucleic acid sequence with mutated HEPN domain.

SEQ ID NO: 124 is an exemplary Cas13d nucleic acid sequence from *Ruminococcus bicirculans*.

SEQ ID NO: 125 is an exemplary Cas13d nucleic acid sequence from *Eubacterium siraeum*.

SEQ ID NO: 126 is an exemplary Cas13d nucleic acid sequence from *Ruminococcus flavefaciens* FD1.

SEQ ID NO: 127 is an exemplary Cas13d nucleic acid sequence from *Ruminococcus albus*.

SEQ ID NO: 128 is an exemplary Cas13d nucleic acid sequence from *Ruminococcus flavefaciens* XPD.

SEQ ID NO: 129 is an exemplary consensus DR nucleic acid sequence for *E. siraeum* Cas13d.

SEQ ID NO: 130 is an exemplary consensus DR nucleic acid sequence for *Rum. Sp.* Cas13d.

SEQ ID NO: 131 is an exemplary consensus DR nucleic acid sequence for *Rum. flavefaciens* strain XPD3002 Cas13d and CasRx.

SEQ ID NOS: 132-137 are exemplary consensus DR nucleic acid sequences.

SEQ ID NO: 138 is an exemplary 50% consensus sequence for seven full-length Cas13d orthologues.

SEQ ID NO: 139 is an exemplary Cas13d nucleic acid sequence from Gut metagenome P1E0.

SEQ ID NO: 140 is an exemplary Cas13d nucleic acid sequence from Anaerobic digester.

SEQ ID NO: 141 is an exemplary Cas13d nucleic acid sequence from *Ruminococcus* sp. CAG:57.

SEQ ID NO: 142 is an exemplary human codon-optimized uncultured Gut metagenome P1E0 Cas13d nucleic acid sequence.

SEQ ID NO: 143 is an exemplary human codon-optimized Anaerobic Digester Cas13d nucleic acid sequence.

SEQ ID NO: 144 is an exemplary human codon-optimized *Ruminococcus flavefaciens* XPD Cas13d nucleic acid sequence.

SEQ ID NO: 145 is an exemplary human codon-optimized *Ruminococcus albus* Cas13d nucleic acid sequence.

SEQ ID NO: 146 is an exemplary processing of the *Ruminococcus* sp. CAG:57 CRISPR array.

15

SEQ ID NO: 147 is an exemplary Cas13d protein sequence from contig embLOBVH01003037.1, human gut metagenome sequence (also found in WGS contigs embLOBXZ01000094.11 and embLOBJF01000033.1).

SEQ ID NO: 148 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 147).

SEQ ID NO: 149 is an exemplary Cas13d protein sequence from contig tpgIDBYI01000091.11 (Uncultivated *Ruminococcus flavefaciens* UBA1190 assembled from bovine gut metagenome).

SEQ ID NOS: 150-152 are exemplary consensus DR nucleic acid sequences (goes with SEQ ID NO: 149).

SEQ ID NO: 153 is an exemplary Cas13d protein sequence from contig tpgIDJXD01000002.11 (uncultivated *Ruminococcus* assembly, UBA7013, from sheep gut metagenome).

SEQ ID NO: 154 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 153).

SEQ ID NO: 155 is an exemplary Cas13d protein sequence from contig OGZCO1000639.1 (human gut metagenome assembly).

SEQ ID NOS: 156-177 are exemplary consensus DR nucleic acid sequences (goes with SEQ ID NO: 155).

SEQ ID NO: 158 is an exemplary Cas13d protein sequence from contig embLOHBM01000764.1 (human gut metagenome assembly).

SEQ ID NO: 159 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 158).

SEQ ID NO: 160 is an exemplary Cas13d protein sequence from contig embLOHCP01000044.1 (human gut metagenome assembly).

SEQ ID NO: 161 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 160).

SEQ ID NO: 162 is an exemplary Cas13d protein sequence from contig embLOGDF01008514.11 (human gut metagenome assembly).

SEQ ID NO: 163 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 162).

SEQ ID NO: 164 is an exemplary Cas13d protein sequence from contig embLOGPN01002610.1 (human gut metagenome assembly).

SEQ ID NO: 165 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 164).

SEQ ID NO: 166 is an exemplary Cas13d protein sequence from contig NFIR01000008.1 (*Eubacterium* sp. An3, from chicken gut metagenome).

SEQ ID NO: 167 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 166).

SEQ ID NO: 168 is an exemplary Cas13d protein sequence from contig NFLV01000009.1 (*Eubacterium* sp. An 11, from chicken gut metagenome).

SEQ ID NO: 169 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 168).

SEQ ID NOS: 171-174 are an exemplary Cas13d motif sequences.

SEQ ID NO: 175 is an exemplary Cas13d protein sequence from contig OJMM01002900 human gut metagenome sequence.

SEQ ID NO: 176 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 175).

SEQ ID NO: 177 is an exemplary Cas13d protein sequence from contig ODAI011611274.1 gut metagenome sequence.

SEQ ID NO: 178 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 177).

SEQ ID NO: 179 is an exemplary Cas13d protein sequence from contig OIZX01000427.1.

16

SEQ ID NO: 180 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 179).

SEQ ID NO: 181 is an exemplary Cas13d protein sequence from contig embIOCVV012889144.1.

SEQ ID NO: 182 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 181).

SEQ ID NO: 183 is an exemplary Cas13d protein sequence from contig OCTW011587266.1

SEQ ID NO: 184 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 183).

SEQ ID NO: 185 is an exemplary Cas13d protein sequence from contig embLOGNF01009141.1.

SEQ ID NO: 186 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 185).

SEQ ID NO: 187 is an exemplary Cas13d protein sequence from contig embLOIEN01002196.1.

SEQ ID NO: 188 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 187).

SEQ ID NO: 189 is an exemplary Cas13d protein sequence from contig e-k87_11092736.

SEQ ID NOS: 190-193 are exemplary consensus DR nucleic acid sequences (goes with SEQ ID NO: 189).

SEQ ID NO: 194 is an exemplary Cas13d sequence from Gut_metagenome_contig6893000291.

SEQ ID NO: 195-197 are exemplary Cas13d motif sequences.

SEQ ID NO: 198 is an exemplary Cas13d protein sequence from Ga0224415_10007274.

SEQ ID NO: 199 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 198).

SEQ ID NO: 200 is an exemplary Cas13d protein sequence from EMG_10003641.

SEQ ID NO: 201 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 200).

SEQ ID NO: 202 is an exemplary Cas13d protein sequence from Ga0129306_1000735.

SEQ ID NO: 203 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 202).

SEQ ID NO: 204 is an exemplary Cas13d protein sequence from Ga0129317_1008067.

SEQ ID NO: 205 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 204).

SEQ ID NO: 206 is an exemplary Cas13d protein sequence from Ga0224415_10048792.

SEQ ID NO: 207 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 206).

SEQ ID NO: 208 is an exemplary Cas13d protein sequence from 160582958_gene49834.

SEQ ID NO: 209 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 208).

SEQ ID NO: 210 is an exemplary Cas13d protein sequence from 250twins_35838_GL0110300.

SEQ ID NO: 211 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 210).

SEQ ID NO: 212 is an exemplary Cas13d protein sequence from 250twins_36050_GL0158985.

SEQ ID NO: 213 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 212).

SEQ ID NO: 214 is an exemplary Cas13d protein sequence from 31009_GL0034153.

SEQ ID NO: 215 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 214).

SEQ ID NO: 216 is an exemplary Cas13d protein sequence from 530373_GL0023589.

SEQ ID NO: 217 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 216).

SEQ ID NO: 218 is an exemplary Cas13d protein sequence from BMZ-11B_GL0037771.

SEQ ID NO: 219 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 218).

SEQ ID NO: 220 is an exemplary Cas13d protein sequence from BMZ-11B_GL0037915.

SEQ ID NO: 221 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 220).

SEQ ID NO: 222 is an exemplary Cas13d protein sequence from BMZ-11B_GL0069617.

SEQ ID NO: 223 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 222).

SEQ ID NO: 224 is an exemplary Cas13d protein sequence from -DLF014_GL0011914.

SEQ ID NO: 225 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 224).

SEQ ID NO: 226 is an exemplary Cas13d protein sequence from EYZ-362B_GL0088915.

SEQ ID NO: 227-228 are exemplary consensus DR nucleic acid sequences (goes with SEQ ID NO: 226).

SEQ ID NO: 229 is an exemplary Cas13d protein sequence from Ga0099364_10024192.

SEQ ID NO: 230 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 229).

SEQ ID NO: 231 is an exemplary Cas13d protein sequence from Ga0187910_10006931.

SEQ ID NO: 232 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 231).

SEQ ID NO: 233 is an exemplary Cas13d protein sequence from Ga0187910_10015336.

SEQ ID NO: 234 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 233).

SEQ ID NO: 235 is an exemplary Cas13d protein sequence from Ga0187910_10040531.

SEQ ID NO: 236 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 235).

SEQ ID NO: 237 is an exemplary Cas13d protein sequence from Ga0187911_10069260.

SEQ ID NO: 238 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 237).

SEQ ID NO: 239 is an exemplary Cas13d protein sequence from MH0288_GL0082219.

SEQ ID NO: 240 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 239).

SEQ ID NO: 241 is an exemplary Cas13d protein sequence from 02.UC29-0_GL0096317.

SEQ ID NO: 242 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 241).

SEQ ID NO: 243 is an exemplary Cas13d protein sequence from PIG-014_GL0226364.

SEQ ID NO: 244 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 243).

SEQ ID NO: 245 is an exemplary Cas13d protein sequence from PIG-018_GL0023397.

SEQ ID NO: 246 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 245).

SEQ ID NO: 247 is an exemplary Cas13d protein sequence from PIG-025_GL0099734.

SEQ ID NO: 248 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 247).

SEQ ID NO: 249 is an exemplary Cas13d protein sequence from PIG-028_GL0185479.

SEQ ID NO: 250 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 249).

SEQ ID NO: 251 is an exemplary Cas13d protein sequence from -Ga0224422_10645759.

SEQ ID NO: 252 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 251).

SEQ ID NO: 253 is an exemplary Cas13d protein sequence from ODAI chimera.

SEQ ID NO: 254 is an exemplary consensus DR nucleic acid sequence (goes with SEQ ID NO: 253).

SEQ ID NO: 255 is an HEPN motif.

SEQ ID NOs: 256 and 257 are exemplary Cas13d nuclear localization signal amino acid and nucleic acid sequences, respectively.

SEQ ID NOs: 258 and 260 are exemplary SV40 large T antigen nuclear localization signal amino acid and nucleic acid sequences, respectively.

SEQ ID NO: 259 is a dCas9 target sequence.

SEQ ID NO: 261 is an artificial *Eubacterium siraeum* nCas1 array targeting ccdB.

SEQ ID NO: 262 is a full 36 nt direct repeat.

SEQ ID NOs: 263-266 are spacer sequences.

SEQ ID NO: 267 is an artificial uncultured *Ruminococcus* sp. nCas1 array targeting ccdB.

SEQ ID NO: 268 is a full 36 nt direct repeat.

SEQ ID NOs: 269-272 are spacer sequences.

SEQ ID NO: 273 is a ccdB target RNA sequence.

SEQ ID NOs: 274-277 are spacer sequences.

SEQ ID NO: 278 is a gRNA sequence.

SEQ ID NO: 279 is a consensus sequence of HEPN1 domain region.

SEQ ID NO: 280-286 are HEPN1 domain regions from seven Cas13d proteins.

SEQ ID NO: 287 is a consensus sequence of HEPN2 domain region.

SEQ ID NO: 288-294 are HEPN2 domain regions from seven Cas13d proteins.

SEQ ID NO: 295 is an exemplary RNA target sequence.

SEQ ID NO: 296-309 are exemplary gRNA sequences with various truncations.

SEQ ID NO: 310 is an exemplary Cas13d protein sequence from 037_-embIOIZA01000315.1.

SEQ ID NO: 311 is an exemplary Cas13d protein sequence from PIG-022_GL0026351.

SEQ ID NO: 312 is an exemplary Cas13d protein sequence from PIG-046_GL0077813.

SEQ ID NO: 313 is an exemplary Cas13d protein sequence from pig_chimera.

DETAILED DESCRIPTION

Unless otherwise noted, technical terms are used according to conventional usage. Definitions of common terms in molecular biology can be found in Benjamin Lewin, *Genes VII*, published by Oxford University Press, 1999; Kendrew et al. (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994; and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995; and other similar references.

As used herein, the singular forms “a,” “an,” and “the,” refer to both the singular as well as plural, unless the context clearly indicates otherwise. As used herein, the term “comprises” means “includes.” Thus, “comprising a nucleic acid molecule” means “including a nucleic acid molecule” without excluding other elements. It is further to be understood that any and all base sizes given for nucleic acids are approximate, and are provided for descriptive purposes, unless otherwise indicated. Although many methods and materials similar or equivalent to those described herein can be used, particular suitable methods and materials are

described below. In case of conflict, the present specification, including explanations of terms, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting. All references, including patent applications and patents, are herein incorporated by reference in their entireties.

In order to facilitate review of the various embodiments of the disclosure, the following explanations of specific terms are provided:

I. Terms

Administration: To provide or give a subject an agent, such as an Cas13d protein (or Cas13d coding sequence) or guide molecule (or coding sequence) disclosed herein, by any effective route. Exemplary routes of administration include, but are not limited to, injection (such as subcutaneous, intramuscular, intradermal, intraperitoneal, intratumoral, and intravenous), transdermal, intranasal, and inhalation routes.

Cas13d (also referred to as CasR, for CRISPR-associated RNase, and Cas13d): An RNA-guided RNA endonuclease enzyme that can cut or bind RNA. Cas13d proteins include one or two HEPN domains (e.g., see SEQ ID NOS: 1-3, 42, 62, 70, 82, 83, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, and 253). Native HEPN domains include the sequence RXXXXXH (SEQ ID NO: 255). Cas13d proteins that include mutated HEPN domain(s), and thus cannot cut RNA, but can process guide RNA, are also encompassed by this disclosure (e.g., see SEQ ID NOS: 2 and 4). An alignment of native Cas13d proteins is shown in FIGS. 18A-18MMM. In addition, Cas13d proteins specifically recognize direct repeat sequences of gRNA having a particular secondary structure (e.g. see FIG. 3C). In one example, Cas13d proteins recognize and/or bind a DR having (1) a loop of about 4 to 8 nt, (2) a stem of 4 to 12 nt, stem formed of complementary nucleotides, which can include a small (e.g., 1 or 2 bp) bulge due to a nt mismatch in the stem, and (3) a bulge or overhang formed of unpaired nt, which can be about 10 to 14 nt (e.g., 5 to 7 to on either side).

In one example, the full length (non-truncated) Cas13d protein is between 870-1080 amino acids long. In one example, the Cas13d protein is derived from a genome sequence of a bacterium from the Order Clostridiales or a metagenomic sequence. In one example, the corresponding DR sequence of an Cas13d protein is located at the 5' end of the spacer sequence in the molecule that includes the Cas13d gRNA. In one example, the DR sequence in the Cas13d gRNA is truncated at the 5' end relative to the DR sequence in the unprocessed Cas13d guide array transcript. In one example, the DR sequence in the Cas13d gRNA is truncated by 5-7 nt at the 5' end by the Cas13d protein. In one example, the Cas13d protein can cut a target RNA flanked at the 3' end of the spacer-target duplex by any of a A, U, G or C ribonucleotide and flanked at the 5' end by any of a A, U, G or C ribonucleotide.

In one example, an Cas13d protein has at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74,

75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253.

In one example, an Cas13d protein contains two HEPN RNase domains which contain a RXXXXXH amino acid motif (SEQ ID NO: 255; where X indicates any amino acid). In addition, an Cas13d protein can include one or more of the following amino acid motifs written in the common Prosite format:

Motif 1: (SEQ ID NO: 195)
 L-x(5)-[FWY]-x(3)-K-[NQS]-[ILM]-[ILMV]-x(2)-N-x(2)-[FWY]-x(2)-[AG]-x(4)-[DE]-x-D

Motif 2: (SEQ ID NO: 196)
 [FWY]-[ILV]-x(2)-[NQS]-[ILV]-x(2)-[DNST]-x(2)-F-x-Y-x(2)-[HKR]-[FHY]

Motif 3: (SEQ ID NO: 197)
 Y-[CDNSV]-x(2)-R-[FWY]-x-[ADNT]-[LM]-[ST]-x(4)-[FWY]

Thus, in some examples, an Cas13d protein having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253, includes the motif of SEQ ID NO: 195, 196 or 197.

Complementarity: The ability of a nucleic acid to form hydrogen bond(s) with another nucleic acid sequence by either traditional Watson-Crick base pairing or other non-traditional types. A percent complementarity indicates the percentage of residues in a nucleic acid molecule which can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 5, 6, 7, 8, 9, 10 out of 10 being 50%, 60%, 70%, 80%, 90%, and 100% complementary). "Perfectly complementary" means that all the contiguous residues of a nucleic acid sequence will hydrogen bond with the same number of contiguous residues in a second nucleic acid sequence. "Substantially complementary" as used herein refers to a degree of complementarity that is at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100% over a region of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, or more nucleotides, or refers to two nucleic acids that hybridize under stringent conditions.

CRISPRs (Clustered Regularly Interspaced Repeats): The CRISPR RNA array is a defining feature of CRISPR systems. The term "CRISPR" refers to the architecture of the array which includes constant direct repeats (DRs) interspaced with the variable spacers. In some examples, a CRISPR array includes at least a DR-spacer-DR-spacer (see FIG. 1A). This feature was used in the disclosed computa-

tional pipeline to identify the novel Cas13d protein family (FIG. 1A). In bacteria, the array is transcribed as one single transcript (containing multiple crRNA units), which is then processed by the Cas13d protein and other RNases into individual crRNAs. CRISPRs are found in approximately 40% of sequenced bacteria genomes and 90% of sequenced archaea. CRISPRs are often associated with cas genes that code for proteins related to CRISPRs (such as the Cas13d proteins provided herein). The disclosed CRISPR/Cas system can be used for RNA targeting, for example to detect a target RNA, modify a target RNA at any desired location, or cut the target RNA at any desired location.

Downregulated or knocked down: When used in reference to the expression of a molecule, such as a target RNA, refers to any process which results in a decrease in production of the target RNA, but in some examples not complete elimination of the target RNA product or target RNA function. In one example, downregulation or knock down does not result in complete elimination of detectable target RNA expression or target RNA activity. In some examples, the target RNA is a coding RNA. In some examples, the target RNA is non-coding RNA. Specific examples of RNA molecules that can be targeted for downregulation include mRNA, miRNA, rRNA, tRNA, nuclear RNA, lincRNA, circular RNA, and structural RNA. In some examples, downregulation or knock down of a target RNA includes processes that decrease translation of the target RNA and thus can decrease the presence of corresponding proteins. The disclosed CRISPR/Cas system can be used to downregulate any target RNA of interest.

Downregulation or knock down includes any detectable decrease in the target RNA. In certain examples, detectable target RNA in a cell or cell free system decreases by at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% (such as a decrease of 40% to 90%, 40% to 80% or 50% to 95%) as compared to a control (such as an amount of target RNA detected in a corresponding normal cell or sample). In one example, a control is a relative amount of expression in a normal cell (e.g., a non-recombinant cell that does not include Cas13d or guide RNA).

Effective amount: The amount of an agent (such as the CRISPR/Cas agents provided herein) that is sufficient to effect beneficial or desired results.

A therapeutically effective amount may vary depending upon one or more of: the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. The beneficial therapeutic effect can include enablement of diagnostic determinations; amelioration of a disease, symptom, disorder, or pathological condition; reducing or preventing the onset of a disease, symptom, disorder or condition; and generally counteracting a disease, symptom, disorder or pathological condition. In one embodiment, an "effective amount" is an amount sufficient to reduce symptoms of a disease, for example by at least 10%, at least 20%, at least 50%, at least 70%, or at least 90% (as compared to no administration of the therapeutic agent).

The term also applies to a dose that will allow for expression of an Cas13d and/or gRNA herein, and that allows for targeting (e.g., detection or modification) of a target RNA.

Increase or Decrease: A statistically significant positive or negative change, respectively, in quantity from a control value. An increase is a positive change, such as an increase

at least 50%, at least 100%, at least 200%, at least 300%, at least 400% or at least 500% as compared to the control value. A decrease is a negative change, such as a decrease of at least 20%, at least 25%, at least 50%, at least 75%, at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or at least 100% decrease as compared to a control value. In some examples the decrease is less than 100%, such as a decrease of no more than 90%, no more than 95% or no more than 99%.

Isolated: An "isolated" biological component (such as an Cas13d protein or nucleic acid, gRNA, or cell containing such) has been substantially separated, produced apart from, or purified away from other biological components in the cell or tissue of an organism in which the component occurs, such as other cells, chromosomal and extrachromosomal DNA and RNA, and proteins. Nucleic acids and proteins that have been "isolated" include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids and proteins. Isolated Cas13d proteins or nucleic acids, or cells containing such, in some examples are at least 50% pure, such as at least 75%, at least 80%, at least 90%, at least 95%, at least 98%, or at least 100% pure.

Label: A compound or composition that is conjugated directly or indirectly to another molecule (such as a nucleic acid molecule) to facilitate detection of that molecule. Specific, non-limiting examples of labels include fluorescent and fluorogenic moieties, chromogenic moieties, haptens, affinity tags, and radioactive isotopes. The label can be directly detectable (e.g., optically detectable) or indirectly detectable (for example, via interaction with one or more additional molecules that are in turn detectable).

Modulate: A change in the content of RNA. Modulation can include, but is not limited to, RNA activation (e.g., upregulation), RNA repression (e.g., downregulation), ribonucleotide deletion, ribonucleotide insertion, ribonucleotide chemical modification, ribonucleotide covalent or non-covalent linkage, and/or ribonucleotide substitution.

Non-naturally occurring or engineered: Terms used herein as interchangeably and indicate the involvement of the hand of man. The terms, when referring to nucleic acid molecules or polypeptides indicate that the nucleic acid molecule or the polypeptide is at least substantially free from at least one other component with which they are naturally associated in nature and as found in nature. In addition, the terms can indicate that the nucleic acid molecules or polypeptides have a sequence not found in nature.

Operably linked: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence (such as a coding sequence of an Cas13d protein) if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein-coding regions, in the same reading frame.

Pharmaceutically acceptable carriers: The pharmaceutically acceptable carriers useful in this invention are conventional. *Remington's Pharmaceutical Sciences*, by E. W. Martin, Mack Publishing Co., Easton, Pa., 15th Edition (1975), describes compositions and formulations suitable for pharmaceutical delivery of an Cas13d protein or nucleic acid molecule (or other molecules needed for modifying RNA using the disclosed CRISPR/Cas system with the disclosed Cas13d proteins).

In general, the nature of the carrier will depend on the particular mode of administration being employed. For instance, parenteral formulations usually comprise injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced salt solutions, aqueous dextrose, glycerol or the like as a vehicle. In addition to biologically-neutral carriers, pharmaceutical compositions to be administered can contain minor amounts of non-toxic auxiliary substances, such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate.

Polypeptide, peptide and protein: Refer to polymers of amino acids of any length. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation, such as conjugation with a labeling component. As used herein the term "amino acid" includes natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino acid analogs and peptidomimetics.

Promoter: An array of nucleic acid control sequences which direct transcription of a nucleic acid. A promoter includes necessary nucleic acid sequences near the start site of transcription. A promoter also optionally includes distal enhancer or repressor elements. A "constitutive promoter" is a promoter that is continuously active and is not subject to regulation by external signals or molecules. In contrast, the activity of an "inducible promoter" is regulated by an external signal or molecule (for example, a transcription factor).

Recombinant or host cell: A cell that has been genetically altered, or is capable of being genetically altered by introduction of an exogenous polynucleotide, such as a recombinant plasmid or vector. Typically, a host cell is a cell in which a vector can be propagated and its nucleic acid expressed. Such cells can be eukaryotic or prokaryotic. The term also includes any progeny of the subject host cell. It is understood that all progeny may not be identical to the parental cell since there may be mutations that occur during replication. However, such progeny are included when the term "host cell" is used.

Regulatory element: A phrase that includes promoters, enhancers, internal ribosomal entry sites (IRES), and other expression control elements (e.g., transcription termination signals, such as polyadenylation signals and poly-U sequences). Such regulatory elements are described, for example, in Goeddel, *GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY* 185, Academic Press, San Diego, Calif. (1990) which is hereby incorporated by reference in its entirety. Regulatory elements include those that direct constitutive expression of a nucleotide sequence in many types of host cells and those that direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). A tissue-specific promoter may direct expression primarily in a desired tissue of interest, such as muscle, neuron, bone, skin, blood, specific organs (e.g., liver, pancreas), or particular cell types (e.g., lymphocytes). Regulatory elements may also direct expression in a temporal-dependent manner, such as in a cell-cycle dependent or developmental stage-dependent manner, which may or may not also be tissue or cell-type specific.

In some embodiments, a vector provided herein includes a pol III promoter (e.g., U6 and H1 promoters), a pol II promoter (e.g., the retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer), the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1a promoter), or both.

Also encompassed by the term "regulatory element" are enhancer elements, such as WPRE; CMV enhancers; the R-U5' segment in LTR of HTLV-I; SV40 enhancer; and the intron sequence between exons 2 and 3 of rabbit β -globin.

RNA Editing: A type of genetic engineering in which a RNA molecule (or ribonucleotides of the RNA) is inserted, deleted or replaced in the genome of an organism using engineered nucleases (such as the Cas13d proteins provided herein), which create site-specific strand breaks at desired locations in the RNA. The induced breaks are repaired resulting in targeted mutations or repairs. The CRISPR/Cas methods disclosed herein, such as those that use an Cas13d, can be used to edit the sequence of one or more target RNAs, such as one associated with cancer (e.g., breast cancer, colon cancer, melanoma), infectious disease (such as HIV, hepatitis, HPV, and West Nile virus), or neurodegenerative disorder (e.g., Huntington's disease or ALS). For example, RNA editing can be used to treat a disease or viral infection.

RNA insertion site: A site of the RNA that is targeted for, or has undergone, insertion of an exogenous polynucleotide. The disclosed methods include use of a disclosed Cas13d protein, which can be used to target a RNA for manipulation at a RNA insertion site.

Sequence identity/similarity: The similarity between amino acid (or nucleotide) sequences is expressed in terms of the similarity between the sequences, otherwise referred to as sequence identity. Sequence identity is frequently measured in terms of percentage identity (or similarity or homology); the higher the percentage, the more similar the two sequences are.

Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith and Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman and Wunsch, *J. Mol. Biol.* 48:443, 1970; Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444, 1988; Higgins and Sharp, *Gene* 73:237, 1988; Higgins and Sharp, *CABIOS* 5:151, 1989; Corpet et al., *Nucleic Acids Research* 16:10881, 1988; and Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:2444, 1988. Altschul et al., *Nature Genet.* 6:119, 1994, presents a detailed consideration of sequence alignment methods and homology calculations.

The NCBI Basic Local Alignment Search Tool (BLAST) (Altschul et al., *J. Mol. Biol.* 215:403, 1990) is available from several sources, including the National Center for Biotechnology Information (NCBI, Bethesda, MD) and on the internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx. A description of how to determine sequence identity using this program is available on the NCBI website on the internet.

Variants of protein and nucleic acid sequences known in the art and disclosed herein are typically characterized by possession of at least about 80%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity counted over the full length alignment with the amino acid sequence using the NCBI Blast 2.0, gapped blastp set to default parameters. For comparisons of amino acid sequences of greater than about 30 amino acids,

the Blast 2 sequences function is employed using the default BLOSUM62 matrix set to default parameters, (gap existence cost of 11, and a per residue gap cost of 1). When aligning short peptides (fewer than around 30 amino acids), the alignment should be performed using the Blast 2 sequences function, employing the PAM30 matrix set to default parameters (open gap 9, extension gap 1 penalties). Proteins with even greater similarity to the reference sequences will show increasing percentage identities when assessed by this method, such as at least 95%, at least 98%, or at least 99% sequence identity. When less than the entire sequence is being compared for sequence identity, homologs and variants will typically possess at least 80% sequence identity over short windows of 10-20 amino acids, and may possess sequence identities of at least 85% or at least 90% or at least 95% depending on their similarity to the reference sequence. Methods for determining sequence identity over such short windows are available at the NCBI website on the internet. One of skill in the art will appreciate that these sequence identity ranges are provided for guidance only; it is entirely possible that strongly significant homologs could be obtained that fall outside of the ranges provided.

Thus, in one example, an Cas13d protein has at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 19, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253.

Subject: A vertebrate, such as a mammal, for example a human. Mammals include, but are not limited to, murines, simians, humans, farm animals, sport animals, and pets. In one embodiment, the subject is a non-human mammalian subject, such as a monkey or other non-human primate, mouse, rat, rabbit, pig, goat, sheep, dog, cat, horse, or cow. In some examples, the subject has a disorder (e.g., viral infection) or genetic disease that can be treated using methods provided herein. In some examples, the subject has a disorder (e.g., viral infection) or genetic disease that can be diagnosed using methods provided herein. In some examples, the subject is a laboratory animal/organism, such as a zebrafish, *Xenopus*, *C. elegans*, *Drosophila*, mouse, rabbit, or rat. Tissues, cells and their progeny of a biological entity obtained in vivo or cultured in vitro are also encompassed.

Therapeutic agent: Refers to one or more molecules or compounds that confer some beneficial effect upon administration to a subject. The beneficial therapeutic effect can include enablement of diagnostic determinations; amelioration of a disease, symptom, disorder, or pathological condition; reducing or preventing the onset of a disease, symptom, disorder or condition; and generally counteracting a disease, symptom, disorder or pathological condition.

Transduced, Transformed and Transfected: A virus or vector "transduces" a cell when it transfers nucleic acid molecules into a cell. A cell is "transformed" or "transfected" by a nucleic acid transduced into the cell when the

nucleic acid becomes stably replicated by the cell, either by incorporation of the nucleic acid into the cellular genome, or by episomal replication.

These terms encompass all techniques by which a nucleic acid molecule can be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of naked DNA by electroporation, lipofection, particle gun acceleration and other methods in the art. In some example the method is a chemical method (e.g., calcium-phosphate transfection), physical method (e.g., electroporation, microinjection, particle bombardment), fusion (e.g., liposomes), receptor-mediated endocytosis (e.g., DNA-protein complexes, viral envelope/capsid-DNA complexes) and biological infection by viruses such as recombinant viruses (Wolff, J. A., ed, *Gene Therapeutics*, Birkhauser, Boston, USA, 1994). Methods for the introduction of nucleic acid molecules into cells are known (e.g., see U.S. Pat. No. 6,110,743). These methods can be used to transduce a cell with the disclosed agents to manipulate its genome.

Transgene: An exogenous gene.

Treating, Treatment, and Therapy: Any success or indicia of success in the attenuation or amelioration of an injury, pathology or condition, including any objective or subjective parameter such as abatement, remission, diminishing of symptoms or making the condition more tolerable to the patient, slowing in the rate of degeneration or decline, making the final point of degeneration less debilitating, improving a subject's physical or mental well-being, or prolonging the length of survival. The treatment may be assessed by objective or subjective parameters; including the results of a physical examination, blood and other clinical tests, and the like. For prophylactic benefit, the disclosed compositions may be administered to a subject at risk of developing a particular disease, condition, or symptom, or to a subject reporting one or more of the physiological symptoms of a disease, even though the disease, condition, or symptom may not have yet been manifested.

Upregulated: When used in reference to the expression of a molecule, such as a target RNA, refers to any process which results in an increase in production of the target RNA. In one example, includes direct upregulation, for example if the target RNA participates in a feedback loop with its own transcription. In one example, includes indirect upregulation, such as by knockdown of an inhibitory miRNA that leads to the activation of a target of that miRNA.

In some examples, the target RNA is a coding RNA. In some examples, the target RNA is non-coding RNA. Specific examples of RNA molecules that can be targeted for upregulation include mRNA, miRNA, rRNA, tRNA, nuclear RNA, and structural RNA. In some examples, upregulation or activation of a target RNA includes processes that increase translation of the target RNA and thus can increase the presence of corresponding proteins. The disclosed CRISPR/Cas system can be used to upregulate any target RNA of interest.

Upregulation includes any detectable increase in target RNA. In certain examples, detectable target RNA expression in a cell or cell free system (such as a cell expressing an Cas13d protein and gRNA disclosed herein) increases by at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 90%, at least 95%, at least 100%, at least 200%, at least 400%, or at least 500% as compared to a control (such an amount of target RNA detected in a corresponding normal cell or sample). In one example, a control is a relative amount of

expression in a normal cell (e.g., a non-recombinant cell that does not include Cas13d or guide RNA).

Under conditions sufficient for: A phrase that is used to describe any environment that permits a desired activity. In one example the desired activity is expression of an Cas13d protein disclosed herein, in combination with other necessary elements, for example to modulate a target RNA.

Vector: A nucleic acid molecule into which a foreign nucleic acid molecule can be introduced without disrupting the ability of the vector to replicate and/or integrate in a host cell. Vectors include, but are not limited to, nucleic acid molecules that are single-stranded, double-stranded, or partially double-stranded; nucleic acid molecules that comprise one or more free ends, no free ends (e.g., circular); nucleic acid molecules that comprise DNA, RNA, or both; and other varieties of polynucleotides known in the art.

A vector can include nucleic acid sequences that permit it to replicate in a host cell, such as an origin of replication. A vector can also include one or more selectable marker genes and other genetic elements known in the art. An integrating vector is capable of integrating itself into a host nucleic acid. An expression vector is a vector that contains the necessary regulatory sequences to allow transcription and translation of inserted gene or genes.

One type of vector is a “plasmid,” which refers to a circular double stranded DNA loop into which additional DNA segments can be inserted, such as by standard molecular cloning techniques. Another type of vector is a viral vector, wherein virally-derived DNA or RNA sequences are present in the vector for packaging into a virus (e.g., retroviruses, replication defective retroviruses, adenoviruses, replication defective adenoviruses, and adeno-associated viruses). Viral vectors also include polynucleotides carried by a virus for transfection into a host cell. In some embodiments, the vector is a lentivirus (such as an integration-deficient lentiviral vector) or adeno-associated viral (AAV) vector.

Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome.

Certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as “expression vectors.” Common expression vectors are often in the form of plasmids. Recombinant expression vectors can comprise a nucleic acid provided herein (such as a guide RNA [which can be expressed from an RNA sequence or a RNA sequence], nucleic acid encoding an Cas13d protein) in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory elements, which may be selected on the basis of the host cells to be used for expression, that is operatively-linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, “operably linked” is intended to mean that the nucleotide sequence of interest is linked to the regulatory element(s) in a manner that allows for expression of the nucleotide sequence (e.g., in an in vitro transcription/translation system or in a host cell when the vector is introduced into the host cell). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression desired, etc. A vector can be introduced into host cells to thereby produce

transcripts, proteins, or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., clustered regularly interspersed short palindromic repeats (CRISPR) transcripts, proteins, enzymes, mutant forms thereof, fusion proteins thereof, etc.).

II. Overview of Several Embodiments

Class 2 CRISPR-Cas systems endow microbes with diverse mechanisms for adaptive immunity. Provided herein is an analysis of prokaryotic genome and metagenome sequences to identify an uncharacterized family of RNA-guided, RNA-targeting CRISPR systems which is classified as Type VI-D. Biochemical characterization and protein engineering of seven distinct orthologs generated a ribonuclease effector derived from *Ruminococcus flavefaciens* XPD3002 (CasRx) with robust activity in human cells. CasRx-mediated knockdown exhibits high efficiency and specificity relative to RNA interference across diverse endogenous transcripts. As one of the most compact single effector Cas enzymes, CasRx can also be flexibly packaged into adeno-associated virus. Virally encoded, catalytically inactive CasRx can be targeted to cis-elements of pre-mRNA to manipulate alternative splicing, alleviating dysregulated tau isoform ratios in a neuronal model of frontotemporal dementia. The results herein present CasRx as a programmable RNA-binding module for efficient targeting of cellular RNA, enabling a general platform for transcriptome engineering and therapeutic methods.

Class 2 CRISPR systems are found throughout diverse bacterial and archaeal life. Using a minimal definition of the CRISPR locus for bioinformatic mining of prokaryotic genome and metagenome sequences, which requires only a CRISPR repeat array and a nearby protein, provided herein is the identification of an uncharacterized, remarkably compact family of RNA-targeting class 2 CRISPR systems designated as Type VI CRISPR-Cas13d.

Because CRISPR systems generally exist as a functional operon within 20 kilobases of genome sequence, even fragmented metagenome reads may be sufficient to recover useful Cas enzymes for bioengineering purposes. CRISPR genome mining strategies described herein and by others (Shmakov et al., 2015), combined with ongoing efforts to profile microbial populations via next-generation sequencing, should contribute mechanistically diverse additions to the genome engineering toolbox.

Two distinct ribonuclease properties of the Cas13d effector, which processes a CRISPR repeat array into mature guides via a HEPN domain-independent mechanism followed by guide sequence-dependent recognition of a complementary activator RNA, were biochemically characterized. This triggers HEPN-mediated RNase activity, enabling Cas13d to cleave both activator and bystander RNAs, a property shared by other RNA-targeting CRISPR systems. Cas13d additionally exhibits no apparent flanking sequence requirements and was found to be active across crRNAs tiling a target RNA, suggesting the ability to target arbitrary single-stranded RNA sequences.

A comprehensive activity reporter screen in human cells of Cas13d orthologs sampled from distinct branches of the Cas13d family revealed that NLS fusions to Cas13d from *Ruminococcus flavefaciens* strain XPD3002 (CasRx) can be engineered for programmable RNA targeting in a eukaryotic context (FIG. 8D). CasRx fusions knocked down a diverse set of 14 endogenous mRNAs and lncRNAs, consistently achieving >90% knockdown with favorable efficiency relative to RNA interference, dCas9-mediated CRISPR interference, and other members of the Cas13 superfamily (FIGS. 10A-10C). Additionally, CasRx interference is mark-

edly more specific than spacer-matching shRNAs, with no detectable off-target changes compared with hundreds for RNA interference.

CasRx is a minimal two-component platform, including an engineered CRISPR-Cas13d effector and an associated guide RNA, and can be fully genetically encoded. Because CasRx is an orthogonally delivered protein, HEPN-inactive dCasRx can be engineered as a flexible RNA-binding module to target specific RNA elements. Importantly, because CasRx uses a distinct ribonuclease activity to process guide RNAs, dCasRx can still be paired with a repeat array for multiplexing applications. The utility of this concept is shown herein by creating a dCasRx splice effector fusion for tuning alternative splicing and resulting protein isoform ratios, applying it in a neuronal model of frontotemporal dementia.

At an average size of 930 aa, Cas13d is the smallest class 2 CRISPR effector characterized in mammalian cells. This allows CasRx effector domain fusions to be paired with a CRISPR array encoding multiple guide RNAs while remaining under the packaging size limit of the versatile adeno-associated virus (AAV) delivery vehicle (Naldini, 2015) for primary cell and in vivo delivery. Further, targeted AAV delivery of CasRx to specific postmitotic cell types such as neurons can mediate long-term expression of a corrective payload that avoids permanent genetic modifications or frequent re-administration (Chiriboga et al., 2016), complementing other nucleic acid targeting technologies such as DNA nuclease editing or antisense oligonucleotides. RNA mis-splicing diseases have been estimated to account for up to 15% of genetic diseases (Hammond and Wood, 2011), highlighting the potential for engineered splice effectors capable of multiplexed targeting. The materials provided herein can be used for RNA targeting for knockdown and splicing, such as live cell labeling and genetic screens to transcript imaging, trafficking, or regulation. CRISPR-Cas13d and engineered variants such as CasRx collectively enable flexible nucleic acid engineering, transcriptome-related study, and therapeutics, expanding the genome editing toolbox beyond DNA to RNA.

Provided herein are methods of targeting (e.g., modifying, detecting) one or more target RNA molecules, such as a clustered regularly interspaced short palindromic repeats (CRISPR)-CRISPR associated (Cas) system-mediated RNA editing method. Such methods can include contacting one or more target RNA molecules with a non-naturally occurring or engineered (e.g., does not naturally occur in the cell or system into which it is introduced) CRISPR-Cas system. Thus in some examples, the disclosed CRISPR-Cas system includes a naturally occurring Cas13d protein (or coding sequence) and a naturally occurring gRNA, but is used in a system or cell in which the Cas1 protein (or coding sequence) and the gRNA are not naturally found. Furthermore, the spacer sequence within the gRNA molecule is not naturally occurring, and has been modified to be complementary to the target RNA molecule.

In some examples, a target RNA is a coding RNA. In some examples, the RNA is non-coding RNA.

The disclosed CRISPR-Cas system can include (1) at least one Cas13d protein or at least one Cas13d nucleic acid coding sequence (such as a mRNA or a vector encoding the at least one Cas13d protein) and (2) at least one CRISPR-Cas system guide nucleic acid molecule (e.g., gRNA) (or at least one nucleic acid molecule encoding the gRNA) having sufficient complementary to a target RNA such that it can hybridize to a target RNA molecule. The Cas13d protein forms a complex with the gRNA, and the gRNA directs the

complex to the one or more target RNA molecules. This targeting can allow the Cas13d-gRNA complex to modify or detect the one or more target RNA molecules. In some examples, the one or more target RNA molecules (or a cell containing the one or more target RNA molecules) are contacted with a complex comprising the at least one Cas13d protein and the at least one gRNA. In some examples, the system includes Mg^{2+} . However, in some examples, the system does not require Mg^{2+} , such as if cleavage of the target RNA is not desired.

In some examples, contacting the one or more target RNA molecules with the non-naturally occurring or engineered CRISPR-Cas system includes introducing into a cell (such as a eukaryotic or prokaryotic cell) containing the one or more target RNA molecules the non-naturally occurring or engineered CRISPR-Cas system, for example using endocytosis (e.g., receptor-mediated endocytosis, micropinocytosis), a liposome, a particle, an exosome, a microvesicle, a gene gun, electroporation, a virus, RNP-antibody fusion (e.g., by tethering an Cas13d RNP to an antibody, antibody fragment, or other targeting moiety [such as ScFv, aptamers, DARPins, nanobodies, affibodies, etc.]), the RNP can be endocytosed into the cell, The RNP could conceivably be tethered to many things other than), or combinations thereof. Thus, cells can be transformed, transduced, transfected, or otherwise contacted with appropriate nucleic acid molecules of the disclosed CRISPR-Cas system. The resulting cells are recombinant cells. In some examples, contacting the one or more target RNA molecules with the non-naturally occurring or engineered CRISPR-Cas system includes contacting a cell-free system (such as a biological or environmental sample, or a cell lysate) containing the one or more target RNA molecules the non-naturally occurring or engineered CRISPR-Cas system (for example in a diagnostic method to detect a target RNA).

In some examples, at least 2, at least 3, at least 4, at least 5, at least 10, or at least 20 different gRNAs are used. For example, such a method could include targeting at least 2, at least 3, at least 4, at least 5, at least 10, or at least 20 different target RNA molecules, targeting at least 2, at least 3, at least 4, at least 5, at least 10, or at least 20 different regions of one or more RNA molecules, or combinations thereof.

Also provided are isolated nucleic acid molecules encoding such Cas13d proteins, such as a cDNA, genomic DNA, RNA, or mRNA. Such isolated nucleic acid molecules can be part of a vector (such as a plasmid or viral vector), and can be operably linked to a promoter. In some examples, an isolated nucleic acid molecule encoding a Cas13d protein has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 124, 125, 126, 127, 128, 139, 140 or 141; or at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 142, 143, 144, or 145. In an additional example, an isolated nucleic acid encodes a Cas13d protein having at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149,

153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253.

In some examples, an isolated nucleic acid molecule encoding at least one Cas13d protein (which can be part of a vector) includes at least one Cas13d protein coding sequence that is codon optimized for expression in a eukaryotic cell, at least one Cas13d protein coding sequence codon optimized for expression in a human cell. In one example, such an Cas13d coding sequence has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 142, 143, 144, or 145, or has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 124, 125, 126, 127, 128, 139, 140 or 141. In an additional example, a eukaryotic cell codon optimized nucleic acid sequence encodes a Cas13d protein having at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253.

In some examples, the gRNA that hybridizes with the one or more target RNA molecules includes one or more direct repeat (DR) sequences, one or more spacer sequences, or one or more sequences comprising DR-spacer-DR-spacer. In some examples, the one or more DR sequences have at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 129, 130, 131, 132, 133, 134, 135, 136, 137, 148, 150, 151, 152, 154, 156, 157, 159, 161, 163, 165, 167, 169, 176, 178, 180, 182, 184, 186, 188, 190, 191, 192, 193, 199, 201, 203, 205, 207, 209, 211, 213, 215, 217, 219, 221, 223, 225, 227, 228, 230, 232, 234, 236, 238, 240, 242, 244, 246, 248, 250, 252, or 254. In one example, the gRNA includes additional sequences, such as an aptamer sequence.

In some examples, a plurality of gRNAs are processed from a single array transcript, wherein each gRNA can be different, for example to target different RNAs or target multiple regions of a single RNA.

In some examples, the DRs are truncated by 1-10 nucleotides (such as 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides) at the 5'end, for example to be expressed as mature pre-processed guide RNAs.

Methods of targeting one or more target RNA molecules are provided. Targeting an RNA molecule can include one or more of cutting or nicking one or more target RNA molecules, activating or upregulating one or more target RNA molecules, activating or suppressing translation of the one or more target RNA molecules, deactivating the one or more target RNA molecules, visualizing, labeling, or detecting the one or more target RNA molecules, binding the one or more target RNA molecules, editing the one or more target RNA

molecules, trafficking the one or more target RNA molecules, and masking the one or more target RNA molecules. In some example, modifying one or more target RNA molecules includes one or more of an RNA base substitution, an RNA base deletion, an RNA base insertion, a break in the target RNA, methylating RNA, and demethylating RNA.

In some examples, such methods are used to treat a disease, such as a disease in a human. In such examples, the one or more target RNA molecules is associated with the disease.

Also provided are isolated proteins, including non-naturally occurring proteins. In some examples, a protein has at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253. In some examples, an isolated protein is an Cas13d ortholog from a prokaryotic genome or metagenome, gut metagenome, an activated sludge metagenome, an anaerobic digester metagenome, a chicken gut metagenome, a human gut metagenome, a pig gut metagenome, a bovine gut metagenome, a sheep gut metagenome, a goat gut metagenome, a capybara gut metagenome, a primate gut metagenome, a termite gut metagenome, a fecal metagenome, a genome from the Order Clostridiales, or the Family Ruminococcaceae. In some examples, an Cas13d ortholog includes an Cas13d ortholog from *Ruminococcus albus*, *Eubacterium siraeum*, a *Ruminococcus flavefaciens* strain XPD3002, *Ruminococcus flavefaciens* FD-1, uncultured *Eubacterium* sp TS28-c4095, uncultured *Ruminococcus* sp., *Ruminococcus bicirculans*, or *Ruminococcus* sp CAG57. Such proteins can include a subcellular localization signal. In some examples, such proteins include a mutation in at least one native HEPN domain.

Also provided are isolated guide RNA (gRNA) molecules. In some examples, an isolated gRNA includes one or more direct repeat (DR) sequences, such as one having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 129, 130, 131, 132, 133, 134, 135, 136, 137, 148, 150, 151, 152, 154, 156, 157, 159, 161, 163, 165, 167, 169, 176, 178, 180, 182, 184, 186, 188, 190, 191, 192, 193, 199, 201, 203, 205, 207, 209, 211, 213, 215, 217, 219, 221, 223, 225, 227, 228, 230, 232, 234, 236, 238, 240, 242, 244, 246, 248, 250, 252, or 254. Such a gRNA can further include one or more spacer sequences specific for (e.g., is complementary to) the target RNA. Such guide RNAs can further be optionally truncated by 1-10 nucleotides (such as 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides) at the 5'end of the DR, for example to generate pre-processed guide RNAs.

Also provided are ribonucleoprotein (RNP) complexes, which include one or more Cas13d proteins provided herein and one or more gRNAs provided herein.

Also provided are recombinant cells that include any Cas13d protein (or nucleic acid molecule encoding Cas13d),

any gRNA, any RNP complex, or any vector, provided herein. In one example, the cell is not a bacterial cell. In one example, the cell is a bacterial cell.

Also provided are compositions that include one or more of any Cas13d protein (or nucleic acid molecule encoding Cas13d), any gRNA or array, any RNP complex, any isolated nucleic acid molecule, any vector, or any cell, provided herein. Such compositions can include a pharmaceutically acceptable carrier.

Also provided are kits. Such kits can include one or more of any Cas13d protein (or nucleic acid molecule encoding Cas13d), any gRNA or array, any RNP complex, any isolated nucleic acid molecule, any vector, any cell, or any composition provided herein. Such reagents can be combined or in separate containers.

In some examples, a Cas13d protein is programmed toward its RNA target by combining the protein (or nucleic acid encoding the protein) with an engineered RNA guide (or nucleic acid encoding RNA guide) consisting of a full or partial direct repeat sequence followed by a "spacer" sequence complementary to the RNA target(s) (or variations thereof, i.e. arrays (DR-spacer-DR-spacer-DR-spacer . . . etc.) or pre-guide RNAs (DR-spacer-DR). Cas13d Proteins can be catalytically inactivated and transformed into RNA binding modules by mutating the conserved RNase HEPN motif (RXXXXH). Exemplary Cas13d proteins and corresponding guides are provided herein (e.g., SEQ ID NOS: 147-170, 175-193 and SEQ ID NOS: 198-254).

A. Cas13d Proteins

Provided herein are novel Cas13d proteins, such as those as shown in in the sequence listing. SEQ ID NOS: 1, 3, 42, 62, 70, 82, 83, 92, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, and 253 provide distinct full length proteins, and SEQ ID NOS: 2, 4-41, 43-61, 63-69, 71-81, 84-91, and 93-113 and 194 provide Cas13d variants and fragments. Such proteins can be used in the disclosed methods, compositions, and kits.

In some examples, an Cas13d protein includes one or more (such as 1 or 2) native HEPN domains. In some examples, an Cas13d protein includes one or more mutated HEPN domains (such as mutant Cas13d protein can process the gRNA, but cannot modify the target RNA). In some examples, an Cas13d protein is no more than 150 kD, no more than 140 kD, no more than 130 kD, no more than 120 kD, such as about 90 to 120 kD, about 100 to 120 kD or about 110 kD.

In addition to the Cas13d proteins provided in Table 1 and in Example 2, the disclosure encompasses Cas13d orthologs from a prokaryotic genome or metagenome, gut metagenome, an activated sludge metagenome, an anaerobic digester metagenome, a chicken gut metagenome, a human gut metagenome, a pig gut metagenome, a bovine gut metagenome, a sheep gut metagenome, a goat gut metagenome, a capybara gut metagenome, a primate gut metagenome, a termite gut metagenome, a fecal metagenome, a genome from the Order Clostridiales, or the Family Ruminococcaceae, such as an Cas13d ortholog from *Ruminococcus albus*, *Eubacterium siraeum*, a *Ruminococcus flavefaciens* strain XPD3002, *Ruminococcus flavefaciens* FD-1, uncultured *Eubacterium* sp TS28-c4095, uncultured *Ruminococcus* sp., *Ruminococcus bicirculans*, lor *Ruminococcus* sp CAG57.

In some examples, an Cas13d protein is at least 800 aa, at least 900 aa, or at least 1000 aa, such as 800 to 1200 aa, 850 to 1050 aa, or 860-1040 aa.

1. Variant Cas13d Sequences

Cas13d proteins, including variants of the sequences provided herein (such as variants of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) are encompassed within this disclosure. In some examples, Cas13d proteins provided herein can contain one or more mutations, such as a single insertion, a single deletion, a single substitution, or combinations thereof. In some examples, the Cas13d protein includes at least 1, at least 5, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, at least 75, at least 100, at least 200 or at least 300 aa insertions, such as 1-20 insertions (for example at the N- or C-terminus or within the protein, such as insertion of a whole small domain), at least 1, at least 5, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, at least 75, at least 100, at least 200 or at least 300 aa deletions (such as deletion of a whole small domain), such as 1-20 deletions (for example at the N- or C-terminus or within the protein), at least 1, at least 5, at least 10, at least 15, at least 20, at least 25, at least 30 aa substitutions, such as 1-20 substitutions, or any combination thereof (e.g., single insertion together with 1-19 substitutions), but retain the ability to bind target RNA molecules complementary to the spacer sequence within the gRNA molecule and/or process an guide array RNA transcript into gRNA molecules and/or retain the ability to cleave target RNA. In some examples, the disclosure provides a variant of any disclosed Cas13d protein (such as SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) having 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 amino acid changes, but retain the ability to bind target RNA molecules complementary to the spacer sequence within the gRNA molecule and/or process an guide array RNA transcript into gRNA molecules. In some examples, any disclosed Cas13d protein (such as SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175,

177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) further includes 1-8 amino acid insertions, 1-15 amino acid deletions, 1-10 amino acid substitutions, or any combination thereof (e.g., 1-15, 1-4, or 1-5 amino acid deletions together with 1-10, 1-5 or 1-7 amino acid substitutions), with the retained ability to bind target RNA molecules complementary to the spacer sequence within the gRNA molecule and/or process an guide array RNA transcript into gRNA molecules. In one example, such variant peptides are produced by manipulating the nucleotide sequence encoding a peptide using standard procedures such as site-directed mutagenesis or PCR. Such variants can also be chemically synthesized.

In some examples, an Cas13d protein includes a motif shown in SEQ ID NO: 195, 196 or 197. Thus, an Cas13d protein having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253, in some examples includes at least one motif shown in SEQ ID NO: 195, 196 or 197.

One type of modification or mutation includes the substitution of amino acids for amino acid residues having a similar biochemical property, that is, a conservative substitution (such as 1-4, 1-8, 1-10, or 1-20 conservative substitutions). Typically, conservative substitutions have little to no impact on the activity of a resulting peptide. For example, a conservative substitution is an amino acid substitution in an Cas13d protein (such as SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) that does not substantially affect the ability of the Cas13d protein to bind target RNA molecules complementary to the spacer sequence within the gRNA molecule and/or process an guide array RNA transcript into gRNA molecules. An alanine scan can be used to identify which amino acid residues in an Cas13d protein (such as SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202,

204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253), can tolerate an amino acid substitution. In one example, the ability of a variant Cas13d protein (such as 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) to modify gene expression in a CRISPR/Cas system, is not altered by more than 25%, for example not more than 20%, for example not more than 10%, when an alanine, or other conservative amino acid, is substituted for 1-4, 1-8, 1-10, or 1-20 native amino acids. Examples of amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative substitutions include: Ser for Ala; Lys for Arg; Gln or His for Asn; Glu for Asp; Ser for Cys; Asn for Gln; Asp for Glu; Pro for Gly; Asn or Gln for His; Leu or Val for Ile; Ile or Val for Leu; Arg or Gln for Lys; Leu or Ile for Met; Met, Leu or Tyr for Phe; Thr for Ser; Ser for Thr; Tyr for Trp; Trp or Phe for Tyr; and Ile or Leu for Val.

One method for identifying regions particularly amenable to insertions, substitutions or deletion is to target stretches of amino acids exhibiting low levels of conservation between orthologs. Such regions are indicated in the conservation graph of the alignment of Cas13d proteins provided in FIG. 1B. Conserved residues of Cas13d are further marked in the Cas13d protein alignment provided in FIG. 18A-18MMM (indicated by symbols “.” “:” or “*” below aligned conserved residues). Examples of deletions and their functional testing are further provided in FIGS. 16A-16B.

Another type of substitution can be achieved by swapping out parts of one ortholog with the homologous region of another ortholog to obtain a combined “chimeric” protein. Such a chimeric protein may combine favorable properties of multiple Cas13d orthologs.

More substantial changes can be made by using substitutions that are less conservative, e.g., selecting residues that differ more significantly in their effect on maintaining: (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation; (b) the charge or hydrophobicity of the polypeptide at the target site; or (c) the bulk of the side chain. The substitutions that in general are expected to produce the greatest changes in polypeptide function are those in which: (a) a hydrophilic residue, e.g., serine or threonine, is substituted for (or by) a hydrophobic residue, e.g., leucine, isoleucine, phenylalanine, valine or alanine; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysine, arginine, or histidine, is substituted for (or by) an electronegative residue, e.g., glutamic acid or aspartic acid; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

Thus, the disclosure provides Cas13d proteins having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39,

40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253 or combinations (e.g., chimeras) thereof.

In one example, an Cas13d protein includes non-naturally occurring amino acids.

2. Cas13d proteins with Other Elements

An Cas13d protein (such as any of SEQ ID NOS: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) can include other elements or domains, for example at the N- or C-terminus (or both). The resulting protein can be referred to as an Cas13d fusion protein. In one example, an Cas13d protein provided herein (such as a native Cas13d or an Cas13d with mutated HEPN domain(s)) includes a subcellular localization signal. Exemplary subcellular localization signals include an organelle localization signal, such as a nuclear localization signal (NLS), nuclear export signal (NES), or mitochondrial localization signal. In one example, an Cas13d protein includes an NLS, such as SPKKKRKVEAS (SEQ ID NO: 256; e.g., encoded by AGCCCCAAGAAgAAGAGaAAGGTGGAGGCCAGC, SEQ ID NO: 257) or GPKKKRKVAAA (SV40 large T antigen NLS, SEQ ID NO: 258; e.g., encoded by ggacctaagaaaagaggaaggtggcggcgcgct, SEQ ID NO: 260). Exemplary NES that can be part of an Cas13d protein include an adenovirus type 5 E1B nuclear export sequence, an HIV nuclear export sequence, a MAPK nuclear export sequence, or a PTK2 nuclear export sequence.

In some examples, the at least one Cas13d protein (such as a native Cas13d or an Cas13d with mutated HEPN domain(s)) further includes one or more effector domains. Exemplary effector domains include proteins and/or enzymes, such that those can cleave RNA (e.g., a PIN endonuclease domain, an NYN domain, an SMR domain from SOT1, or an RNase domain from Staphylococcal nuclease), those that can affect RNA stability (e.g., tristetraprolin (TTP) or domains from UPF1, EXOSC5, and STAU1), those that can edit a nucleotide or ribonucleotide (e.g., a cytidine deaminase, PPR protein, adenosine deaminase, ADAR family protein, or APOBEC family protein), those that can activate translation (e.g., eIF4E and other translation initiation factors, a domain of the yeast poly(A)-binding protein or GLD2), those that can repress translation (e.g., Pumilio or FBF PUF proteins, deadenylases, CAF1, Argonaute proteins), those that can methylate RNA (e.g., domains from m6A methyltransferase factors such as METTL14, METTL3, or WTAP), those that can demethylate RNA (e.g., human alkylation repair homolog 5 or Alkbh5), those that can affect splicing (e.g., the RS-rich domain of SRSF1, the Gly-rich domain of hnRNP A1, the alanine-rich motif of RBM4, or the proline-rich motif of

DAZAP1), those that can enable affinity purification or immunoprecipitation (e.g., FLAG, HA, biotin, or HALO tags), and those that can enable proximity-based protein labeling and identification (e.g., a biotin ligase (such as BirA) or a peroxidase (such as APEX2) in order to biotinylate proteins that interact with the target RNA).

In some examples, the Cas13d protein and effector module combination can constitute a transcriptional sensor. For example, the transcriptional sensor can be comprised of at least one Cas13d protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4), at least one gRNA containing at least one spacer sequence specific for the target RNA, and an effector module such as an optionally split fluorescent protein or probe (e.g., a split Venus fluorescent protein, a split GFP, a split enhanced GFP, a split mCherry, a split super-folder mCherry, and other fluorescent protein variants such as ECFP, YFP, RFP, and derivatives or fragments thereof); an optionally split luminescent protein or probe (e.g. *Gaussia*, Firefly, NanoLuc, or *Renilla* variants); an optionally split enzyme (e.g., ubiquitin or TEV protease); a FRET-compatible protein pair; one or more transcription factor(s) fused to Cas13d via cleavable linkers (e.g., an artificial GAL4, zinc finger, transcriptional activator like effector (TALE), CRISPR-Cas9, CRISPR-Cpf1, or TetR-based transcription factor or an endogenous transcription factor); a split intein that trans-splices a protein to restore its function such as a transcription factor (e.g., an intein from *Rhodothermus marinus* or DnaE); a kinase-substrate pair that activates upon phosphorylation (e.g., TYK2-STAT3); one, two, or more monomers that activate upon dimerization or multimerization (e.g., caspase 9); or one or more proteins that induce conformational and functional change upon interaction. In one example, the spatial proximity of two or more Cas13d proteins and gRNAs due to binding a particular transcript would activate the effector module, resulting in a detectable signal or detectable activity in the cell.

In one example, the effector domain is fused to a protein that specifically recognizes and binds an RNA aptamer, such as one that can be appended to or inserted within a gRNA molecule (e.g., an MS2, PP7, Q β , and other aptamers). This aptamer-effector domain fusion can be used to target the target RNA because the Cas13d and gRNA complex will guide the aptamer protein-effector domain in proximity to the target RNA.

In another example, the aptamer can be directly inserted into the gRNA molecule to permit detection of a target RNA, such as a fluorophore aptamer (e.g., Spinach, Mango, etc.).

In some examples, the Cas13d protein (such as a native Cas13d or an Cas13d with mutated HEPN domain(s)) includes a purification tag, such as an HA-tag, His-tag (such as 6-His), Myc-tag, E-tag, S-tag, calmodulin tag, FLAG-tag, GST-tag, MBP-tag, and the like. Such tags are in some examples at the N- or C-terminal end of the Cas13d protein.

In some examples, an Cas13d protein (such as a native Cas13d or an Cas13d with mutated HEPN domain(s)) includes one or more subcellular localization signals, effector domains, and purification tags.

In some examples, an Cas13d protein may be split into multiple fragments, which are then expressed individually. Such fragments of Cas13d may be optionally fused to other protein domains. In one example, an Cas13d can be split into two halves, which are then fused to two parts of an inducible heterodimer pair. Upon induction of heterodimer binding, the Cas13d halves are recruited to each other to form an active protein. Such a system would allow for the inducible control of Cas13d activity. Useful heterodimer pairs include two proteins that dimerize upon light illumination or through

administration of a small molecule compound, amongst others. Specific examples of heterodimer pair include but are not limited to: light inducible Magnets proteins, the light inducible iLID-SspB pair, the light inducible Cryptochrome2-CIB1 dimer and the small molecule inducible FKBP protein. In another example of a split Cas13d design, two halves of the Cas13d protein may be fused to protein trans-splicing domains. Such a design would enable the separate expression of two halves which are reconstituted into a full length protein once expressed inside a cell. An example of such transsplicing domains includes the Intein system.

One method for identifying regions particularly amenable to splitting of the protein, is to identify stretches of amino acids exhibiting low levels of conservation between orthologs. Such regions are indicated in the conservation graph of the alignment of Cas13d proteins provided in FIG. 1B. Regions of conserved residues of Cas13d are further marked in the Cas13d protein alignment provided in FIG. 18A-18MMM (indicated by symbols . : or * below aligned conserved residues).

3. Generation of Cas13d Proteins

In one example, the Cas13d protein is expressed in vitro, for example, in a prokaryotic cell (e.g., bacteria such as *Lactobacillus*, *Lactococcus*, *Bacillus* (such as *B. subtilis*), *Escherichia* (such as *E. coli*), *Salmonella typhimurium*, and *Clostridium*), archaea cell, plant or plant cell, fungal cell (e.g., *Neurospora*), yeast cell (e.g., *Saccharomyces* or *Pichia* (such as *S. cerevisiae* or *P. pastoris*), *Kluyveromyces lactis*), insect cell (e.g., SF9 cells), or mammalian cells (e.g., 293 cells, or immortalized mammalian myeloid and lymphoid cell lines). Once expressed, the Cas13d protein can be isolated and/or purified (e.g., using chromatography or immunological separation). In some examples, as tag on the Cas13d protein permits isolation of the protein from a culture media. Exemplary procedures include ammonium sulfate precipitation, affinity columns, column chromatography, and the like (see, generally, R. Scopes, *Protein Purification*, Springer-Verlag, N.Y., 1982). Substantially pure compositions of at least about 90 to 95% homogeneity, such as 98% to 99% homogeneity, can be used in the methods provided herein. For example, a purified preparation of an Cas13d protein can be used as an alternative to expressing the Cas13d protein from a nucleic acid molecule in the CRISPR/Cas system.

In addition to recombinant methods, Cas13d proteins disclosed herein can also be constructed in whole or in part using native chemical ligation and/or expressed protein ligation.

B. Nucleic Acid Molecules Encoding Cas13d Proteins

Nucleic acid molecules encoding an Cas13d protein are encompassed by this disclosure. Nucleic acid molecules include DNA, genomic DNA, cDNA, mRNA, and RNA sequences which encode an Cas13d peptide. Such nucleic acid molecules can include naturally occurring or non-naturally occurring nucleotides or ribonucleotides. Exemplary nucleic acid molecules that encode the novel Cas13d proteins shown in SEQ ID NOS: 1, 3, 42, 62, 70, 82, 83, 92 and 104, are shown in SEQ ID NOS: 124-128, 139, 140, and 141. Also provided are codon optimized nucleic acid molecules that encode the novel Cas13d proteins, for example those optimized for expression in a mammalian cells, such as a human cell (SEQ ID NOS: 114-123 and 142-145). For example, SEQ ID NOS: 114, 118, and 122 provide nucleic acid molecules optimized for expression in human cells, and

which encode for mutant HEPN sites. SEQ ID NOS: 116 and 120 provide nucleic acid molecules optimized for expression in human cells, and which includes an N-terminal nuclear localization (NLS) coding sequence (namely, SPKKKRRKVEAS). SEQ ID NO: 117 and 121 provide nucleic acid molecules optimized for expression in human cells, and which include N-terminal and C-terminal NLS coding sequences (namely, SPKKKRRKVEAS, SEQ ID NO: 256, and GPKKKRRKVAAA SEQ ID NO: 258, respectively).

In one example, a nucleic acid sequence encodes an Cas13d protein having at least 60%, at least 70%, at least 75%, at least 80%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 99% or at least 99% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253. Such nucleic acid sequences can be generated based on the amino acid sequences provided herein, and the genetic code. In one example, an Cas13d nucleic acid sequence has at least 70%, at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 124, 125, 126, 127, 128, 139, 140 or 141. In one example, an Cas13d nucleic acid sequence is optimized for expression in mammalian cells, such as human cells, such as one having at least 70%, at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 142, 143, 144, or 145.

One of skill can readily construct a variety of clones containing functionally equivalent nucleic acids, such as nucleic acids which differ in sequence but which encode the same Cas13d protein sequence. Silent mutations in the coding sequence result from the degeneracy (i.e., redundancy) of the genetic code, whereby more than one codon can encode the same amino acid residue. Thus, for example, leucine can be encoded by CTT, CTC, CTA, CTG, TTA, or TTG; serine can be encoded by TCT, TCC, TCA, TCG, AGT, or AGC; asparagine can be encoded by AAT or AAC; aspartic acid can be encoded by GAT or GAC; cysteine can be encoded by TGT or TGC; alanine can be encoded by GCT, GCC, GCA, or GCG; glutamine can be encoded by CAA or CAG; tyrosine can be encoded by TAT or TAC; and isoleucine can be encoded by ATT, ATC, or ATA. Tables showing the standard genetic code can be found in various sources (see, for example, Stryer, 1988, *Biochemistry*, 3rd Edition, W.H. Freeman and Co., NY).

Based on the genetic code, nucleic acid sequences coding for any Cas13d sequence can be generated. In some examples, such a sequence is optimized for expression in a host or target cell, such as a host cell used to express the Cas13d protein or a cell in which the disclosed methods are practiced (such as in a mammalian cell, such as a human cell). Codon preferences and codon usage tables for a particular species can be used to engineer isolated nucleic acid molecules encoding an Cas13d (such as one encoding a protein having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4,

5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253 that takes advantage of the codon usage preferences of that particular species). For example, the Cas13d proteins disclosed herein can be designed to have codons that are preferentially used by a particular organism of interest. In one example, an Cas13d nucleic acid sequence is optimized for expression in human cells, such as one having at least 70%, at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 142, 143, 144, or 145.

A nucleic acid encoding an Cas13d protein (such as one encoding a protein having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) can be cloned or amplified by in vitro methods, such as the polymerase chain reaction (PCR), the ligase chain reaction (LCR), the transcription-based amplification system (TAS), the self-sustained sequence replication system (3SR) and the Q β replicase amplification system (QB). In addition, nucleic acids encoding an Cas13d protein (such as one encoding a protein having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) can be prepared by cloning techniques. Examples of appropriate cloning and sequencing techniques, and instructions sufficient to direct persons of skill through cloning are found in Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual* 2nd ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989, and Ausubel et al., (1987) in "Current Protocols in Molecular Biology," John Wiley and Sons, New York, N.Y.

Nucleic acid sequences encoding an Cas13d protein (such as one encoding a protein having at least 80%, at least 85%,

at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) can be prepared by any suitable method including, for example, cloning of appropriate sequences or by direct chemical synthesis by methods such as the phosphotriester method of Narang et al., *Meth. Enzymol.* 68:90-99, 1979; the phosphodiester method of Brown et al., *Meth. Enzymol.* 68:109-151, 1979; the diethylphosphoramidite method of Beaucage et al., *Tetra. Lett.* 22:1859-1862, 1981; the solid phase phosphoramidite triester method described by Beaucage & Caruthers, *Tetra. Letts.* 22(20):1859-1862, 1981, for example, using an automated synthesizer as described in, for example, Needham-VanDevanter et al., *Nucl. Acids Res.* 12:6159-6168, 1984; and, the solid support method of U.S. Pat. No. 4,458,066. Chemical synthesis produces a single stranded oligonucleotide. This can be converted into double stranded DNA by hybridization with a complementary sequence, or by polymerization with a DNA polymerase using the single strand as a template. One of skill would recognize that while chemical synthesis of DNA is generally limited to sequences of about 100 bases, longer sequences may be obtained by the ligation of shorter sequences.

In one example, an Cas13d protein (such as a protein having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) is prepared by inserting the cDNA which encodes the Cas13d protein into a plasmid or vector. The insertion can be made so that the Cas13d protein is read in frame so that the Cas13d protein is produced.

The Cas13d nucleic acid coding sequence (such as one having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226,

229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) can be inserted into an expression vector including, but not limited to a plasmid, virus or other vehicle that can be manipulated to allow insertion or incorporation of sequences and can be expressed in either prokaryotes or eukaryotes. Hosts can include microbial, yeast, insect, plant and mammalian organisms. The vector can encode a selectable marker, such as a thymidine kinase gene or antibiotic resistance gene.

Nucleic acid sequences encoding an Cas13d protein (such as one encoding a protein having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) can be operatively linked to expression control sequences. An expression control sequence operatively linked to an Cas13d coding sequence is ligated such that expression of the Cas13d protein coding sequence is achieved under conditions compatible with the expression control sequences. The expression control sequences include, but are not limited to appropriate promoters, enhancers, transcription terminators, a start codon (i.e., ATG) in front of an Cas13d protein-encoding gene, splicing signal for introns, maintenance of the correct reading frame of that gene to permit proper translation of mRNA, and stop codons.

In one embodiment, vectors are used for expression in yeast such as *S. cerevisiae*, *P. pastoris*, or *Kluyveromyces lactis*. Exemplary promoters for use in yeast expression systems include but are not limited to: the constitutive promoters plasma membrane H⁺-ATPase (PMA1), glyceraldehyde-3-phosphate dehydrogenase (GPD), phosphoglycerate kinase-1 (PGK1), alcohol dehydrogenase-1 (ADH1), and pleiotropic drug-resistant pump (PDR5). In addition, many inducible promoters are of use, such as GAL1-10 (induced by galactose), PHO5 (induced by low extracellular inorganic phosphate), and tandem heat shock HSE elements (induced by temperature elevation to 37° C.). Promoters that direct variable expression in response to a titratable inducer include the methionine-responsive MET3 and MET25 promoters and copper-dependent CUP1 promoters. Any of these promoters may be cloned into multicopy (2 μ) or single copy (CEN) plasmids to give an additional level of control in expression level. The plasmids can include nutritional markers (such as URA3, ADE3, HIS1, and others) for selection in yeast and antibiotic resistance (AMP) for propagation in bacteria. Plasmids for expression on *K. lactis* are known, such as pKLAC1.

Viral vectors can also be prepared that encode an Cas13d (such as one encoding a protein having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76,

77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253). Exemplary viral vectors include polyoma, SV40, adenovirus, vaccinia virus, adeno-associated virus, herpes viruses including HSV and EBV, lentivirus, Sindbis viruses, alphaviruses and retroviruses of avian, murine, and human origin. Baculovirus (*Autographa californica* multinuclear polyhedrosis virus; AcMNPV) vectors can be used and obtained from commercial sources. Other suitable vectors include retrovirus vectors, orthopox vectors, avipox vectors, fowlpox vectors, capripox vectors, suipox vectors, adenoviral vectors, herpes virus vectors, alpha virus vectors, baculovirus vectors, Sindbis virus vectors, vaccinia virus vectors and poliovirus vectors. Specific exemplary vectors are poxvirus vectors such as vaccinia virus, fowlpox virus and a highly attenuated vaccinia virus (MVA), adenovirus, baculovirus and the like. Pox viruses of use include orthopox, suipox, avipox, and capripox virus. Orthopox include vaccinia, ectromelia, and raccoon pox. One example of an orthopox of use is vaccinia. Avipox includes fowlpox, canary pox and pigeon pox. Capripox include goatpox and sheeppox. In one example, the suipox is swinepox. Other viral vectors that can be used include other DNA viruses such as herpes simplex virus and adenoviruses, and RNA viruses such as retroviruses and polio.

Viral vectors that encode an Cas13d protein (such as one encoding a protein having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) can include at least one expression control element operatively linked to the nucleic acid sequence encoding the Cas13d protein. The expression control elements control and regulate the expression of the Cas13d nucleic acid sequence. Exemplary expression control elements that can be used include, but are not limited to, lac system, operator and promoter regions of phage lambda, yeast promoters and promoters derived from polyoma, adenovirus, retrovirus or SV40. In one example the promoter is CMV, U6, CBh, CMW, Cbh, EF1a. In one example, the promoter is a cell type specific promoter, such as synapsin or GFAP, or an inducible promoter, such as a tetracycline inducible promoter. Additional operational elements include, but are not limited to, leader sequence, termination codons, polyadenylation signals and any other sequences necessary for the appropriate transcription and subsequent translation of the nucleic acid sequence encoding the Cas13d protein in the host system. The expression vector can contain additional elements necessary for the transfer and subsequent replication of the expression vector containing the nucleic acid sequence in the host system. Examples

of such elements include, but are not limited to, origins of replication and selectable markers.

In one example, the vector includes a polyA signal after the Cas13d protein coding sequence, a WPRE signal for expression in viral vectors, or combinations thereof.

In one example, the method uses direct delivery of an mRNA that encodes for an Cas13d protein.

C. Guide Nucleic Acid Molecules

The disclosure provides guide nucleic acid molecules, such as guide RNA (gRNA or crRNA, CRISPR (guide) RNA), which can be used in the methods, compositions, and kits provided herein. Such molecules can include naturally occurring or non-naturally occurring nucleotides or ribonucleotides (such as LNAs or other chemically modified nucleotides or ribonucleotides, for example to protect a guide RNA from degradation). In some examples, the guide sequence is RNA. The guide nucleic acid can include modified bases or chemical modifications (e.g., see Latorre et al., *Angewandte Chemie* 55:3548-50, 2016). A guide sequence directs a Cas13d protein to a target RNA, thereby targeting the RNA (e.g., modifying or detecting the RNA).

Guide molecules include one or more regions referred to as spacers. A spacer has sufficient complementarity with a target RNA sequence to hybridize with the target RNA and direct sequence-specific binding of an Cas13d protein to the target RNA. Thus, the spacer is the variable portion of the guide sequence. In some examples, a spacer has 100% complementarity to a target RNA (or region of the RNA to be target), but a spacer can have less than 100% complementarity to a target RNA, such as at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% complementarity to a target RNA.

A guide sequence can also include one or more direct repeats (DRs). The DR is the constant portion of the guide, which contains strong secondary structure (FIG. 3C), which facilitate interaction between a Cas13d protein and the guide molecule. Each ortholog has a slightly different DR sequence (e.g., SEQ ID NOS: 129, 130, 131, 132, 133, 134, 135, 136, 137, 148, 150, 151, 152, 154, 156, 157, 159, 161, 163, 165, 167, and 169). In one example, the gRNA includes at least one DR sequence having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 129, 130, 131, 132, 133, 134, 135, 136, 137, 148, 150, 151, 152, 154, 156, 157, 159, 161, 163, 165, 167, 169, 176, 178, 180, 182, 184, 186, 188, 190, 191, 192, 193, 199, 201, 203, 205, 207, 209, 211, 213, 215, 217, 219, 221, 223, 225, 227, 228, 230, 232, 234, 236, 238, 240, 242, 244, 246, 248, 250, 252, or 254 (such as 1, 2, 3, or 4 of such DR sequences).

In one example, a guide sequence includes a constant DR on its 5'-end and a variable spacer on its 3' end. In one example includes the sequence DR-spacer-DR-spacer. In some examples, the sequence DR-spacer is repeated two or more times, such as at least 3 times or at least 4 times. This type of sequence is called a guide array.

Guide molecules generally exist in various states of processing. In one example, an unprocessed guide RNA is 36nt of DR followed by 30-32 nt of spacer. The guide RNA is processed (truncated/modified) by Cas13d itself or other RNases into the shorter "mature" form. In some embodiments, an unprocessed guide sequence is about, or at least about 30, 35, 40, 45, 50, 55, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, or more nucleotides (nt) in length. In some embodiments, a processed guide sequence is about 44 to 60 nt (such as 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64,

65, 66, 67, 68, 69, or 70 nt). In some embodiments, an unprocessed spacer is about 28-32 nt long (such as 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 nt) while the mature (processed) spacer can be about 10 to 30 nt, 10 to 25 nt, 14 to 25 nt, 20 to 22 nt, or 14-30 nt (such as 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 nt). In some embodiments, an unprocessed DR is about 36 nt (such as 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40 or 41 nt), while the processed DR is about 30 nt (such as 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 nt).

The ability of a guide sequence to direct sequence-specific binding of a CRISPR complex to a target RNA may be assessed by any suitable assay. For example, the components of a CRISPR system sufficient to form a CRISPR complex, including the guide sequence to be tested, may be provided to a host cell having the corresponding target RNA molecule, such as by transfection with vectors encoding the components of the CRISPR sequence, followed by an assessment of preferential cleavage within the target sequence. Similarly, cleavage of a target RNA sequence may be evaluated in a test tube by providing the target RNA, components of a CRISPR complex, including the guide sequence to be tested and a control guide sequence different from the test guide sequence, and comparing binding or rate of cleavage at the target RNA between the test and control guide sequence reactions. Other assays are possible, and will occur to those skilled in the art.

Also provided are vectors, such as a viral vector or plasmid (e.g., retrovirus, lentivirus, adenovirus, adeno-associated virus, or herpes simplex virus), that includes a guide nucleic acid molecule. Exemplary vectors are described herein. In some examples, the guide nucleic acid molecule is operably linked to a promoter or expression control element (examples of which are provided elsewhere in this application). As described elsewhere herein, such vectors can include other elements, such as a gene encoding a selectable marker, such as an antibiotic, such as puromycin, hygromycin, or a detectable marker such as GFP or other fluorophore.

In one example, a plurality of gRNAs are part of an array (which can be part of a vector, such as a viral vector or plasmid). For example, a guide array including the sequence DR-spacer-DR-spacer-DR-spacer, can include three unique unprocessed gRNAs (one for each DR-spacer sequence). Once introduced into a cell or cell-free system, the array is processed by the Cas13d protein into the three individual mature gRNAs. This allows for multiplexing, e.g. the delivery of multiple gRNAs to a cell or system to target multiple target RNAs or multiple positions within a single target RNA (or combinations thereof).

D. Vectors that encode Cas13d and Guide Nucleic Acid Molecules

The disclosure provides vectors, such as plasmids and viral vectors as described elsewhere herein, which include one or more guide molecule coding sequences (e.g., to permit targeting of one or more RNA molecules), and one or more Cas13d protein coding sequences. Such vectors can be used in the methods, compositions, and kits provided herein. Such vectors can include naturally occurring or non-naturally occurring nucleotides or ribonucleotides. Such vectors can include a single promoter operably linked to the guide molecule (which can be part of an array that includes at least two different guide molecules) and the Cas13d protein coding sequence. Alternatively, the guide molecule (which can be part of an array that includes at least two different guide molecules) and the Cas13d protein coding sequence can be operably linked to different promoters.

E. Recombinant Cells and Cell-Free Systems

Cells that include a non-native Cas13d protein, a non-native Cas13d protein coding sequence, a guide molecule (or coding sequence), or combinations thereof, are provided. Such recombinant cells can be used in the methods, compositions, and kits provided herein. Nucleic acid molecules encoding an Cas13d protein disclosed herein and/or nucleic acid molecules encoding a guide molecule can be introduced into cells to generate transformed (e.g., recombinant) cells. In some examples, such cells are generated by introducing one or more non-native Cas13d proteins and one or more guide molecules (e.g., gRNAs) into the cell, for example as a ribonucleoprotein (RNP) complex.

Similarly, cell free systems, such as those generated from lysed cells (or those that include an Cas13d RNP in a test tube or other vessel, into which in vitro transcribed or chemically synthesized target RNAs are added), which include a, Cas13d protein, a Cas13d protein coding sequence, a guide molecule (or coding sequence), or combinations thereof, are provided. Such cell free systems can be used in the methods, compositions, and kits provided herein. In some examples, one or more non-native Cas13d proteins and one or more guide molecules (e.g., gRNAs) are added to a cell free system, for example as a RNP complex.

Thus, cells and cell-free systems containing an Cas13d protein (such as a protein having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 138, 147, 149, 153, 155, 158, 160, 162, 164, 166, 168, 170, 175, 177, 179, 181, 183, 185, 187, 189, 194, 198, 200, 202, 204, 206, 208, 210, 212, 214, 216, 218, 220, 222, 224, 226, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, or 253) are disclosed. Similarly, cells and cell-free systems containing a guide molecule, such as one having at least one DR sequence having at least 80%, at least 85%, at least 90%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 129, 130, 131, 132, 133, 134, 135, 136, or 137, and in some examples also at least one spacer sequence complementary to a target RNA, are provided.

Such recombinant cells (e.g., which can be used to generate a cell-free system) can be eukaryotic or prokaryotic. Examples of such cells include, but are not limited to bacteria, archaea, plant, fungal, yeast, insect, and mammalian cells, such as *Lactobacillus*, *Lactococcus*, *Bacillus* (such as *B. subtilis*), *Escherichia* (such as *E. coli*), *Clostridium*, *Saccharomyces* or *Pichia* (such as *S. cerevisiae* or *P. pastoris*), *Kluyveromyces lactis*, *Salmonella typhimurium*, *Drosophila* cells, *C. elegans* cells, *Xenopus* cells, SF9 cells, C129 cells, *Neurospora*, and immortalized mammalian cell lines (e.g., HeLa cells, myeloid cell lines, and lymphoid cell lines).

In one example, the cell is a prokaryotic cell, such as a bacterial cell, such as *E. coli*.

In one example, the cell is a eukaryotic cell, such as a mammalian cell, such as a human cell. In one example, the cell is primary eukaryotic cell, a stem cell, a tumor/cancer cell, a circulating tumor cell (CTC), a blood cell (e.g., T cell, B cell, NK cell, Tregs, etc.), hematopoietic stem cell,

specialized immune cell (e.g., tumor-infiltrating lymphocyte or tumor-suppressed lymphocytes), a stromal cell in the tumor microenvironment (e.g., cancer-associated fibroblasts, etc.) In one example, the cell is a brain cell (e.g., neurons, astrocytes, microglia, retinal ganglion cells, rods/cones, etc.) of the central or peripheral nervous system).

In one example, a cell is part of (or obtained from) a biological sample, such as a biological specimen containing genomic DNA, RNA (e.g., mRNA), protein, or combinations thereof, obtained from a subject. Examples include, but are not limited to, peripheral blood, serum, plasma, urine, saliva, sputum, tissue biopsy, fine needle aspirate, surgical specimen, and autopsy material. Such cells can also be used to generate a cell free system.

In one example the cell (or cell free system) is from a tumor, such as a hematological tumor (e.g., leukemias, including acute leukemias (such as acute lymphocytic leukemia, acute myelocytic leukemia, acute myelogenous leukemia and myeloblastic, promyelocytic, myelomonocytic, monocytic and erythroleukemia), chronic leukemias (such as chronic myelocytic (granulocytic) leukemia, chronic myelogenous leukemia, and chronic lymphocytic leukemia), polycythemia vera, lymphoma, Hodgkin's disease, non-Hodgkin's lymphoma (including low-, intermediate-, and high-grade), multiple myeloma, Waldenström's macroglobulinemia, heavy chain disease, myelodysplastic syndrome, mantle cell lymphoma and myelodysplasia) or solid tumor (e.g., sarcomas and carcinomas: fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, and other sarcomas, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, lymphoid malignancy, pancreatic cancer, breast cancer, lung cancers, ovarian cancer, prostate cancer, hepatocellular carcinoma, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, Wilms' tumor, cervical cancer, testicular tumor, bladder carcinoma, and CNS tumors (such as a glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma and retinoblastoma).

In one example the cell (or cell free system) is obtained from an environmental sample, such as a water, soil, or air sample.

F. Compositions & Kits

Compositions and kits that include a Cas13d protein, a Cas13d protein coding sequence, a guide molecule (or coding sequence), or combinations thereof, are provided. In one example, the composition or kit includes an RNP complex composed of one or more Cas13d proteins and one or more guide molecules (e.g., gRNAs). In one example, the composition or kit includes a vector encoding an Cas13d protein, a guide molecule, or both. In one example, the composition or kit includes a cell, such as a bacterial cell or eukaryotic cell, that includes a non-native Cas13d protein, a non-native Cas13d protein coding sequence, a guide molecule (or coding sequence), or combinations thereof. In one example, the composition or kit includes a cell-free system that includes a non-native Cas13d protein, a non-native Cas13d protein coding sequence, a guide molecule (or coding sequence), or combinations thereof.

Such compositions can include a pharmaceutically acceptable carrier (e.g., saline, water, PBS). In some examples, the composition is a liquid, lyophilized powder, or cryopreserved.

In some examples, the kit includes a delivery system (e.g., liposome, a particle, an exosome, a microvesicle, a viral vector, or a plasmid), and/or a label (e.g., a peptide or antibody that can be conjugated either directly to an Cas13d RNP or to a particle containing the Cas1 RNP to direct cell type specific uptake/enhance endosomal escape/enable blood-brain barrier crossing etc.). In some examples, the kits further include cell culture or growth media, such as media appropriate for growing bacterial, plant, insect, or mammalian cells.

In some examples, such parts of a kit are in separate containers.

G. Targeting RNA

The disclosed Cas13d proteins (and coding sequences), and guide molecules (e.g., gRNA and coding sequences) can be used in a CRISPR/Cas system to target one or more RNA molecules, such as those present in a sample (such as a biological sample, environmental sample (e.g., soil, air or water sample), and the like. In one example, the target RNA is a coding RNA. In one example, the target RNA is a nuclear RNA. In other examples, the target RNA is non-coding RNA (such as functional RNA, siRNA, microRNA, snRNA, snoRNA, piRNA, scaRNA, tRNA, rRNA, lincRNA, or lincRNA). Such RNA targeting methods can be performed in vitro (such as in cell culture or in a cell-free system), or in vivo (such as in an organism, embryo, or mammal)

The CRISPR/Cas system provided herein includes two general components: (1) an Cas13d protein or its coding sequence (whose expression can be driven by a promoter) and (2) a guide nucleic acid molecule, such as RNA (gRNA), which is specific for the target RNA (whose expression can also be driven by promoter. When introduced into cells (or to a cell free system) for example (1) as Cas13d mRNA and Cas13d gRNA, (2) as part of a single vector or plasmid or divided into multiple vectors or plasmids, (3) as separate Cas13d protein and guide molecules, or (4) as an RNP complex of the Cas13d protein and guide molecule, the guide molecule guides the Cas13d to the target RNA. If the Cas13d protein has a native HEPN domain(s) or is fused to an appropriate effector domain bearing RNase activity, the RNA can be cut. If the Cas13d protein has a mutated HEPN domain(s), a guide array can be processed into mature gRNAs, but the target RNA is not cut. Using this system, RNA sequences are easily targeted, for example edited or detected, optionally with an effector domain.

1. Introduction of Cas13d Protein Directly into a Cell

In one example, the Cas13d protein is expressed in a recombinant cell, such as *E. coli*, and purified. The resulting purified Cas13d protein, along with an appropriate guide molecule specific for the target RNA, is then introduced into a cell or organism where one or more RNAs can be targeted. In some examples, the Cas13d protein and guide nucleic acid molecule are introduced as separate components into the target cell/organism. In other examples, the purified Cas13d protein is complexed with the guide nucleic acid (e.g., gRNA), and this ribonucleoprotein (RNP) complex is introduced into target cells (e.g., using transfection or injection). In some examples, the Cas13d protein and guide molecule are injected into an embryo (such as a human, mouse, zebrafish, or *Xenopus* embryo).

Once the Cas13d protein and guide nucleic acid molecule are in the cell, one or more RNAs can be targeted.

2. Expression of Cas13d from Nucleic Acids

In one example, the Cas13d protein is expressed from a nucleic acid molecule in a cell containing a target RNA, for example an RNA to be detected or modified. In some such examples, the Cas13d protein is expressed from a vector, such as a viral vector or plasmid introduced into a cell or into a cell-free system. This results in the production of the Cas13d protein in the cell, organism, or system. In addition, these nucleic acid molecules can be co-expressed in the cell/organism/system with the guide nucleic acid molecule (e.g., gRNA) specific for the target RNA.

In one example, multiple plasmids or vectors are used for RNA targeting. The nucleic acid molecule encoding the Cas13d can be provided for example on one vector or plasmid, and the guide nucleic acid molecule (e.g., gRNA) on another plasmid or vector. Multiple plasmids or viral vectors can be mixed and introduced into cells (or a cell free system) at the same time, or separately.

In some examples, multiple nucleic acid molecules are expressed from a single vector or plasmid. For example, a single vector can include the nucleic acid molecule encoding the Cas13d, and a separate vector can include the guide molecule.

In some examples a plurality of different guide molecules (e.g., gRNAs), one for each target (such as 1, 2, 3, 4, 5, or 10 different targets), are present on a single array and/or vector. In one example, the method includes delivering a plurality of gRNAs (such as at least 2, at least 3, at least 4, at least 5, at least 10, at least 20, or at least 50 different gRNAs), which are part of an array (which can be part of a vector, such as a viral vector or plasmid). Once introduced into a cell or cell-free system, the array is processed by the Cas13d protein into the individual mature gRNAs.

The nucleic acid molecules expressed from the vector can be under the control of a promoter and optionally contain selection markers (such as antibiotic resistance).

In some examples, the protein and guide molecules are expressed by an embryo (such as a zebrafish or *Xenopus* embryo). The Cas13d protein can be expressed from injected plasmid DNA, injected mRNA, or stably integrated copies into the animal genome. The gRNA can be directly injected or expressed from a vector or stably integrated copies into the animal genome.

3. Targets

One or more RNAs can be targeted by the disclosed methods, such as at least 1, at least 2, at least 3, at least 4 or at least 5 different RNAs in a cell, cell-free system, or organism, such as 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 different RNAs. In one example, the RNA is associated with a disease such as cystic fibrosis, Huntington's disease, Tay-Sachs, Fragile X syndrome, Fragile X-associated tremor/ataxia syndrome, Duchenne muscular dystrophy, myotonic dystrophy, spinal muscular atrophy, spinocerebellar ataxia, or familial ALS. In one example, the RNA is associated with cancer (e.g., a cancer of the lung, breast, colon, liver, pancreas, prostate, bone, brain, skin (e.g., melanoma), or kidney). Examples of target RNAs include, but are not limited to those associated with cancer (e.g., BCR-ABL, Ras, Raf, p53, BRCA1, BRCA2, CXCR4, beta-catenin, HER2, and CDK4).

In one example, the RNA is associated with viral infection, such as infection by a positive-strand RNA viruses, such as Picornaviruses (such as Aphthoviridae [for example foot-and-mouth-disease virus (FMDV)]), Cardioviridae; Enteroviridae (such as Coxsackie viruses, Echoviruses, Enteroviruses, and Polioviruses); Rhinoviridae (Rhinoviruses); Hepataviridae (Hepatitis A viruses); Togaviruses

(examples of which include rubella; alphaviruses (such as Western equine encephalitis virus, Eastern equine encephalitis virus, and Venezuelan equine encephalitis virus)); Flaviviruses (examples of which include Dengue virus, West Nile virus, and Japanese encephalitis virus); Calciviridae (which includes Norovirus and Sapovirus); or Coronaviruses (examples of which include SARS coronaviruses, such as the Urbani strain), or a negative-strand RNA virus, such as Orthomyxoviruses (such as the influenza virus), Rhabdoviruses (such as Rabies virus), and Paramyxoviruses (examples of which include measles virus, respiratory syncytial virus, or parainfluenza viruses), or a DNA viral infection (such as infection by Herpesviruses (such as Varicella-zoster virus, for example the Oka strain; cytomegalovirus; and Herpes simplex virus (HSV) types 1 and 2), Adenoviruses (such as Adenovirus type 1 and Adenovirus type 41), Poxviruses (such as Vaccinia virus), or Parvoviruses (such as Parvovirus B19).

In one example, the RNA is associated with a bacterial infection or property of a bacterial infection, such as bacterial resistance, persistence, or antibiotic resistance. Detection of these RNAs can be used for diagnostic methods, while editing these RNAs in cell-based or cell-free systems can be used for therapeutic methods.

4. Methods of Detecting RNA

In one example, the method of targeting an RNA results in detecting, visualizing, or labeling a target RNA. For example, by using at least one Cas13d protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4), at least one gRNA containing at least one spacer sequence specific for the target RNA, and an effector module, the target RNA will be recognized by Cas13d but will not be cut or nicked while the effector module becomes activated. In some examples, such a method is used to detect a target RNA. Such a method can be used in a cell or cell free system to determine if a target RNA is present, such as in a tumor cell. In some examples, the cell or cell free system is obtained from a tissue sample, blood sample or saliva sample.

In one example, the method of detecting RNA comprises of an Cas13d protein fused to a fluorescent protein or other detectable label along with a gRNA containing a spacer sequence specific for the target RNA. Binding of Cas13d to the target RNA can be visualized by microscopy or other methods of imaging. In another example, RNA aptamer sequences can be appended to or inserted within the gRNA molecule, such as MS2, PP7, Q β , and other aptamers. The introduction of proteins that specifically bind to these aptamers, e.g. the MS2 phage coat protein, fused to a fluorescent protein or other detectable label can be used to detect the target RNA because the Cas13d-gRNA-target RNA complex will be labeled via the aptamer interaction.

In another example, the method of detecting RNA is a transcriptional sensor (e.g., as part of a synthetic circuit) for diagnostics or therapeutics. For example, the transcriptional sensor can be comprised of at least one Cas13d protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4), at least one gRNA containing at least one spacer sequence specific for the target RNA, and an effector module such as an optionally split fluorescent protein or probe; an optionally split luminescent protein or probe; an optionally split enzyme that catalyzes a detectable reaction such as ubiquitin or TEV protease; a FRET-compatible protein pair; one or more transcription factor(s) fused to Cas13d via cleavable linkers; a split intein that trans-splices a protein to restore its function such as a transcription factor; a kinase-substrate pair that activates upon phosphorylation; one, two, or more monomers that activate upon dimerization or multimeriza-

tion; or one or more proteins that induce conformational and functional change upon interaction. In one example, the spatial proximity of two or more Cas13d proteins and gRNAs due to binding a particular transcript would activate the effector module, resulting in a detectable signal or detectable activity in the cell.

For example, the transcriptional sensor could allow a cancer-specific transcript, inflammation-specific transcript, disease-specific transcript, or cell state-specific transcript to be detected. A synthetic circuit containing a Cas13d-based system that is able to sense particular transcripts could encode conditional logic, e.g. requiring target detection to up- or down-regulate a gene for therapeutic application.

In one example, the method results in a detectable agent being bound to the target RNA, which can be detected. For example, two separate Cas13d fusion proteins that each include part of a fluorophore (e.g., GFP), and two different gRNAs with different spacer sequences that target regions of an RNA in close proximity, can be used. When the two parts bind to the target RNA in proximity, the two parts of the fluorophore form a complete fluorophore, thereby generating a detectable signal.

In one example, the method results in RNA detection, for example by triggering a response such as expression of a second gene, modification of a protein, translocation of a protein or RNA to a different location, induction of cell death via suicide gene, induction of cell proliferation, induction of a transgene that enables a secondary function, induction of a permanent change in DNA sequence to enable storing a memory of past transcriptional events, or altering the RNA to enable pulldown.

In one embodiment, two halves of a transcription factor could be linked to two separate Cas13d via a split intein system. The Cas13d proteins are provided with two different gRNAs with different spacer sequences that target regions of an RNA in close proximity. Upon binding to the target RNA in proximity, the split inteins trans-splice a reconstituted transcription factor (TF) so that it can translocate to the nucleus and turn on a target gene or cluster of target genes. In one example, the target gene could be an endogenous gene in the cell. In another example, the target gene could be a transgene expressed on a vector or introduced through genetic engineering, such as a fluorescent protein or toxin.

5. Methods of Detecting RNA in Cell-Free Systems

In one example, the method of detecting a target RNA in a cell-free system results in a detectable label or enzyme activity. For example, by using at least one Cas13d protein (e.g., SEQ ID NO: 3, 42, 62, 70, 82, 83, and 92), at least one gRNA containing at least one spacer sequence specific for the target RNA, and a detectable label, the target RNA will be recognized by Cas13d. The binding of the target RNA by Cas13d triggers its RNase activity, which can lead to the cleavage of the target RNA as well as the detectable label.

In one example, the detectable label is an RNA linked to a fluorescent probe and quencher. The intact detectable RNA links the fluorescent probe and quencher, suppressing fluorescence. Upon cleavage by Cas13d of the detectable RNA, the fluorescent probe is released from the quencher and displays fluorescent activity. Such a method can be used to determine if a target RNA is present in a lysed cell sample, lysed tissue sample, blood sample, saliva sample, environmental sample (such as a water, soil, or air sample), or other lysed cell or cell-free sample. Such a method can also be used to detect a pathogen, such as a virus or bacteria, or diagnose a disease state, such as a cancer.

In one example, the detection of the target RNA aids in the diagnosis of disease and/or pathological state, or the exist-

tence of a viral or bacterial infection. For example, Cas13d-mediated detection of non-coding RNAs such as PCA3 can be used to diagnose prostate cancer if detected in patient urine. In another example, Cas13d-mediated detection of the lncRNA-AA174084, which is a biomarker of gastric cancer, can be used to diagnose gastric cancer.

6. Methods of Editing Target RNA

In one example, the method of targeting an RNA results in editing the sequence of a target RNA. For example, by using an Cas13d protein with a non-mutated HEPN domain (e.g., SEQ ID NOS: 1, 3, 42, 62, 70, 82, 83, and 92), and a gRNA containing at least one a spacer sequence specific for the target RNA, the target RNA can be cut or nicked at a precise location. In some examples, such a method is used to decrease expression of a target RNA, which will decrease translation of the corresponding protein. Such a method can be used in a cell where increased expression of an RNA is not desired. In one example, the RNA is associated with a disease such as cystic fibrosis, Huntington's disease, Tay-Sachs, Fragile X syndrome, Fragile X-associated tremor/ataxia syndrome, muscular dystrophy, myotonic dystrophy, spinal muscular atrophy, spinocerebellar ataxia, or familial ALS. In another example, the RNA is associated with cancer (e.g., a cancer of the lung, breast, colon, liver, pancreas, prostate, bone, brain, skin (e.g., melanoma), or kidney). Examples of target RNAs include, but are not limited to those associated with cancer (e.g., PD-L1, BCR-ABL, Ras, Raf, p53, BRCA1, BRCA2, CXCR4, beta-catenin, HER2, and CDK4). Editing such target RNAs can have a therapeutic effect.

In another example, the RNA is expressed in an immune cell. The target RNA could, for example, code for a protein leading to the repression of a desirable immune response, such as infiltration of a tumor. Knock-down of such an RNA could enable progression of such a desirable immune response (e.g., PD1, CTLA4, LAG3, TIM3). In another example, the target RNA encodes a protein resulting in the undesirable activation of an immune response, for example in the context of an autoimmune disease such as multiple sclerosis, Crohn's disease, lupus, or rheumatoid arthritis.

In one example, targeting the target RNA allows for decreasing expression of the target protein encoded by the RNA. For example, by using an Cas13d fusion protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4) and a translational repression domain (such as Pumilio or FBF PUF proteins, deadenylases, CAF1, Argonaute proteins, and others), and a guide RNA containing at least one spacer sequence specific for the target RNA, expression of a target RNA can be decreased.

In some examples, Cas13d can be fused to a ribonuclease (such as a PIN endonuclease domain, an NYN domain, an SMR domain from SOT1, or an RNase domain from Staphylococcal nuclease) or a domain that affects RNA stability (such as tristetraprolin or domains from UPF1, EXOSC5, and STAU1).

In another example, RNA aptamer sequences can be appended to or inserted within the gRNA molecule, such as MS2, PP7, Q β , and other aptamers. Proteins that specifically bind to these aptamers, e.g. the MS2 phage coat protein, can be fused to a translational repression domain, a ribonuclease, or a domain that affects RNA stability. This aptamer-effector domain fusion can be used to target the target RNA because the Cas13d and gRNA complex will guide the aptamer protein-effector domain in proximity to the target RNA.

Such a method can be used in a cell where increased expression of an RNA is not desired, such as when an expressed RNA is associated with a disease such as cystic

fibrosis, Huntington's disease, Tay-Sachs, Fragile X syndrome, Fragile X-associated tremor/ataxia syndrome, muscular dystrophy, myotonic dystrophy, spinal muscular atrophy, spinocerebellar ataxia, or familial ALS. In another example, the target RNA is associated with cancer (e.g., a cancer of the lung, breast, colon, liver, pancreas, prostate, bone, brain, skin (e.g., melanoma), or kidney). Examples of target RNAs include, but are not limited to those associated with cancer (e.g., PD-L1, BCR-ABL, Ras, Raf, p53, BRCA1, BRCA2, CXCR4, beta-catenin, HER2, and CDK4). Editing such target RNAs would have a therapeutic effect.

In another example, the RNA is expressed in an immune cell. The target RNA could, for example, code for a protein leading to the repression of a desirable immune response, such as infiltration of a tumor. Knock-down of such an RNA could enable progression of such a desirable immune response (e.g., PD1, CTLA4, LAG3, TIM3). In another example, the target RNA could encode a protein resulting in the undesirable activation of an immune response, for example in the context of an autoimmune disease such as multiple sclerosis, Crohn's disease, lupus, or rheumatoid arthritis.

In one example, targeting the target RNA allows for activating or increasing expression of the target RNA. For example, by using an Cas13d fusion protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4) and a translational activation domain (such as eIF4E and other translation initiation factors, a domain of the yeast poly(A)-binding protein or GLD2), and a guide RNA containing at least one a spacer sequence specific for the target RNA, expression of a target RNA can be increased. Aptamer introduction into the gRNA with a cognate aptamer-binding protein fused to a translational activation domain can also be used. In one example, RNA aptamer sequences are appended to or inserted within the gRNA molecule, such as MS2, PP7, Q β , and other aptamers. The introduction of proteins that specifically bind to these aptamers, e.g. the MS2 phage coat protein, fused to a translational activation domain can be used to target the target RNA because the Cas13d and gRNA complex will bring the aptamer protein-translational activation domain in proximity to the target RNA.

In some examples, such a method is used to increase the activity or expression of a target RNA, which will increase translation of the corresponding protein (if the RNA is a coding RNA). Such a method can be used in a cell where increased expression of an RNA is desired, such as a heterozygous genetic disease or disorders caused by copy number variation. Increasing translation of a desired protein product could be therapeutic in nature.

In another example, increasing the expression of a target RNA (such as Cyclin B1) can render the target cell (such as cancers) more sensitive to drugs (such as chemotherapeutic agents).

In one example, targeting the target RNA allows for one or more RNA base substitutions, RNA base edits, RNA base deletions, RNA base insertions, or combinations thereof, in the target RNA. In some examples, the Cas13d protein with a mutated HEPN domain is associated, either via direct fusion or a gRNA-aptamer modification, an effector domain that allows base edits (such as a cytidine deaminase, PPR protein, adenosine deaminase, ADAR family protein, or APOBEC family protein). In some examples, such a method is used to modify an RNA sequence, edit an RNA mutation, or modify an RNA transcript (e.g., gene therapy), for

example to treat diseases such as ALS and melanoma or genetic disorders caused by undesired splice sites, such as Leber congenital amaurosis.

In one example, targeting the target RNA allows for methylating the target RNA. Some examples may use a chimeric Cas13d protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4) associated either via direct fusion or a gRNA-aptamer modification with a methylation domain (e.g., m6A), and a guide RNA containing at least one a spacer sequence specific for the target RNA. In some examples, such a method is used to combat aberrant RNA demethylation. In one example, such a method is used to modify the methylation levels of pluripotency transcripts such as NANOG or KLF4 for example to decrease their stability in breast cancer cells, which can suppress the acquisition of breast cancer stem cell phenotypes that are associated with increased proliferation and cancer stem cell formation.

In one example, targeting the target RNA allows for demethylating the target RNA. Some examples can use Cas13d protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4), a guide RNA containing at least one a spacer sequence specific for the target RNA, and a demethylation domain (e.g., human alkylation repair homolog 5 or Alkbh5). The demethylation domain can be associated either via direct fusion to the Cas13d protein or via a gRNA-aptamer modification. In some examples, such a method is used to reverse aberrant RNA methylation, for example to treat myeloid leukemia by decreasing m⁶A levels.

In one example, targeting the target RNA allows for binding to the target RNA. For example, by using a Cas13d protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4) and a guide RNA containing at least one a spacer sequence specific for the target RNA, molecules can be bound or tethered to a target RNA. In some examples, such a method is used to capture the target RNA (e.g., immunoprecipitation). This can be used as part of a kit to identify the proteins interacting with a specific RNA transcript. In one example, an epitope tagged Cas13d (e.g. FLAG, HA, biotin, HALO tag) can be targeted to specific target RNAs and cross-linked via fixation (e.g. with paraformaldehyde or glutaraldehyde). Immunoprecipitation of Cas13d with an epitope-recognizing antibody allows for the identification of co-immunoprecipitated proteins via Western blot or mass spectrometry.

In another example, Cas13d can be fused to a biotin ligase (such as BirA) or a peroxidase (such as APEX2) in order to biotinylate proteins that interact with the target RNA. Labeled proteins can then be pulled down with streptavidin beads followed by mass spectrometry or Western blot.

In some examples, biotinylated Cas13d could be targeted to ribosomal RNA sequences with a gRNA. Streptavidin bead-mediated pulldown can be used to deplete rRNA for RNA sequencing library preparation.

In one example, targeting the target RNA allows for masking the target RNA. For example, by using an Cas13d protein with a mutated or intact HEPN domain and a guide RNA containing at least one a spacer sequence specific for the target RNA, a target RNA can be masked from RNA-binding proteins or RNA-binding elements such as miRNAs.

In some examples, the Cas13d can be used to mask RNA binding sites from RNA-binding proteins (RBPs). In another example, Cas13d can mask miRNA binding sites. For example, the liver-specific miR-122 forms a complex with Hepatitis C viral RNA which protects it from degradation. A HEPN-active Cas13d protein could be targeted to the

miRNA-122 binding site on the viral RNA to synergistically combat HCV infections by simultaneously reversing miRNA-122-mediated protection and directly degrading HCV RNA. In some examples, such a method is used to preserve or protect the target RNA molecule, for example to protect the target RNA from degradation. For example, by targeting AU-rich elements in the 3' UTR of a target gene, a HEPN-mutated Cas13d can block binding of RNA-binding proteins such as tristetraprolin (TTR) or AUF1, which lead to degradation of the target transcript.

In one example, targeting the target RNA allows for changing splicing of the target RNA. Both the direct binding of splice acceptor and/or donor sites as well as splice effector domains can be used to manipulate splicing. For example, by using an Cas13d protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4), a guide RNA containing at least one a spacer sequence specific for the target RNA, and optionally an effector domain that affects splicing (such as the RS-rich domain of SRSF1, the Gly-rich domain of hnRNP A1, the alanine-rich motif of RBM4, or the proline-rich motif of DAZAP1), alternative splicing of the RNA can be achieved.

In some examples, such a method is used for exon inclusion, for example to include exon 2 of acid alpha-glucosidase (GAA) to treat Pompe disease or to include exon 7 of SMN2 to treat spinal muscular atrophy (SMA). In some examples, such a method is used for exon exclusion, for example to restore the reading frame of dystrophin to treat Duchenne muscular dystrophy or to shift the splicing of the Bcl-x pre-mRNA from the antiapoptotic long isoform to the proapoptotic short isoform to treat cancer.

In some examples, the method uses the Cas13d protein with a mutated HEPN domain to mask splice acceptor or donor sites, for example to create neoantigens to make cold tumors hot. By affecting the splicing of certain target pre-mRNAs, this method can generate novel exon-exon junctions that can lead to the creation of neo-epitopes in cancer cells. This can make a cancer cell vulnerable to the immune system due to the display of unnatural antigens. In other examples, this method can be used to dynamically manipulate isoform ratios or to restore reading frame of a protein (e.g., dystrophin for Duchenne's muscular dystrophy).

In one example, targeting the target RNA allows for controlling transcript trafficking of the target RNA. For example, by using an Cas13d fusion protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4) and a subcellular localization signal or export sequence, a guide RNA containing at least one a spacer sequence specific for the target RNA. In some examples, such a method is used to traffic the target RNA molecule to a particular organelle or cytosolic compartment, or even export the target RNA transcript, for example to endosomes for extracellular release

In another example, the method can affect RNA trafficking. For example, the zipcode binding protein ZBP1 specifically recognizes an RNA sequence 5'-CGGAC(C/A-CA-C/U) that leads to localization of certain transcripts to the leading edge of fibroblasts. By masking or manipulating particular RNA zipcodes or regulatory sequences from recognition by regulatory protein complexes, this method can affect RNA localization or trafficking within a cell.

In one example, the target RNA is a nuclearly localized RNA. For example, by using an Cas13d protein with a non-mutated HEPN domain (e.g., SEQ ID NOS: 1, 3, 42, 62, 70, 82, 83, and 92) fused to a nuclear localization signal and a guide RNA containing at least one a spacer sequence specific for the target nuclear RNA, the nuclearly localized RNA can be targeted and degraded. In some examples, such

a method is used to degrade the target nuclear RNA molecule, for example to knock-down a non-coding nuclear RNA such as HOTAIR, which is associated with metastatic progression in breast cancer.

In one example, the target RNA is viral RNA or transcript of a DNA virus. For example, an Cas13d protein with a non-mutated HEPN domain (e.g., SEQ ID NOS: 1, 3, 42, 62, 70, 82, 83, and 92) and a guide RNA containing at least one spacer sequence specific for the target RNA can be used. In some examples, such a method is used to treat an RNA viral infection (such as infection by a positive-strand RNA viruses, such as Picornaviruses (such as Aphthoviridae [for example foot-and-mouth-disease virus (FMDV)], Cardioviridae; Enteroviridae (such as Coxsackie viruses, Echoviruses, Enteroviruses, and Polioviruses); Rhinoviridae (Rhinoviruses)); Hepataviridae (Hepatitis A viruses); Togaviruses (examples of which include rubella; alphaviruses (such as Western equine encephalitis virus, Eastern equine encephalitis virus, and Venezuelan equine encephalitis virus)); Flaviviruses (examples of which include Dengue virus, West Nile virus, and Japanese encephalitis virus); Calciviridae (which includes Norovirus and Sapovirus); or Coronaviruses (examples of which include SARS coronaviruses, such as the Urbani strain), or a negative-strand RNA virus, such as Orthomyxoviruses (such as the influenza virus), Rhabdoviruses (such as Rabies virus), and Paramyxoviruses (examples of which include measles virus, respiratory syncytial virus, or parainfluenza viruses), or a DNA viral infection (such as infection by Herpesviruses (such as Varicella-zoster virus, for example the Oka strain; cytomegalovirus; and Herpes simplex virus (HSV) types 1 and 2), Adenoviruses (such as Adenovirus type 1 and Adenovirus type 41), Poxviruses (such as Vaccinia virus), or Parvoviruses (such as Parvovirus B19), for example by cutting the viral RNA or transcript of a DNA virus. Thus, such methods can be used as an RNA-based antiviral or antimicrobial.

EXAMPLE 1

Materials and Methods

This example describes the materials and methods used to obtain the results shown in Examples 2-7.
Cell Culture of Human Embryonic Kidney (HEK) Cell Line 293FT

Human embryonic kidney (HEK) cell line 293FT (Thermo Fisher) was maintained in DMEM (4.5 g/L glucose), supplemented with 10% FBS (GE Life Sciences) and 10 mM HEPES at 37° C. with 5% CO₂. Upon reaching 80-90% confluency, cells were dissociated using TrypLE Express (Life Technologies) and passaged at a ratio of 1:2. Cell Culture of Human Bone Osteosarcoma Epithelial Cell Line U2OS

Human bone osteosarcoma epithelial U2OS were maintained in DMEM (4.5 g/L glucose) supplemented with 10% FBS and 10 mM HEPES at 37° C. with 5% CO₂. Cells were passaged at a 1:3 ratio upon reaching 70% confluence. This cell line was not authenticated.

Maintenance of induced pluripotent stem cells and neuronal differentiation

Stable human iPSC lines containing the FTDP-17 IVS10+16 mutation or age- and sex-matched control lines were obtained from the laboratory of Fen-Biao Gao (Biswas et al., 2016). Briefly, cells obtained from one male patient with the MAPT IVS10 +16 mutation and two separate lines from one male control patient were reprogrammed into hiPSCs

(Almeida et al., 2012). iPSCs were transduced with lentivirus containing a doxycycline-inducible Ngn2 cassette. Lentiviral plasmids were a gift from S. Schafer and F. Gage. iPSCs were then passaged with Accutase and plated into a Matrigel-coated 6-well plate with mTESR media containing ROCK inhibitor Y-27632 (10 μM, Cayman) at 500,000 cells per well. On day 1, media was changed with mTESR. On day 2, media was changed to mTESR containing doxycycline (2 μg/ml, Sigma) to induce Ngn2 expression. On day 3, culture media was replaced with Neural Induction media (NIM, DMEM/F12 (Life Technologies) containing BSA (0.1 mg/ml, Sigma), apo-transferrin (0.1 mg/ml, Sigma), putrescine (16 μg/ml, Sigma), progesterone (0.0625 μg/ml, Sigma), sodium selenite (0.0104 μg/ml, Sigma), insulin (5 μg/ml, Roche), BDNF (10 ng/ml, Peprotech), SB431542 (10 μM, Cayman), LDN-193189 (0.1 μM, Sigma), laminin (2 μg/ml, Life Technologies), doxycycline (2 μg/ml, Sigma) and puromycin (Life Technologies)). NIM media was changed daily. Following 3 days of puromycin selection, immature neuronal cells were passaged with Accumax (Innovative Cell Technologies) and plated onto 96-well plates coated with poly-D-lysine and Matrigel in Neural Maturation media (NMM; 1:1 Neurobasal/DMEM (Life Technologies) containing B27 (Life Technologies), BDNF (10 ng/ml, Peprotech), N-Acetyl cysteine (Sigma), laminin (2 μg/ml, Life Technologies), dbcAMP (49 μg/ml, Sigma) and doxycycline (2 μg/ml, Sigma). Media was replaced the next day (day 7) with NMM containing AraC (2 μg/ml, Sigma) to eliminate any remaining non-differentiated cells. On day 8, AraC was removed and astrocytes were plated on top of neurons to support neuron cultures in NMM containing hbEGF (5 ng/ml, Peprotech). Cells were transduced with AAV on day 10 and assayed on day 24.

Computational Pipeline for Cas13d Identification

We obtained whole genome, chromosome, and scaffold-level prokaryotic genome assemblies from NCBI Genome in June 2016 and compared CRISPRfinder, PILER-CR, and CRT for identifying CRISPR repeats. The 20 kilobase flanking regions around each putative CRISPR repeat was extracted to identify nearby proteins and predicted proteins using Python. Candidate Cas proteins were required to be >750 aa in length and within 5 proteins of the repeat array, and extracted CRISPR loci were filtered out if they contained Cas genes associated with known CRISPR systems such as types I-III CRISPR. Putative effectors were clustered into families via all-by-all BLASTp analysis followed by single-linkage hierarchical clustering where a bit score of at least 60 was required for cluster assignment. Each cluster of at least 2 proteins was subjected to BLAST search against the NCBI non-redundant (nr) protein database, requiring a bit score >200 to assign similarity. The co-occurrence of homologous proteins in each expanded cluster to a CRISPR array was analyzed and required to be >70%. Protein families were sorted by average amino acid length and multiple sequence alignment for each cluster was performed using Clustal Omega and the Geneious aligner with a Blosum62 cost matrix. The RxxxxH HEPN motif was identified in the Cas13d family on the basis of this alignment. TBLASTN was performed on all predicted Cas13d effectors against public metagenome whole genome shotgun sequences without predicted open reading frames (ORFs). The Cas13d family was regularly updated via monthly BLAST search on genome and metagenome databases to identify any newly deposited sequences. New full-length homologs and homologous fragments were aligned using Clustal Omega and clustered using PhyML 3.2. CRISPRDetect was used to predict the direction of direct repeats in the Cas13d array and

DR fold predictions were performed using the Andronescu 2007 RNA energy model at 37° C. Sequence logos for Cas13d direct repeats were generated using Geneious 10. Protein Expression and Purification

Recombinant Cas13d proteins were PCR amplified from genomic DNA extractions of cultured isolates or metagenomic samples and cloned into a pET-based vector with an N-terminal His-MBP fusion and TEV protease cleavage site. The resulting plasmids were transformed into Rosetta2 (DE3) cells (Novagen), induced with 200 μM IPTG at OD₆₀₀ 0.5, and grown for 20 hours at 18° C. Cells were then pelleted, freeze-thawed, and resuspended in Lysis Buffer (50 mM HEPES, 500 mM NaCl, 2 mM MgCl₂, 20 mM Imidazole, 1% v/v Triton X-100, 1 mM DTT) supplemented with 1× protease inhibitor tablets, 1 mg/mL lysozyme, 2.5 U/mL Turbo DNase (Life Technologies), and 2.5 U/mL salt active nuclease (Sigma Aldrich). Lysed samples were then sonicated and clarified via centrifugation (18,000×g for 1 hour at 4° C.), filtered with 0.45 μM PVDF filter and incubated with 50 mL of Ni-NTA Superflow resin (Qiagen) per 10 L of original bacterial culture for 1 hour. The bead-lysate mixture was applied to a chromatography column, washed with 5 column volumes of Lysis Buffer, and 3 column volumes of Elution Buffer (50 mM HEPES, 500 mM NaCl, 300 mM Imidazole, 0.01% v/v Triton X-100, 10% glycerol, 1 mM DTT). The samples were then dialyzed overnight into TEV Cleavage Buffer (50 mM Tris-HCl, 250 mM KCl, 7.5% v/v glycerol, 0.2 mM TCEP, 0.8 mM DTT, TEV protease) before cation exchange (HiTrap SP, GE Life Sciences) and gel filtration (Superdex 200 16/600, GE Life Sciences). Purified, eluted protein fractions were pooled and frozen at 4 mg/mL in Protein Storage Buffer (50 mM Tris-HCl, 1M NaCl, 10% glycerol, 2 mM DTT).

Preparation of Guide and Target RNAs

Oligonucleotides carrying the T7 promoter and appropriate downstream sequence were synthesized (IDT) and annealed with an antisense T7 oligo for crRNAs and PCR-amplified for target and array templates. Homopolymer target RNAs were synthesized by Synthego. The oligo anneal and PCR templates were in vitro transcribed with the Hiscribe T7 High Yield RNA Synthesis kit (New England Biolabs) at 31° C. for 12 hours. For labeled targets, fluorescently labelled aminoallyl-UTP atto 680 (Jena Biosciences) was additionally added at 2 mM. Guide RNAs were purified with RNA-grade Agencourt AMPure XP beads (Beckman Coulter) and arrays and targets were purified with MEGAclear Transcription Clean-Up Kit (Thermo Fisher) and frozen at -80° C. For ssDNA and dsDNA targets, corresponding oligonucleotide sequences were synthesized (IDT) and either gel purified, or PCR amplified and then subsequently gel purified respectively.

Biochemical Cleavage Reactions

Purified EsCas13d protein and guide RNA were mixed (unless otherwise indicated) at 2:1 molar ratio in RNA Cleavage Buffer (25 mM Tris pH 7.5, 15 mM Tris pH 7.0, 1 mM DTT, 6 mM MgCl₂). The reaction was prepared on ice and incubated at 37° C. for 15 minutes prior to the addition of target at 1:2 molar ratio relative to EsCas13d. The reaction was subsequently incubated at 37° C. for 45 minutes and quenched with 1 μL of enzyme stop solution (10 mg/mL Proteinase K, 4M Urea, 80 mM EDTA, 20 mM Tris pH 8.0) at 37° C. for 15 minutes. The reaction was then denatured with 2× RNA loading buffer (2×: 13 mM Ficoll, 8 M Urea, 25 mM EDTA), at 85° C. for 10 minutes, and separated on a 10% TBE-Urea gel (Life Technologies). Gels containing labeled targets were visualized on the Odyssey

Imager (Li-Cor); unlabeled array or target cleavage gels were stained with SYBR Gold prior to imaging via Gel Doc EZ system (Bio-Rad).

Transient transfection of Human Cell Lines

Engineered Cas13 coding sequences were cloned into a standardized plasmid expression backbone containing an EF1a promoter and prepared using the Nucleobond Xtra Midi EF Kit (Machery Nagel) according to the manufacturer's protocol. NLS-LwaCas13a-msfGFP and PspCas13b-NES-HIV were PCR amplified from Addgene #103854, and #103862, respectively, a gift from Feng Zhang. Cas13d pre-gRNAs and gRNAs were cloned into a minimal backbone containing a U6 promoter. shRNAs and guides for LwaCas13a were cloned into the same backbone and position matched to their corresponding guide RNA at the 3' of the target sequence. Matched gRNAs for PspCas13b were moved to the closest 5'-G nucleotide.

For transient transfection, HEK 293FT cells were plated at a density of 20,000 cells per well in a 96-well plate and transfected at >90% confluence with 200 ng of Cas13 expression plasmid and 200 ng of gRNA expression plasmid using Lipofectamine 2000 (Life Technologies) according to the manufacturer's protocol. Transfected cells were harvested 48-72 hours post-transfection for flow cytometry, gene expression analysis, or other downstream processing.

For reporter assays, HEK 293FT cells were transfected in 96-well format with 192 ng of Cas13d expression plasmid, 192 ng of guide expression plasmid, and 12 ng of mCherry expression plasmid with Lipofectamine 2000 (Life Technologies). Cells were harvested after 48 hours and analyzed by flow cytometry.

U2OS cells were plated at a density of 20,000 cells per well in a 96-well plate and transfected at >90% confluence with 100 ng of Cas13d expression plasmid using Lipofectamine 3000 (Life Technologies) according to the manufacturer's protocol and processed for immunocytochemistry after 48 h.

Flow Cytometry

Cells were dissociated 48 hours post-transfection with TrypLE Express and resuspended in FACS Buffer (1× DPBS^{-/-}, 0.2% BSA, 2 mM EDTA). Flow cytometry was performed in 96-well plate format using a MACSQuant VYB (Miltenyi Biotec) and analyzed using FlowJo 10. RG6 was a gift from Thomas Cooper (Addgene plasmid # 80167) and modified to replace EGFP with mTagBFP2. All represented samples were assayed with three biological replicates. In the mCherry reporter assay, data is representative of at least 20,000 gated events per condition. In the splicing reporter assay, data is representative of at least 2,500 gated events per condition.

Gene Expression Analysis

Cells were lysed 48 hours post-transfection with DTT-supplemented RLT buffer and total RNA was extracted using RNeasy Mini Plus columns (Qiagen). 200 ng of total RNA was then reverse transcribed using random hexamer primers and Revertaid Reverse Transcriptase (Thermo Fisher) at 25° C. for 10 min, 37° C. for 60 min, and 95° C. for 5 min followed by qPCR using 2× Taqman Fast Advanced Master Mix (Life Technologies) and Taqman probes for GAPDH and the target gene as appropriate (Life Technologies and IDT). Taqman probe and primer sets were generally selected to amplify cDNA across the Cas13 or shRNA target site position to prevent detection of cleaved transcript fragments (see Table S4 of Konermann et al., Cell 173:1-12, 2018, herein incorporated by reference in its entirety). qPCR was carried out in 5 μL multiplexed reactions and 384-well format using the LightCycler 480 Instrument II (Roche).

Fold-change was calculated relative to GFP-transfected vehicle controls using the ddCt method. One-way or two-way ANOVA with multiple comparison correction was used to assess statistical significance of transcript changes using Prism 7.

Immunohistochemistry

For immunohistochemical analysis, U2OS cells were cultured on 96-well optically clear plates (Greiner Bio-One), transfected as previously described, then fixed in 4% PFA (Electron Microscopy Sciences) diluted in PBS (Gibco) and washed with 0.3M glycine (Sigma) in PBS to quench PFA. Samples were blocked and permeabilized in a PBS solution containing 8% donkey serum (Jackson ImmunoResearch), 8% goat serum (Cell Signaling Technologies), and 0.3% Triton-X 100 (Sigma) for one hour, followed by primary antibody incubation in 1% BSA (Fisher Bioreagents), 1% goat serum, and 0.25% Triton-X overnight at 4° C. Samples were washed 3 times with PBS containing 0.1% BSA and 0.1% Triton-X 100 before incubating with fluorophore-conjugated secondary antibodies in PBS with 0.05% Triton-X 100 and 1% BSA at room temperature for one hour. Cells were washed with PBS with 0.1% Triton-X, stained with DAPI, and then covered with Mounting Media (Ibidi) before imaging. Primary antibody, HA-Tag 6E2 (Cell Signaling, 2367), was used at a 1:100 dilution as per manufacturer's instructions. Secondary antibodies used were goat anti-mouse IgG1-Alexa-Fluor 647 (Thermo Fisher, A21240) and Anti-Mouse IgG1 CF 633 (Sigma, SAB4600335). Confocal images were taken using a Zeiss Airyscan LSM 880 followed by image processing in Zen 2.3 (Zeiss).

Bacterial Small RNA Sequencing and Analysis

E. coli DH5 α cells were transfected with pACYC184 carrying the CRISPR-Cas13d locus derived from an uncultured *Ruminococcus* sp. strain. Cells were harvested in stationary phase, rinsed in PBS, resuspended in TRIzol (Life Technologies), transferred to Lysing Matrix B tubes containing 0.1 mm silica beads (MP Biomedicals), and homogenized on a Bead Mill 24 (Fisher Scientific) for three 30-second cycles. Total RNA was isolated by phenol-chloroform extraction, then purified using the DirectZol Mini-prep Kit (Zymo Research). RNA quality was assessed on an Agilent 2200 TapeStation followed by Turbo DNase treatment (Ambion). Total RNA was treated with T4 Polynucleotide Kinase (NEB) and rRNA-depleted using the Ribo-Zero rRNA Removal Kit for bacteria (Illumina). RNA was treated with RNA 5' polyphosphatase, poly(A)-tailed with *E. coli* poly(A) polymerase, and ligated with 5' RNA sequencing adapters using T4 RNA ligase 1 (NEB). cDNA was generated via reverse transcription using an oligo-dT primer and M-MLV RT/RNase Block (AffinityScript, Agilent) followed by PCR amplification and barcoding. Resulting libraries were sequenced on Illumina MiSeq, demultiplexed using custom Python scripts, and aligned to the Cas13d CRISPR locus using Bowtie 2. Alignments were visualized with Geneious.

Ngn2 Lentivirus Preparation

Low passage HEK 293FT cells were transfected with Polyethylenimine Max (PEI, Polysciences) and Ngn2 target plasmid plus pMDG.2 and psPAX2 packaging plasmids (a gift from Didier Trono, Addgene #12259 and #12260) in DMEM +10% FBS media during plating. The following day, media was changed to serum-free chemically defined minimal medium (Ultraplex supplemented with Glutamax, Lonza). Viral supernatant was harvested 48 h later, clarified through a 0.45 micron PVDF filter (Millipore) and concentrated using ultracentrifugation.

AAV preparation

Low passage HEK 293FT cells were transfected with Polyethylenimine Max (PEI, Polysciences) and AAV target plasmid plus AAV1 serotype and pAdDeltaF6 helper packaging plasmids (UPenn Vector Core) in DMEM+10% FBS media during plating. The following day, 60% of the media was changed to chemically defined minimal medium (Ultraplex supplemented with Glutamax, Lonza). 48 h later, AAV-containing supernatant was harvested and clarified through a 0.45 μ m PVDF filter (Millipore) and concentrated using precipitation by polyethylene glycol (PEG virus precipitation kit #K904, Biovision) following the manufacturer's protocol.

RNA-Seq Library Preparation and Sequencing

48 h after transfection, total RNA was extracted from 293FT cells using the RNeasy Plus Mini kit from Qiagen. Stranded mRNA libraries were prepared using the NEBNext II Ultra Directional RNA Library Prep Kit from New England Biolabs (Cat# E7760S) and sequenced on an Illumina NextSeq500 with 42 nt paired end reads. ~15M total reads were demultiplexed per condition.

RNA-Seq Analysis

Sequenced reads were quality-tested using FASTQC and aligned to the hg19 human genome using the 2.5.1b STAR aligner (Dobin et al., 2013). Mapping was carried out using default parameters (up to 10 mismatches per read, and up to 9 multi-mapping locations per read). The genome index was constructed using the gene annotation supplied with the hg19 Illumina iGenomes collection (Illumina) and sjdbOverhang value of 100. Uniquely mapped reads were quantified across all gene exons using the top-expressed isoform as proxy for gene expression with the HOMER analysis suite (Heinz et al., 2010), and differential gene expression was carried out with DESeq2 v 1.14.1 (Love et al., 2014) using triplicates to compute within-group dispersion and contrasts to compare between targeting and non-targeting conditions. Significant differentially expressed genes were defined as having a false discovery rate (FDR) <0.01 and a log₂ fold change >0.75. Volcano plots were generated in R 3.3.2 using included plotting libraries and the alpha() color function from the scales 0.5.0 package.

Statistics

All values are reported as mean \pm SD or mean \pm SEM as indicated in the appropriate figure legends. For comparing two groups, a one-tailed student's t-test was used and statistical significance was determined using the Holm-Sidak method with alpha=0.05. A one-way ANOVA with Tukey multiple hypothesis correction was used to assess significance between more than two groups. Two-way ANOVA was used when comparing across two factors (i.e., RNA targeting modality and guide position) and adjusted for multiple hypothesis correction by Sidak's multiple comparisons test. For comparing groups that were found to not meet the assumption of a normal distribution by a D'Agostino and Pearson normality test, the non-parametric Friedman test with Dunn's multiple comparison adjustment was performed. PRISM 7.0 was used for all statistical analysis. Sample sizes were not determined a priori. At least three biological replicates were used for each experiment, as indicated specifically in each figure.

Sequencing data reported herein can be found in the NCBI Gene Expression Omnibus under GEO Series accession number GSE108519.

Additional details on the materials and methods used, such as sequences (e.g., Tables S1 to S5), can be found in Konermann et al., Cell 173:1-12, 2018, herein incorporated by reference in its entirety.

Computational Identification of a Type VI-Like Cas Ribonuclease Family

This example describes methods used to identify previously undetected or uncharacterized RNA-targeting CRISPR-Cas systems by developing a computational pipeline for class 2 CRISPR-Cas loci, which require only a single nuclease for CRISPR interference such as Cas9, Cas12a (formerly Cpf1), or Cas13a (formerly C2c2) (Makarova et al., 2015; Shmakov et al., 2015). To improve upon previous strategies for bioinformatic mining of CRISPR systems, which focus on discovering sets of conserved Cas genes involved in spacer acquisition (Shmakov et al., 2015), the minimal requirements for a CRISPR locus to be the presence of a CRISPR repeat array and a nearby effector nuclease were defined. Using the CRISPR array as a search anchor, all prokaryotic genome assemblies and scaffolds were obtained from the NCBI WGS database and adapted algorithms for de novo CRISPR array detection (Bland et al., 2007; Edgar, 2007; Grissa et al., 2007) to identify 21,175 putative CRISPR repeat arrays (FIG. 1A).

Up to 20 kilobases (kb) of genomic DNA sequence flanking each CRISPR array was extracted to identify predicted protein-coding genes in the immediate vicinity. Candidate loci containing signature genes of known class 1 and class 2 CRISPR-Cas systems such as Cas3 or Cas9 were excluded from further analysis, except for Cas12a and Cas13a to judge the ability of the pipeline to detect and cluster these known class 2 effector families. To identify new class 2 Cas effectors, it was required that candidate proteins be >750 residues in length and within 5 protein-coding genes of the repeat array, as large proteins closely associated with CRISPR repeats are key characteristics of known single effectors. The resulting proteins were classified into 408 putative protein families using single-linkage hierarchical clustering based on homology.

To discard protein clusters that reside in close proximity to CRISPR arrays due to chance or overall abundance in the genome, additional homologous proteins to each cluster were identified from the NCBI non-redundant protein database and their proximity to a CRISPR array determined. Reasoning that true Cas genes would have a high co-occurrence rate with CRISPR repeats, >70% of the proteins for each expanded cluster were required to exist within 20 kb of a CRISPR repeat. These remaining protein families were analyzed for nuclease domains and motifs.

Among the candidates, which include the recently described Cas13b system (Smargon et al., 2017), a family of uncharacterized putative class 2 CRISPR-Cas systems encoding a candidate CRISPR-associated ribonuclease containing 2 predicted HEPN ribonuclease motifs (Anantharaman et al., 2013) were identified (FIG. 2A). Importantly, they are among the smallest class 2 CRISPR effectors described to date (~930 aa). The Type VI CRISPR-Cas13 superfamily is exemplified by sequence-divergent, single-effector signature nucleases and the presence of two HEPN domains. Other than these two RxxxxH HEPN motifs (FIG. 3A), the candidate effectors have no significant sequence similarity to previously described Cas13 enzymes, so his family of putative CRISPR ribonucleases is designated as Type VI Cas13d, or Type VI-D (FIG. 3B).

CRISPR-Cas13d systems are derived from gut-resident microbes, so we sought to expand the Cas13d family via alignment to metagenomic contigs from recent large-scale microbiome sequencing efforts. Comparison of Cas13d pro-

teins against public metagenome sequences without predicted open reading frames (ORFs) identified additional full-length systems as well as multiple effector and array fragments that cluster in several distinct branches (FIG. 1B).

To generate full-length Cas13d ortholog proteins and loci from the different branches of the Cas13d protein family, genomic DNA samples we43 obtained from associated assemblies and performed targeted Sanger sequencing to fill in gaps due to incomplete sequencing coverage, such as for the metagenomic ortholog 'Anaerobic digester metagenome' (Adm) (Treu et al., 2016).

Cas13d CRISPR loci are largely clustered within benign, Gram-positive gut bacteria of the genus *Ruminococcus*, and exhibit a surprising diversity of CRISPR locus architectures (FIG. 2A). With the exception of the metagenomic AdmCas13d system, Cas13d systems lack the key spacer acquisition protein Cas1 (Yosef et al., 2012) within their CRISPR locus, highlighting the utility of a class 2 CRISPR discovery pipeline without Cas1 or Cas2 gene requirements. Cas13d direct repeats (DRs) are highly conserved in length and predicted secondary structure (FIG. 3C), with a 36 nt length, an 8-10 nt stem with A/U-rich loop, and a 5'-AAAAC motif at the 3' end of the direct repeat (FIG. 3D). This conserved 5'-AAAAC motif has been shown to be specifically recognized by a type II Cas2 spacer acquisition complex (Wright and Doudna, 2016). In fact, Cas1 can be found in relative proximity to some Cas13d systems (within 10-30 kb for PIE0 and Rfx) while the remaining Cas13d-containing bacteria contain Cas1 elsewhere in their genomes, likely as part of another CRISPR locus.

EXAMPLE 3

CRISPR-Cas13d Possesses Dual RNase Activities

To demonstrate that the Cas13d repeat array is transcribed and processed into CRISPR guide RNAs (gRNA), the Cas13d CRISPR locus was cloned from an uncultured *Ruminococcus* sp. sample (Ur) into a bacterial expression plasmid. CRISPR systems tend to form self-contained operons with the necessary regulatory sequences for independent expression, facilitating heterologous expression in *E. coli* (Gasiunas et al., 2012). RNA sequencing (Heidrich et al., 2015) revealed processing of the array into ~52 nt mature gRNAs, with a 30 nt 5' direct repeat followed by a variable 3' spacer that ranged from 14-26 nt in length (FIG. 2B).

To characterize Cas13d properties in vitro, *Eubacterium siraeum* Cas13d protein (EsCas13d) was purified based on its robust recombinant expression in *E. coli* (FIGS. 4A-4C) and found that EsCas13d was solely sufficient to process its matching CRISPR array into constituent guides without additional helper ribonucleases (FIG. 2C, Table S1 of Konermann et al., "Transcriptome Engineering with RNA-Targeting Type VI-D CRISPR Effectors," Cell 173:1-12, 2018, herein incorporated by reference in its entirety), a property shared by some class 2 CRISPR-Cas systems (East-Seletsky et al., 2016; Fonfara et al., 2016; Smargon et al., 2017). Furthermore, inactivating the positively charged catalytic residues of the HEPN motifs (Anantharaman et al., 2013) (dCas13d: R295A, H300A, R849A, H854A) did not affect array processing, indicating a distinct RNase activity dictating gRNA biogenesis analogous to Cas13a (East-Seletsky et al., 2016; Liu et al., 2017).

Cas effector proteins typically form a binary complex with mature gRNA to generate an RNA-guided surveillance ribonucleoprotein capable of cleaving foreign nucleic acids for immune defense (van der Oost et al., 2014). To assess if

Cas13d has programmable RNA targeting activity as indicated by the presence of two HEPN motifs, EsCas13d protein was paired with an array or a mature gRNA along with a cognate *in vitro*-transcribed target. Based on the RNA sequencing results, a mature gRNA containing a 30 nt direct repeat and an intermediate spacer length of 22 nt was selected (nucleotides 6-36 of SEQ ID NO: 129, followed by 22 bases complementary to the RNA target).

Cas13d was able to efficiently cleave the complementary target ssRNA with both the unprocessed array and mature gRNA in a guide-sequence dependent manner, while non-matching spacer sequences abolished Cas13d activity (FIG. 5A). Substitution with dCas13d or the addition of EDTA to the cleavage reaction also abolished guide-dependent RNA targeting, indicating that Cas13d targeting is HEPN- and Mg²⁺-dependent (FIG. 5B). To determine the minimal spacer length for efficient Cas13d targeting, a series of spacer truncations ranging from the unprocessed 30 nt length down to 10 nt were generated (FIG. 6A). Cleavage activity dropped significantly below a 21 nt spacer length, confirming the choice of a 22 nt spacer (FIG. 6B).

RNA-targeting class 2 CRISPR systems have been proposed to act as sensors of foreign RNAs (Abudayyeh et al., 2016; East-Seletsky et al., 2016), where general RNase activity of the effector nuclease is triggered by a guide-matching target. To assay for a similar property in Cas13d, RNase activity of the binary EsCas13d:gRNA complex was monitored in the presence of a matching RNA target. It was observed that EsCas13d can be activated by target RNA to cleave bystander RNA targets (FIG. 3C), albeit inefficiently relative to its activity on the complementary ssRNA target. Bystander cleavage is guide sequence- and HEPN-dependent, as the presence of non-matching bystander target alone was insufficient to induce cleavage while substitution of dCas13d or addition of EDTA abolished activity. These results indicate that bystander RNase activity may be a general property of RNA-targeting class 2 systems in CRISPR adaptive bacterial immunity (FIG. 3D).

To assess the generalizability of Cas13d reprogramming, twelve guides tiling a complementary RNA target were generated and efficient cleavage in all cases was observed (FIG. 7A). Cas13d was unable to cleave a ssDNA (FIG. 6C) or dsDNA (FIG. 6D) version of the ssRNA target, indicating that Cas13d is an RNA-specific nuclease. Further, RNA target cleavage did not depend on the protospacer flanking sequence (PFS) (FIG. 7A) in contrast to other RNA-targeting class 2 systems, which require a 3'-H (Abudayyeh et al., 2016) or a double-sided, DR-proximal 5'-D and 3'-NAN or NNA (Smargon et al., 2017). Although a slight bias against an adenine PFS was initially observed (FIG. 6E), varying the target PFS base with a constant guide sequence resulted in no significant differences (P=0.768) in targeting efficiency (FIG. 6F).

While DNA-targeting class 2 CRISPR systems (Gasiunas et al., 2012; Jinek et al., 2012; Zetsche et al., 2015) and some RNA-targeting class 1 systems tend to cleave at defined positions relative to the target-guide duplex (Samai et al., 2015; Zhang et al., 2016), the Cas13d cleavage pattern varies for different targets (FIGS. 5A, 5C, 6H) and remains remarkably similar despite the guide sequence position (FIG. 7A). This indicates that Cas13d may preferentially cleave specific sequences or structurally accessible regions in the target RNA. Cas13d was tested activity on targets containing variable homopolymer repeats in the loop region of a hairpin or as a linear single-stranded repeat. EsCas13d

exhibited significant preference for uracil bases in both target structures, with lower but detectable activity at all other bases (FIG. 7B).

Cas enzymes are found in nearly all archaea and about half of bacteria (Hsu et al., 2014; van der Oost et al., 2014), spanning a wide range of environmental temperatures. To determine the optimal temperature range for Cas13d activity, a spectrum of cleavage temperature conditions from 16-62° C. was tested and observed maximal activity in the 24-41° C. range (FIG. 6G, 6H). This temperature range is compatible with a wide range of prokaryotic and eukaryotic hosts, indicating Cas13d can be adapted for RNA targeting in different cells and organisms.

EXAMPLE 4

Cell-Based Activity Screen of Engineered Orthologs

The Cas13d nuclease was used as a flexible tool for programmable RNA targeting in mammalian cells. CRISPR orthologs from distinct bacterial species commonly exhibit variable activity (Abudayyeh et al., 2017; East-Seletsky et al., 2017), especially upon heterologous expression in human cells (Ran et al., 2015; Zetsche et al., 2017). Highly active Cas13d orthologs were identified in a eukaryotic cell-based mCherry reporter screen.

By synthesizing human codon-optimized versions of 7 orthologs from distinct branches within the Cas13d family (FIG. 1B), mammalian expression plasmids carrying the catalytically active and HEPN-inactive proteins were generated. Each protein was then optionally fused to N- and C-terminal nuclear localization signals (NLS). These Cas13d effector designs were HA-tagged and paired with two distinct guide RNA architectures, either with a 30 nt spacer flanked by two direct repeat sequences to mimic an unprocessed guide RNA (pre-gRNA) or a 30 nt direct repeat with 22 nt spacer (gRNA) predicted to mimic mature guide RNAs (FIG. 8A). For each guide design, four distinct spacer sequences complementary to the mCherry transcript were then pooled to minimize potential spacer-dependent variability in targeting efficiency. The ability of Cas13d to knockdown mCherry protein levels was determined in a human embryonic kidney (HEK) 293FT cell-based reporter assay.

48 hours post-transfection, flow cytometry indicated that RfxCas13d and AdmCas13d efficiently knocked down mCherry protein levels by up to 92% and 87% (P<0.0003), respectively, relative to a non-targeting control guide (FIG. 8B). In contrast, EsCas13d along with RaCas13d and RffCas13d exhibited limited activity in human cells. Furthermore, none of the HEPN-inactive Rfx-dCas13d constructs significantly affected mCherry fluorescence, indicating HEPN-dependent knockdown (P>0.43 for all cases). Robust nuclear translocation of the Rfx and AdmCas13d NLS fusion constructs was observed via immunocytochemistry, while the wild-type effectors remain primarily extranuclear (FIG. 8C).

Proceeding with RfxCas13d and AdmCas13d as lead candidates, we next compared their ability to knockdown endogenous transcripts. To determine the optimal ortholog and guide architecture, the capability of Rfx and AdmCas13d construct variants to target β -1,4-N-acetylgalactosaminyl transferase 1 (B4GALNT1) transcripts was systematically assayed. In each condition, four guides containing distinct spacer sequences tiling the B4GALNT1 transcript were pooled. The RfxCas13d-NLS fusion targeted

B4GALNT1 more efficiently than wild-type RfxCas13d and both variants of AdmCas13d, with both the gRNA and pre-gRNA mediating potent knockdown (~82%, $P < 0.0001$) (FIG. 8D). Cas13d-NLS from *Ruminococcus flavefaciens* strain XPD3002 was therefore selected for the remaining experiments (CasRx).

EXAMPLE 5

Programmable RNA Knockdown in Human Cells with CasRx

Because Cas13d is capable of processing its own CRISPR array, this property was leveraged for the simultaneous delivery of multiple targeting guides in a simple single-vector system (FIG. 9A). Arrays encoding four spacers that each tile the transcripts of mRNAs (B4GALNT1 and ANXA4) or nuclear localized lncRNAs (HOTTIP and MALAT1) consistently facilitated robust (>90%) RNA knockdown by CasRx ($P < 0.0001$) (FIG. 9B).

CasRx was compared to more established technologies for transcript knockdown or repression, by comparing CasRx-mediated RNA interference to dCas9-mediated CRISPR interference (Gilbert et al., 2014; Gilbert et al., 2013) and spacer sequence-matched shRNAs via transient transfection (FIG. 9C). For CRISPRi-based repression, the most potent dCas9 guide for B4GALNT1 from previous reports was analyzed (Gilbert et al., 2014; Zalatan et al., 2015). Across 3 endogenous transcripts, CasRx outperformed shRNAs (11/11) and CRISPRi (4/4) in each case (FIG. 9D), exhibiting a median knockdown of 96% compared to 65% for shRNA and 53% for CRISPRi after 48 hours. In addition, knockdown by CasRx was compared to two recently described Cas13a and Cas13b effectors (Abudayyeh et al., 2017; Cox et al., 2017) (FIG. 10A). Across three genes and eight guide RNAs, CasRx mediated significantly greater transcript knockdown than both LwaCas13a-msfGFP-NLS and PspCas13b-NES (median: 97% compared to 80% and 66% respectively, $P < 0.0001$) (FIG. 10B).

RNAi has been widely used to disrupt any gene of interest due to a combination of simple re-targeting principles, scalable synthesis, knockdown potency, and ease of reagent delivery. However, widespread off-target transcript silencing has been a consistent concern (Jackson et al., 2003; Sigoillot et al., 2012), possibly due to the entry of RNAi reagents into the endogenous miRNA pathway (Doench et al., 2003; Smith et al., 2017). Consistent with these reports, upon RNA sequencing of human cells transfected with a B4GALNT1-targeting shRNA, widespread off-target transcriptional changes relative to a non-targeting shRNA were observed (>500 significant off-target changes, $P < 0.01$, FIGS. 9E, 9G). In contrast, transcriptome profiling of spacer-matched CasRx guide RNAs revealed no significant off-target changes other than the targeted transcript (FIG. 9F). This indicates that the moderate bystander cleavage observed in vitro (FIG. 5C) may not result in observable off-target transcriptome perturbation in mammalian cells. A similar pattern was observed when targeting ANXA4 (FIGS. 11A-11B), with over 900 significant off-target changes resulting from shRNA targeting compared to zero with CasRx (FIG. 9G).

To confirm that CasRx interference is broadly applicable, a panel of 11 additional genes with diverse roles in cancer, cell signaling, and epigenetic regulation were selected and 3 guides per gene were screened. CasRx consistently mediated high levels of transcript knockdown across genes with a median reduction of 96% (FIG. 9H). Each tested guide

mediated at least 80% knockdown, underscoring the consistency of the CasRx system for RNA interference.

EXAMPLE 6

Splice Isoform Engineering with dCasRx

The experiments on RNA targeting with CasRx revealed that target RNA and protein knockdown is dependent on the catalytic activity of the HEPN domains (FIGS. 8B, 5B). The same guide sequences mediating efficient knockdown with CasRx failed to significantly reduce mCherry levels when paired with catalytically inactive dCasRx (FIG. 8B), indicating that targeting of dCasRx to the coding portion of mRNA does not necessarily perturb protein translation. This observation indicated the possibility of utilizing dCasRx for targeting of specific coding and non-coding elements within a transcript to study and manipulate RNA. To validate this concept, the utility of the dCasRx system was expanded by creating a splice effector.

Alternative splicing is generally regulated by the interaction of cis-acting elements in the pre-mRNA with positive or negative trans-acting splicing factors, which can mediate exon inclusion or exclusion (Matera and Wang, 2014; Wang et al., 2015). It was reasoned that dCasRx binding to such motifs may be sufficient for targeted isoform perturbation. For proof-of-concept, distinct splice elements were identified in a bichromatic splicing reporter containing DsRed upstream of mTagBFP2 in two different reading frames following an alternatively spliced exon (Orengo et al., 2006) (FIG. 12A). Inclusion or exclusion of this second exon toggles the reading frame and resulting fluorescence, facilitating quantitative readout of splicing patterns by flow cytometry. To mediate exon skipping, four guide RNAs were designed to target the intronic branchpoint nucleotide, splice acceptor site, putative exonic splice enhancer, and splice donor of exon 2.

One widespread family of negative splice factors are the highly conserved heterogeneous nuclear ribonucleoproteins (hnRNPs), which typically inhibit exon inclusion via a C-terminal, glycine-rich domain (Wang et al., 2015). The splicing reporter was targeted with dCasRx and engineered fusions to the Gly-rich C-terminal domain of hnRNP1, one of the most abundant hnRNP family members (FIG. 12B).

Guide position appears to be a major determinant of the efficiency of engineered exon skipping. While each guide position mediated a significant increase in exon exclusion ($P < 0.0001$ in all cases) relative to the non-targeting guide, targeting the splice acceptor resulted in the most potent exon exclusion (increase from 8% basal skipping to 65% for dCasRx alone and 75% with hnRNP1 fusion). By comparison, dLwaCas13a-msfGFP-NLS mediated significantly lower levels of exon skipping across all four positions (19% skipping for splice acceptor guide) (FIGS. 10C and 10D, $P < 0.0001$).

Targeting all 4 positions simultaneously with a CRISPR array achieved higher levels of exon skipping than individual guides alone (81% for dCasRx and 85% for hnRNP1 fusion, $P < 0.006$ compared to SA guide) (FIG. 12B). These results indicate that dCasRx allows for tuning of isoform ratios through varying guide placement and suggest that it can be leveraged as an efficient RNA binding module in human cells for targeting and manipulation of specific RNA elements.

Viral Delivery of dCasRx to a Neuronal Model of Frontotemporal Dementia

The Cas13d family averages 930 amino acids in length, in contrast to Cas9 (~1100 aa to ~1400 aa depending on subtype, with compact outliers such as CjCas9 or SaCas9), Cas13a (1250 aa), Cas13b (1150 aa), and Cas13c (1120 aa) (FIG. 3B) (Chylinski et al., 2013; Cox et al., 2017; Hsu et al., 2014; Kim et al., 2017; Shmakov et al., 2015; Smargon et al., 2017). Although adeno-associated virus (AAV) is a versatile vehicle for transgene delivery and gene therapy due to its broad range of capsid serotypes, low levels of insertional mutagenesis, and lack of apparent pathogenicity, its limited packaging capacity (~4.7 kb) makes it challenging to effectively deliver many single effector CRISPR enzymes (Abudayyeh et al., 2017; Ran et al., 2015; Swiech et al., 2015). The remarkably small size of Cas13d effectors render them uniquely suited for all-in-one AAV delivery with a CRISPR array, an optional effector domain, and requisite expression or regulatory elements (FIG. 12C).

Frontotemporal Dementia with Parkinsonism linked to Chromosome 17 (FTDP-17) is an autosomal dominant major neurodegenerative disease caused by diverse point mutations in MAPT, the gene encoding for tau. Tau exists as two major isoforms in human neurons, 4R and 3R, which are distinguished by the presence or absence of tau exon 10 and thus contain 4 or 3 microtubule binding domains. The balance of these two isoforms is generally perturbed in FTDP-17 as well as other tauopathies, driving the progression of neurodegeneration (Boeve and Hutton, 2008). Some forms of FTD are caused by mutations in the intron following MAPT exon 10 which disrupt an intronic splice silencer and elevate the expression of 4R tau (Kar et al., 2005), thereby inducing pathological changes (Schoch et al., 2016).

It was reasoned that dCasRx targeted to MAPT exon 10 could induce exon exclusion to alleviate dysregulated 4R/3R tau ratios. Patient-derived human induced pluripotent stem cells (hiPSCs) were differentiated into cortical neurons via Neurogenin-2 directed differentiation for 2 weeks (Zhang et al., 2013). Postmitotic neurons were then transduced with AAV1 carrying dCasRx (FIG. 12D) paired with a repeat array containing 3 spacers that target the exon 10 splice acceptor and two putative exonic splice enhancers (FIG. 12E). dCasRx-mediated exon exclusion was able to reduce the relative 4R/3R tau ratio by nearly 50% relative to a BFP vehicle control (FIG. 12F) and to a level similar to unaffected control neurons, demonstrating that CasRx can be exploited for transcriptional modulation in primary cell types via AAV delivery.

EXAMPLE 8

RNA Targeting in Human Cells Using Cas13d

RNA can be targeted in human cells using the active Cas13d nuclease. As a proof of concept, human U-2 OS bone osteosarcoma cells were stably integrated with an mCherry reporter and transfected with plasmids encoding human codon optimized Cas13d and guide RNAs targeting the mCherry transcript (FIG. 13).

Cas13d proteins were also fused with N- and C-terminal NLS sequences (SPKKKRKRKVEAS, SEQ ID NO: 256, for the N-terminal NLS and GPKKKRKRKVEAS, SEQ ID NO: 258, C-terminal NLS) to understand if nuclear localization can affect mCherry knockdown (these are denoted the 2x

NLS constructs). Guide RNAs were either provided in a vector with a U6 promoter operably linked to a 36 nt DR-30 nt spacer-36 nt DR sequence, which mimics the unprocessed CRISPR guide array (denoted DR36), or to a 30nt DR-22 nt spacer to mimic the processed, mature gRNA (denoted gRNA). The DR36 construct is presumed to be processed by Cas13d into mature gRNAs within the cell. The spacer sequences within the DR36 or gRNA molecules were either complementary to the mCherry target RNA (on-target mCherry) or computationally optimized to avoid complementarity to mCherry or any endogenous human transcript (non-targeting mCherry).

mCherry knockdown was quantified via flow cytometry and normalized to a transfection control. The non-targeting mCherry guides do not affect mCherry protein levels via flow cytometry, presumably because the mCherry transcripts are not targeted. However, the on-target mCherry guides paired with 4 different Cas13d orthologues exhibited significant mCherry knockdown (FIG. 13). “XPD” refers to *Ruminococcus flavefaciens* XPD3002 Cas13d (SEQ ID NO: 92); “P1E0” refers to Gut metagenome P1E0 Cas13d (SEQ ID NO: 83); “AnDig” refers to Anaerobic digester gut metagenome Cas13d (SEQ ID NO: 42); “Uncultured” refers to uncultured *Ruminococcus* sp. Cas13d.

EXAMPLE 9

In Vivo RNA Targeting Using Cas13d

RNA can be targeted in mouse models of cancer. To observe which cells in the mouse are expressing EGFR, a guide RNA is designed that includes one or more spacer regions complementary to mouse EGFR and is combined with Cas13d having a mutated HEPN domain (such as SEQ ID NO: 2 or 4), and a biotin label. The gRNA and Cas13d coding sequence are cloned into a viral vector (such as a lentivirus) which is used to infect the mice by tail vein injection at a titer to insure 100% infection rates. A fluorescent streptavidin label is administered to the mice. Cells expressing EGFR are visualized and detected with the appropriate excitation frequency for the fluorescent label. Alternatively, Cas13d is delivered in its active form in vivo to mediate target knock-down.

EXAMPLE 10

Treatment of Cancer

Human subjects with histologically confirmed stage 1, EGFR+ breast cancer can be treated with the disclosed methods. Each subject is administered a complex comprising an active Cas13d or a Cas13d protein mutated in the HEPN domain (such as SEQ ID NO: 2 or 4), a guide RNA targeting EGFR, and a toxin, after receiving lumpectomy surgery. Treated individuals are monitored for breast cancer recurrence.

EXAMPLE 11

Treatment of HIV Infection

Human subjects with HIV infection can be treated with the disclosed methods. Each subject is administered a construct comprising an active Cas13d or a Cas13d protein mutated in the HEPN domain (such as SEQ ID NO: 2 or 4),

a guide RNA targeting HIV Nef protein, and a toxin. Treated individuals are monitored for HIV progression.

EXAMPLE 12

Treatment of Huntington's Disease

Human subjects with Huntington's Disease can be treated with the disclosed methods. Each individual is administered a construct comprising Cas13d, a guide RNA targeting the Huntington mutation. Treated individuals are monitored for disease progression.

EXAMPLE 13

Alternative Splicing Using Cas13d

Cas13d splice effectors can be used for therapeutic protein restoration (for example that results from a mutation or deletion), gene knockdown via frameshift induction, tuning or restoring a desired isoform ratio, or inducing a desired dominant splice isoform (FIG. 14). Alternative splicing is generally regulated by the interaction of cis-acting elements in the pre-mRNA with positive or negative trans-acting splicing factors, which can mediate exon inclusion or exclusion. dCas13d and Cas13d, with optional fusion to positive or negative splicing factors, can be used as splicing effectors that target to said cis-acting elements in the pre-mRNA to manipulate splicing. Such elements can include exonic splicing enhancers or suppressors, intronic splicing enhancers or suppressors, splice acceptor and splice donor sites, and more generally protein- or RNA-interacting motifs or elements on that particular pre-mRNA, mRNA, or other RNA species, such as a non-coding RNA, tRNA, miRNA, and the like.

Additionally, that the effects of Cas13d-based splice effectors can be guide position-dependent. This can be exploited to perturb or discover particular motifs or sites in an RNA transcript such as protein-binding sites, via steric hindrance, blocking, recruitment, or effector-mediated interaction. For example, the interaction between non-coding RNAs and particular chromatin remodeling complexes can be perturbed. Access of the ribosomal binding site and other elements can be blocked in a 5' or 3' UTR (or to recruit appropriate effector domains) to decrease, increase, or otherwise manipulate translation.

Targeting or tiling Cas13d guides along a pre-mRNA can be used to discover or map new cis-acting elements such as intronic or exonic splice enhancers. This has been exploited in a therapeutic context in the case of the dystrophin gene for optimal antisense oligonucleotide positioning and can also be used for optimal Cas13d positioning. This can be also used to map, mask, or otherwise perturb RNA zipcodes or other cis-acting elements to affect trafficking and localization, chromatin remodeling, polyadenylation, RNA stability and half-life, or levels of nonsense-mediated decay.

In one example, targeting an RNA allows for changing splicing of the target RNA. Both the direct binding of splice acceptor and/or donor sites as well as splice effector domains can be used to manipulate splicing. For example, by using a dCas13d protein with a mutated HEPN domain (e.g., SEQ ID NO: 2 or 4), a guide RNA containing at least one a spacer sequence specific for the target RNA, and optionally an effector domain that affects splicing (such as the RS-rich domain of SRSF1, the Gly-rich domain of hnRNPA1, the alanine-rich motif of RBM4, or the proline-rich motif of DAZAP1), alternative splicing of the RNA can be achieved.

In some examples, such a method is used for exon inclusion, for example to include exon 2 of acid alpha-glucosidase (GAA) to treat Pompe disease or to include exon 7 of SMN2 to treat spinal muscular atrophy (SMA). In some examples, such a method is used for exon exclusion, for example to restore the reading frame of dystrophin to treat Duchenne muscular dystrophy, to shift the splicing of the Bcl-x pre-mRNA from the antiapoptotic long isoform to the proapoptotic short isoform to treat cancer, to shift the splicing of the MAPT transcript to affect ratios of 3R and 4R tau, or to manipulate the splicing of the lamin A transcript in the case of Hutchinson-Gilford progeria syndrome or other genetic diseases of aging.

In some examples, the method uses an Cas13d protein, optionally with a mutated HEPN domain, to mask splice acceptor or donor sites, for example to create neoantigens to make cold tumors hot. By affecting the splicing of certain target pre-mRNAs, this method can generate novel exon-exon junctions that can lead to the creation of neo-epitopes in cancer cells. This can make a cancer cell vulnerable to the immune system due to the display of unnatural antigens. In other examples, this method can be used to dynamically manipulate isoform ratios or to restore reading frame of a protein (e.g., dystrophin for Duchenne's muscular dystrophy).

EXAMPLE 14

AAV Delivery of Cas13d

As described in the examples above, Cas13d can be effectively packaged into AAV to mediate expression in cell types that are not amenable to plasmid delivery or for in vivo delivery of Cas13d. AAV delivery of nuclease active Cas13d can be used to mediate RNA target knock-down in the cell type of interest. Due to its small size compared to other single-effector CRISPR nucleases, Cas13d can be packaged together with a guide RNA or an array containing multiple guide RNAs in a single AAV vector.

EXAMPLE 15

Nucleic Acid-Based Diagnostics with Cas13d

Cas13d enzymes can be exploited for nucleic acid-based diagnostics within the context of a cell, using cell-free lysate derived from a cell, or a cell-free system containing an engineered Cas13d enzyme and guide RNA to facilitate formation of a ribonucleoprotein complex. Said guide RNA can be provided in the form of a pre-guide RNA, a mature guide RNA, or an array containing one or more spacer sequences. The components can also be provided in the form of a DNA or RNA precursor encoding for the Cas13d enzyme and appropriate guide RNA design via an in vitro transcription/translation system to facilitate the generation of the necessary components. These components of the diagnostic kit comprise the "sensor" module.

Such a method can be used to determine if a target RNA is present in a test sample. Such a method can also be used to detect a pathogen, such as a virus or bacteria, or diagnose a disease state, such as a cancer (e.g., wherein the target RNA is specific for a particular microbe or disease). Such a method can also be used to test the purity or identity of an environmental sample or agricultural sample, such as seed or soil.

The "sensor" module will then be challenged with a test sample in the form of RNA. Said test sample can be, but is

not limited to, a genomic DNA sample that is converted into RNA—for example via in vitro transcription—or a direct RNA sample. These samples can be extracted from biological material such as patient samples (e.g., cells, tissue, blood, plasma, serum, saliva, urine, tumor biopsy, cell free DNA or RNA, exosomes, carrier vesicles or particles) and environmental samples (e.g., soil, water, air, seed, or plant samples). In one embodiment to improve diagnostic sensitivity, nucleic acid molecules in the sample are amplified using amplification techniques, such as polymerase chain reaction, recombinase polymerase amplification, loop mediated isothermal amplification, nucleic acid sequence based amplification, strand displacement amplification, rolling circle amplification, ligase chain reaction, and others (e.g., those that use isothermal amplification). Said amplification techniques can optionally employ nucleic acid conversion techniques such as transcription or reverse transcription with randomized primers or targeted primers.

If the sensor module recognizes a cognate target in the test sample, it will activate an RNase activity. This RNase activity can be detected by using a detectable label. In one example, the detectable label includes an RNA linked to a fluorophore and quencher. The intact detectable RNA links the fluorophore and quencher, suppressing fluorescence. Upon cleavage by Cas13d of the detectable RNA, the fluorophore is released from the quencher and displays detectable fluorescent activity.

In another example, cleavage of the reporter RNA releases a non-fluorescent molecule, which can be converted into a visible signal (e.g., visible by eye). In one example, cleavage of the reporter RNA releases a molecule that can be detected via lateral flow. A molecule that can be detected by lateral flow is any molecule that can be bound specifically by antibodies. In one example, the Cas13d protein along with the guide RNA detecting the target and the reporter RNA conjugated to the reporter molecule can be delivered as a single system in the form of a dry test strip. Upon incubation with the test sample, Cas13d protein, a3d, guide RNA and the reporter RNA are rehydrated and in the presence of the RNA target. The Cas13d protein will cleave the reporter RNA, resulting in the migration of the reporter molecule in the test strip via lateral flow and a resulting positive test line signal by binding to antibodies localized there. Such a shelf stable dry detection system not requiring special (frozen) storage could for example prove advantageous in situations where detection of a target RNA or DNA is performed outside a centralized laboratory facility such as a doctor's office, a hospital, a pharmacy, during field work, in an agricultural setting and so forth.

Cas13d is active over a broad range of temperatures making such an application outside of a controlled laboratory environment feasible.

EXAMPLE 16

Cas13d as a Diagnostic for RNA or DNA Transcribed into RNA In Vitro

Cas13d is capable of converting the presence of a matched target RNA into a visible signal in a minimal diagnostic in vitro system. FIGS. 15A-15D (A) Cas13d is converted into an active RNase complex upon binding a target matching the spacer sequence of the guide RNA. It is capable of cleaving gRNA-complementary target RNA or non-complementary bystander RNAs. (B) Cas13d target-dependent RNase activity can be converted into a detectable signal, for example through cleavage of a labeled detector

RNA that is cleaved only in the presence of a target matching the spacer of the Cas13d guide RNA. In this example, the detector RNA contains a fluorophore, 'F', and a quencher 'Q', that abolishes fluorescence. Only upon bystander RNA cleavage is the fluorophore liberated from the quencher and fluorescence is generated. (C) Cas13d from *E. siraeum* produces a visible signal only in the presence of a perfectly matched target and not in the presence of different mismatched targets. (D) Cas13d from *R. flavefaciens* strain XPD3002 produces a visible signal only in the presence of a perfectly matched target and not in the presence of different mismatched targets.

Thus, the system disclosed herein can be part of a lateral flow device (or other solid support), that can be used in diagnostics. The presence of an RNA or DNA sequence can be converted into a signal than can then be detected by conventional lateral flow.

EXAMPLE 17

Cas13d Modifications

FIGS. 16A-16B, Cas13d is amenable to modifications including truncations of regions with low conservation among orthologs. The alignment of Cas13d orthologs in FIG. 16A shows regions with high (green bars) and low (red bars) conservation.

EXAMPLE 18

Targeting Transcripts In Vivo

The ccdB gene in bacterial cells was targeted in vivo using different nCas1 orthologues, *Eubacterium siraeum* nCas1 (Es_nCas1; SEQ ID NO: 1); *Eubacterium siraeum* nCas1 with mutated HEPN domains (Es_nCas1 HEPN -/-; SEQ ID NO: 2); uncultured *Ruminococcus* sp. nCas1 (uncul_nCas1; SEQ ID NO: 3) and uncultured *Ruminococcus* sp. nCas1 with mutated HEPN domains (uncul_nCas1 HEPN -/-; SEQ ID NO: 4) (FIG. 17A). Chemically competent *E. coli* (strain BW25141-DE3) cells were transformed with (1) an arabinose-inducible ccdB plasmid, and (2) a second plasmid (targeting vector) carrying a compatible origin of replication, nCas1 protein coding sequence, and nCas1 guide array containing 4 spacer sequences targeting the ccdB transcript (FIG. 17A).

1. Artificial *Eubacterium siraeum* nCas1 Array Targeting ccdB

a. (SEQ ID NO: 261)
GAACUACACCCGUGCAAAAUGCAGGGGUCUAAAACUAACGGC
 UCUCUCUUUUUAGGUGUAAAC**GAACUACACCCGUGCAAAAU**
GCAGGGGUCUAAAACUUUUAUCUGACAGCAGACGUGCACUGGCC
AGAACUACACCCGUGCAAAAUGCAGGGGUCUAAAACCAUCAU
GCGCCAGCUUUCAUCCCGUAUUGGAACUACACCCGUGCAAAA
UGCAGGGGUCUAAAACUAAUGGCGUUUUUGAUGUCAUUUUCCGG
 GUCCGCGUA
 i. Full 36 nt direct repeat: (SEQ ID NO: 262)
GAACUACACCCGUGCAAAAUGCAGGGGUCUAAAAC

75

-continued

ii. Spacer 1: (SEQ ID NO: 263)
 UAACGGCUCUCUCUUUUUAUAGGUGUAAACC
 iii. Spacer 2: (SEQ ID NO: 264)
 CUUUUAUCUGACAGCAGACGUGCACUGGCCA
 iv. Spacer 3: (SEQ ID NO: 265)
 CAUCAUGCGCCAGCUUUUCAUCCCCGAUUG
 v. Spacer 4: (SEQ ID NO: 266)
 UAAUGGCGUUUUUGAUGUCAUUUUCGCGGUCGCCUGA

2. Artificial Uncultured *Ruminoccus* sp. nCas1 Array Targeting ccdB

a. (SEQ ID NO: 267)
 CUACUACACUGGUGCAAUUUUGCACUAGUCUAAAACUAACGGCU
 CUCUCUUUUUAUAGGUGUAAACCCUACUACACUGGUGCAAUUUG
 CACUAGUCUAAAACUUUAUCUGACAGCAGACGUGCACUGGCCAC
 UACUACACUGGUGCAAUUUUGCACUAGUCUAAAACCAUCAUGCG
 CCAGCUUUUCAUCCCCGAUUGCUACUACACUGGUGCAAUUUGC
 ACUAGUCUAAAACUAAUGGCGUUUUUGAUGUCAUUUUCGCGGUC
 CGC

i. Full 36 nt direct repeat: (SEQ ID NO: 268)
 CUACUACACUGGUGCAAUUUUGCACUAGUCUAAAAC

ii. Spacer 1: (SEQ ID NO: 269)
 UAACGGCUCUCUCUUUUUAUAGGUGUAAACC

iii. Spacer 2: (SEQ ID NO: 270)
 CUUUUAUCUGACAGCAGACGUGCACUGGCCA

iv. Spacer 3: (SEQ ID NO: 271)
 CAUCAUGCGCCAGCUUUUCAUCCCCGAUUG

v. Spacer 4: (SEQ ID NO: 272)
 UAAUGGCGUUUUUGAUGUCAUUUUCGCGGUCGCCUGA

3. Target RNA (ccdB Sequence)

a. (SEQ ID NO: 273)
 AUGCAGUUUUAAGGUUUUACACCUUAUAAAAGAGAGAGCGGUUAUCG
 UCUGUUUGUGGAGUACAGAGUGAUUUUAUUGACACGCCCGGGCG
 ACGGAUGGUGAUCCCCUGGCCAGUGCACGUCUGCUGUCAGAU
 AAGUCUCCCGUGAACUUUACCGGUGGUGCAUAUCGGGGAUGAA
 AGCUGGCGCAUGAUGACCCGAUAUGGCCAGUGUGCCGGUCUC
 CGUUUAUCGGGGAAGAAGUGGCUAUCUCAGCCACCGCGAAAAUG
 ACAUCAAAAACGCCAUUAACCUGAUGUUUUGGGGAAUA

i. spacer 1 target: (SEQ ID NO: 274)
 GGUUUAACCCUUAUAAAAGAGAGCGGUUA

76

-continued

ii. spacer 2 target: (SEQ ID NO: 275)
 UGGCCAGUGCAGCUGCUGCUGCAGAUAAAAG

iii. spacer 3 target: (SEQ ID NO: 276)
 CAUAUCGGGGAUGAAAGCUGGCGCAUGAUG

iv. spacer 4 target: (SEQ ID NO: 277)
 ACCGCGAAAUGACAUCAAAAACGCCAUUA

v. dCas9 target (underlined): (SEQ ID NO: 259)
UUAUCGUCUGUUUGGGAUG

The transformed bacteria were plated on 2 mM arabinose plates to induce ccdB expression and harvested after 24 hours. Total RNA was extracted with Trizol followed by random hexamer-mediated reverse transcription and Taqman probe-based qPCR.

Mutation of the two HEPN domains in each nCas1 protein demonstrate no targeting of ccdB (FIG. 17B, HEPN -/-), while active wild-type nCas1 proteins knock downed the expression of ccdB (FIG. 17B, Es_nCas1; uncul_nCas1). Because dCas9-(catalytically inactive, or “dead” Cas9)-mediated transcriptional repression through binding of the DNA sequence downstream of the transcriptional start site is a current standard in the field, we validate the assay by targeting dCas9 to the ccdB promoter to repress transcription of the ccdB gene. Taken together, this data demonstrates guide RNA-specific degradation of target transcripts in vivo within prokaryotic cells in a HEPN domain-dependent manner.

REFERENCES

Abudayyeh, O.O., Gootenberg, J. S., Essletzbichler, P., Han, S., Joung, J., Belanto, J. J., Verdine, V., Cox, D. B. T., Kellner, M. J., Regev, A., et al. (2017). RNA targeting with CRISPR-Cas13. *Nature*.
 Abudayyeh, O. O., Gootenberg, J. S., Konermann, S., Joung, J., Slaymaker, I. M., Cox, D. B., Shmakov, S., Makarova, K. S., Semenova, E., Minakhin, L., et al. (2016). C2c2 is a single-component programmable RNA-guided RNA-targeting CRISPR effector. *Science* 353, aaf5573.
 Almeida, S., Zhang, Z., Coppola, G., Mao, W., Futai, K., Karydas, A., Geschwind, M. D., Tartaglia, M. C., Gao, F., Gianni, D., et al. (2012). Induced pluripotent stem cell models of progranulin-deficient frontotemporal dementia uncover specific reversible neuronal defects. *Cell Rep* 2, 789-798.
 Anantharaman, V., Makarova, K. S., Burroughs, A. M., Koonin, E. V., and Aravind, L. (2013). Comprehensive analysis of the HEPN superfamily: identification of novel roles in intra-genomic conflicts, defense, pathogenesis and RNA processing. *Biol Direct* 8, 15.
 Batra, R., Nelles, D. A., Pirie, E., Blue, S. M., Marina, R. J., Wang, H., Chaim, I. A., Thomas, J. D., Zhang, N., Nguyen, V., et al. (2017) Elimination of Toxic Microsatellite Repeat Expansion RNA by RNA-Targeting Cas9. *Cell* 170, 899-912 e810.
 Birmingham, A., Anderson, E. M., Reynolds, A., Ilsley-Tyree, D., Leake, D., Fedorov, Y., Baskerville, S., Maksimova, E., Robinson, K., Karpilow, J., et al. (2006). 3' UTR seed matches, but not overall identity, are associated with RNAi off-targets. *Nat Methods* 3, 199-204.

Biswas, M. H. U., Almeida, S., Lopez-Gonzalez, R., Mao, W., Zhang, Z., Karydas, A., Geschwind, M. D., Biernat, J., Mandelkow, E. M., Futai, K., et al. (2016). MMP-9 and MMP-2 Contribute to Neuronal Cell Death in iPSC Models of Frontotemporal Dementia with MAPT Mutations. *Stem Cell Reports* 7, 316-324.

Bland, C., Ramsey, T. L., Sabree, F., Lowe, M., Brown, K., Kyrpides, N. C., and Hugenholtz, P. (2007). CRISPR recognition tool (CRT): a tool for automatic detection of clustered regularly interspaced palindromic repeats. *BMC Bioinformatics* 8, 209.

Bovee, B. F., and Hutton, M. (2008). Refining frontotemporal dementia with parkinsonism linked to chromosome 17: introducing FTDP-17 (MAPT) and FTDP-17 (PGRN). *Arch Neurol* 65, 460-464.

Cheong, C. G., and Hall, T. M. (2006). Engineering RNA sequence specificity of Pumilio repeats. *Proc Natl Acad Sci U S A* 103, 13635-13639.

Chiriboga, C. A., Swoboda, K. J., Darras, B. T., Iannaccone, S. T., Montes, J., De Vivo, D. C., Norris, D. A., Bennett, C. F., and Bishop, K. M. (2016). Results from a phase I study of nusinersen (ISIS-SMN(Rx)) in children with spinal muscular atrophy. *Neurology* 86, 890-897.

Chylinski, K., Le Rhun, A., and Charpentier, E. (2013). The tracrRNA and Cas9 families of type II CRISPR-Cas immunity systems. *RNA Biol* 10, 726-737.

Cox, D. B. T., Gootenberg, J. S., Abudayyeh, O. O., Franklin, B., Kellner, M. J., Joung, J., and Zhang, F. (2017). RNA editing with CRISPR-Cas13. *Science* 358, 1019-1027.

Deltcheva, E., Chylinski, K., Sharma, C. M., Gonzales, K., Chao, Y., Pirzada, Z. A., Eckert, M. R., Vogel, J., and Charpentier, E. (2011). CRISPR RNA maturation by trans-encoded small RNA and host factor RNase III. *Nature* 471, 602-607.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15-21.

Doench, J. G., Petersen, C. P., and Sharp, P. A. (2003). siRNAs can function as miRNAs. *Genes Dev* 17, 438-442.

Doudna, J. A., and Charpentier, E. (2014). Genome editing. The new frontier of genome engineering with CRISPR-Cas9. *Science* 346, 1258096.

Du, Y. C., Gu, S., Zhou, J., Wang, T., Cai, H., Macinnes, M. A., Bradbury, E. M., and Chen, X. (2006). The dynamic alterations of H2AX complex during DNA repair detected by a proteomic approach reveal the critical roles of Ca(2+)/calmodulin in the ionizing radiation-induced cell cycle arrest. *Mol Cell Proteomics* 5, 1033-1044.

East-Seletsky, A., O'Connell, M. R., Burstein, D., Knott, G. J., and Doudna, J. A. (2017). RNA Targeting by Functionally Orthogonal Type VI-A CRISPR-Cas Enzymes. *Mol Cell* 66, 373-383 e373.

East-Seletsky, A., O'Connell, M. R., Knight, S. C., Burstein, D., Cate, J. H., Tjian, R., and Doudna, J. A. (2016). Two distinct RNase activities of CRISPR-C2c2 enable guide-RNA processing and RNA detection. *Nature* 538, 270-273.

Edgar, R. C. (2007). PILER-CR: fast and accurate identification of CRISPR repeats. *BMC Bioinformatics* 8, 18.

Fonfara, I., Richter, H., Bratovic, M., Le Rhun, A., and Charpentier, E. (2016). The CRISPR-associated DNA-cleaving enzyme Cpf1 also processes precursor CRISPR RNA. *Nature* 532, 517-521.

Gasiunas, G., Barrangou, R., Horvath, P., and Siksnys, V. (2012). Cas9-crRNA ribonucleoprotein complex mediates specific DNA cleavage for adaptive immunity in bacteria. *Proc Natl Acad Sci U S A* 109, E2579-2586.

Gilbert, L. A., Horlbeck, M. A., Adamson, B., Villalta, J. E., Chen, Y., Whitehead, E. H., Guimaraes, C., Panning, B., Ploegh, H. L., Bassik, M. C., et al. (2014). Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell* 159, 647-661.

Gilbert, L. A., Larson, M. H., Morsut, L., Liu, Z., Brar, G. A., Torres, S. E., Stern-Ginossar, N., Brandman, O., Whitehead, E. H., Doudna, J. A., et al. (2013). CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell* 154, 442-451.

Grissa, I., Vergnaud, G., and Pourcel, C. (2007). CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35, W52-57.

Hammond, S. M., and Wood, M. J. (2011). Genetic therapies for RNA mis-splicing diseases. *Trends Genet* 27, 196-205.

Heidrich, N., Dugar, G., Vogel, J., and Sharma, C. M. (2015). Investigating CRISPR RNA Biogenesis and Function Using RNA-seq. *Methods Mol Biol* 1311, 1-21.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh, H., and Glass, C. K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* 38, 576-589.

Hsu, P. D., Lander, E. S., and Zhang, F. (2014). Development and applications of CRISPR-Cas9 for genome engineering. *Cell* 157, 1262-1278.

Jackson, A. L., Bartz, S. R., Schelter, J., Kobayashi, S. V., Burchard, J., Mao, M., Li, B., Cavet, G., and Linsley, P. S. (2003). Expression profiling reveals off-target gene regulation by RNAi. *Nat Biotechnol* 21, 635-637.

Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J. A., and Charpentier, E. (2012). A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337, 816-821.

Kar, A., Kuo, D., He, R., Zhou, J., and Wu, J. Y. (2005). Tau alternative splicing and frontotemporal dementia. *Alzheimer Dis Assoc Disord* 19 Suppl 1, S29-36.

Kim, E., Koo, T., Park, S. W., Kim, D., Kim, K., Cho, H. Y., Song, D. W., Lee, K. J., Jung, M. H., Kim, S., et al. (2017). In vivo genome editing with a small Cas9 orthologue derived from *Campylobacter jejuni*. *Nat Commun* 8, 14500.

Liu, L., Li, X., Wang, J., Wang, M., Chen, P., Yin, M., Li, J., Sheng, G., and Wang, Y. (2017). Two Distant Catalytic Sites Are Responsible for C2c2 RNase Activities. *Cell* 168, 121-134 e112.

Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15, 550.

Makarova, K. S., Wolf, Y. I., Alkhnbashi, O. S., Costa, F., Shah, S. A., Saunders, S. J., Barrangou, R., Brouns, S. J., Charpentier, E., Haft, D. H., et al. (2015). An updated evolutionary classification of CRISPR-Cas systems. *Nat Rev Microbiol* 13, 722-736.

Matera, A. G., and Wang, Z. (2014). A day in the life of the spliceosome. *Nat Rev Mol Cell Biol* 15, 108-121.

Naldini, L. (2015). Gene therapy returns to centre stage. *Nature* 526, 351-360.

O'Connell, M. R., Oakes, B. L., Sternberg, S. H., East-Seletsky, A., Kaplan, M., and Doudna, J. A. (2014).

Programmable RNA recognition and cleavage by CRISPR/Cas9. *Nature* 516, 263-266.

Orengo, J. P., Bundman, D., and Cooper, T. A. (2006). A bichromatic fluorescent reporter for cell-based screens of alternative splicing. *Nucleic Acids Res* 34, e148.

Peabody, D. S. (1993). The RNA binding site of bacteriophage MS2 coat protein. *EMBO J* 12, 595-600.

Ran, F. A., Cong, L., Yan, W. X., Scott, D. A., Gootenberg, J. S., Kriz, A. J., Zetsche, B., Shalem, O., Wu, X., Makarova, K. S., et al. (2015). In vivo genome editing using *Staphylococcus aureus* Cas9. *Nature* 520, 186-191.

Samai, P., Pyenson, N., Jiang, W., Goldberg, G. W., Hatoum-Aslan, A., and Marraffini, L. A. (2015). Co-transcriptional DNA and RNA Cleavage during Type III CRISPR-Cas Immunity. *Cell* 161, 1164-1174.

Schena, M., Shalon, D., Davis, R. W., and Brown, P. O. (1995). Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* 270, 467-470.

Schoch, K. M., DeVos, S. L., Miller, R. L., Chun, S. J., Norrbom, M., Wozniak, D. F., Dawson, H. N., Bennett, C. F., Rigo, F., and Miller, T. M. (2016). Increased 4R-Tau Induces Pathological Changes in a Human-Tau Mouse Model. *Neuron* 90, 941-947.

Shendure, J., Balasubramanian, S., Church, G. M., Gilbert, W., Rogers, J., Schloss, J. A., and Waterston, R. H. (2017). DNA sequencing at 40: past, present and future. *Nature advance online publication*.

Shmakov, S., Abudayyeh, O. O., Makarova, K. S., Wolf, Y. I., Gootenberg, J. S., Semenova, E., Minakhin, L., Joung, J., Konermann, S., Severinov, K., et al. (2015). Discovery and Functional Characterization of Diverse Class 2 CRISPR-Cas Systems. *Mol Cell* 60, 385-397.

Sigoillot, F. D., Lyman, S., Huckins, J. F., Adamson, B., Chung, E., Quattrocchi, B., and King, R. W. (2012). A bioinformatics method identifies prominent off-targeted transcripts in RNAi screens. *Nat Methods* 9, 363-366.

Smargon, A. A., Cox, D. B., Pyzocha, N. K., Zheng, K., Slaymaker, I. M., Gootenberg, J. S., Abudayyeh, O. A., Essletzbichler, P., Shmakov, S., Makarova, K. S., et al. (2017). Cas13b Is a Type VI-B CRISPR-Associated RNA-Guided RNase Differentially Regulated by Accessory Proteins Csx27 and Csx28. *Mol Cell* 65, 618-630 e617.

Smith, I., Greenside, P. G., Natoli, T., Lahr, D. L., Wadden, D., Tirosh, I., Narayan, R., Root, D. E., Golub, T. R., Subramanian, A., et al. (2017). Evaluation of RNAi and CRISPR technologies by large-scale gene expression profiling in the Connectivity Map. *PLoS Biol* 15, e2003213.

Swiech, L., Heidenreich, M., Banerjee, A., Habib, N., Li, Y., Trombetta, J., Sur, M., and Zhang, F. (2015). In vivo interrogation of gene function in the mammalian brain using CRISPR-Cas9. *Nat Biotechnol* 33, 102-106.

Treu, L., Kougias, P. G., Campanaro, S., Bassani, I., and Angelidaki, I. (2016). Deeper insight into the structure of the anaerobic digestion microbial community; the biogas microbiome database is expanded with 157 new genomes. *Bioresour Technol* 216, 260-266.

van der Oost, J., Westra, E. R., Jackson, R. N., and Wiedenheft, B. (2014). Unravelling the structural and mechanistic basis of CRISPR-Cas systems. *Nat Rev Microbiol* 12, 479-492.

Wang, Y., Liu, J., Huang, B. O., Xu, Y.M., Li, J., Huang, L. F., Lin, J., Zhang, J., Min, Q. H., Yang, W. M., et al. (2015). Mechanism of alternative splicing and its regulation. *Biomed Rep* 3, 152-158.

Wright, A. V., and Doudna, J. A. (2016). Protecting genome integrity during CRISPR immune adaptation. *Nat Struct Mol Biol* 23, 876-883.

Yang, X., Zou, P., Yao, J., Yun, D., Bao, H., Du, R., Long, J., and Chen, X. (2010). Proteomic dissection of cell type-specific H2AX-interacting protein complex associated with hepatocellular carcinoma. *J Proteome Res* 9, 1402-1415.

Yosef, I., Goren, M. G., and Qimron, U. (2012). Proteins and DNA elements essential for the CRISPR adaptation process in *Escherichia coli*. *Nucleic Acids Res* 40, 5569-5576.

Zalatan, J. G., Lee, M. E., Almeida, R., Gilbert, L. A., Whitehead, E. H., La Russa, M., Tsai, J. C., Weissman, J. S., Dueber, J. E., Qi, L. S., et al. (2015). Engineering complex synthetic transcriptional programs with CRISPR RNA scaffolds. *Cell* 160, 339-350.

Zetsche, B., Gootenberg, J. S., Abudayyeh, O. O., Slaymaker, I. M., Makarova, K. S., Essletzbichler, P., Volz, S. E., Joung, J., van der Oost, J., Regev, A., et al. (2015). Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. *Cell* 163, 759-771.

Zetsche, B., Strecker, J., Abudayyeh, O. O., Gootenberg, J. S., Scott, D. A., and Zhang, F. (2017). A Survey of Genome Editing Activity for 16 Cpf1 orthologs. *bioRxiv*.

Zhang, J., Graham, S., Tello, A., Liu, H., and White, M. F. (2016). Multiple nucleic acid cleavage modes in divergent type III CRISPR systems. *Nucleic Acids Res* 44, 1789-1799.

Zhang, Y., Pak, C., Han, Y., Ahlenius, H., Zhang, Z., Chanda, S., Marro, S., Patzke, C., Acuna, C., Covy, J., et al. (2013). Rapid single-step induction of functional neurons from human pluripotent stem cells. *Neuron* 78, 785-798.

In view of the many possible embodiments to which the principles of the disclosure may be applied, it should be recognized that the illustrated embodiments are only examples of the invention and should not be taken as limiting the scope of the invention. Rather, the scope of the invention is defined by the following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

SEQUENCE LISTING

The patent contains a lengthy "Sequence Listing" section. A copy of the "Sequence Listing" is available in electronic form from the USPTO web site (<https://seqdata.uspto.gov/?pageRequest=docDetail&DocID=US11316812B2>). An electronic copy of the "Sequence Listing" will also be available from the USPTO upon request and payment of the fee set forth in 37 CFR 1.19(b)(3).

We claim:

1. A non-naturally occurring or engineered clustered regularly interspaced short palindromic repeats (CRISPR)-associated (Cas) system for targeting one or more target RNA molecules, comprising:

- (a) at least one Cas13d protein comprising at least 95% sequence identity to SEQ ID NO: 42, or a nucleic acid molecule encoding the at least one Cas13d protein; and
- (b) at least one CRISPR-Cas system guide RNA (gRNA) comprising one or more direct repeat (DR) sequences and one or more spacer sequences, which hybridizes with the one or more target RNA molecules, or at least one nucleic acid molecule encoding the gRNA,

whereby the Cas13d protein forms a complex with the gRNA, wherein the gRNA directs the complex to the one or more target RNA molecules and targets the one or more target RNA molecules.

2. An isolated cell comprising the system of claim 1.

3. A ribonucleoprotein complex comprising the system of claim 1, wherein the ribonucleoprotein complex comprises the (a) at least one Cas13d protein comprising at least 95% sequence identity to SEQ ID NO: 42 and the (b) at least one CRISPR-Cas system gRNA comprising one or more DR sequences and one or more spacer sequences, which hybridizes with the one or more target RNA molecules,

wherein the gRNA directs the ribonucleoprotein complex to the one or more target RNA molecules and targets the one or more target RNA molecules.

4. The system of claim 1, wherein targeting one or more target RNA molecules comprises one or more of cutting the one or more target RNA molecules, nicking the one or more target RNA molecules, increasing expression of the one or more target RNA molecules as compared to an amount of target RNA in a corresponding cell or sample not contacted with the at least one Cas13d protein, decreasing expression of the one or more target RNA molecules as compared to an amount of target RNA in a corresponding cell or sample not contacted with the at least one Cas13d protein, visualizing or detecting the one or more target RNA molecules, labeling the one or more target RNA molecules, capturing molecules bound or tethered to the one or more target RNA molecules, enriching the one or more target RNA molecules, depleting the one or more target RNA molecules, editing the one or more target RNA molecules, trafficking the one or more target RNA molecules, splicing or perturbing the splicing of the one or more target RNA molecules, and masking the one or more target RNA molecules.

5. The system of claim 1, wherein the at least one Cas13d protein comprises at least 96% sequence identity to SEQ ID NO: 42.

6. The system of claim 1, wherein the at least one Cas13d protein comprises at least 97% sequence identity to SEQ ID NO: 42.

7. The system of claim 1, wherein the at least one Cas13d protein comprises at least 98% sequence identity to SEQ ID NO: 42.

8. The system of claim 1, wherein the at least one Cas13d protein comprises at least 99% sequence identity to SEQ ID NO: 42.

9. The system of claim 1, wherein the at least one Cas13d protein comprises SEQ ID NO: 42.

10. The system of claim 1, wherein the at least one Cas13d protein further comprises one or more subcellular localization signals, effector domains, purification tags, affinity tags, or combinations thereof.

11. The system of claim 1, wherein the at least one gRNA comprises one or more DR sequences comprising at least 95% sequence identity to SEQ ID NO: 135.

12. The system of claim 1, wherein the at least one gRNA comprises one or more DR sequences comprising at least 96% sequence identity to SEQ ID NO: 135.

13. The system of claim 1, wherein the at least one gRNA comprises one or more DR sequences comprising at least 97% sequence identity to SEQ ID NO: 135.

14. The system of claim 1, wherein the at least one gRNA comprises one or more DR sequences comprising at least 98% sequence identity to SEQ ID NO: 135.

15. The system of claim 1, wherein the at least one gRNA comprises one or more DR sequences comprising at least 99% sequence identity to SEQ ID NO: 135.

16. The system of claim 1, wherein the at least one gRNA comprises one or more DR sequences comprising SEQ ID NO: 135.

17. The system of claim 1, wherein the at least one gRNA comprises a nucleic acid aptamer.

18. The system of claim 17 wherein the nucleic acid aptamer inserted into the gRNA comprises MS2, PP7, or Q β .

19. The system of claim 1, wherein the at least one gRNA comprises modified nucleotides.

20. The isolated cell of claim 2, wherein the cell is a eukaryotic cell or a bacterial cell.

21. The isolated cell of claim 20, wherein the eukaryotic cell is a mammalian cell.

22. The isolated cell of claim 20, wherein the eukaryotic cell is a plant cell.

23. The system of claim 1, wherein the at least one Cas13d protein further comprises a nuclear localization signal (NLS).

24. The system of claim 23, wherein the NLS comprises SEQ ID NO: 256 or SEQ ID NO: 258.

25. The system of claim 1, wherein the at least one Cas13d protein further comprises a nuclear export signal (NES).

26. The system of claim 25, wherein the NES comprises an adenovirus type 5 E1B nuclear export sequence, an HIV nuclear export sequence, a MAPK nuclear export sequence, or a PTK2 nuclear export sequence.

27. The system of claim 1, wherein the at least one Cas13d protein further comprises an effector domain that can cleave RNA, edit a nucleotide, edit a ribonucleotide, methylate RNA, or demethylate RNA.

28. The system of claim 1, wherein the at least one Cas13d protein is catalytically inactive for ribonuclease activity.

29. The system of claim 28, wherein the catalytically inactive Cas13d protein comprises one or more mutated HEPN domains.

30. The system of claim 28, wherein the catalytically inactive Cas13d protein comprises a mutated HEPN1 domain or a mutated HEPN2 domain.

* * * * *