

US010160795B2

# (12) United States Patent

# Sullivan et al.

# (54) NEUTRALIZING ANTIBODIES TO EBOLA VIRUS GLYCOPROTEIN AND THEIR USE

- (71) Applicants: The United States of America, as represented by the Secretary, Department of Health and Human Services, Washington, DC (US); Institute for Research in Biomedicine, Bellinzona (CH); The United States of America as represented by the Secretary of the Army, Washington, DC (US); Humabs BioMed SA, Bellinzona (CH)
- (72) Inventors: Nancy Sullivan, Bethesda, MD (US);
  Sabue Mulangu, Bethesda, MD (US);
  Davide Corti, Bellinzona (CH);
  Antonio Lanzavecchia, Bellinzona (CH); Barney Graham, Bethesda, MD (US); Jean-Jacques
  Muyembe-Tamfun, Kinshasa (CD);
  John Trefry, Frederick, MD (US);
  Julie Ledgerwood, Bethesda, MD (US); Daphne Stanley, Bethesda, MD (US)
- (73) Assignees: The United States of America, as represented by the Secretary, Department of Health and Human Services, Washington, DC (US); Institute for Research in Biomedicine, Bellinzona (CH); The United States of America as represented by the Secretary of the Army, Washington, DC (US); Humabs BioMed SA, Bellinzona (CH)
- (\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 25 days.
- (21) Appl. No.: 15/526,661
- (22) PCT Filed: Nov. 13, 2015
- (86) PCT No.: PCT/US2015/060733
  § 371 (c)(1),
  (2) Date: May 12, 2017
- (87) PCT Pub. No.: WO2016/077789PCT Pub. Date: May 19, 2016

# (65) **Prior Publication Data**

US 2018/0291088 A1 Oct. 11, 2018

# **Related U.S. Application Data**

(60) Provisional application No. 62/087,087, filed on Dec. 3, 2014, provisional application No. 62/080,094, filed on Nov. 14, 2014.

# (10) Patent No.: US 10,160,795 B2

# (45) **Date of Patent:** Dec. 25, 2018

(51)	Int. Cl.	
. ,	C07K 16/10	(2006.01)
	A61K 39/00	(2006.01)
	A61K 39/395	(2006.01)
	A61K 45/06	(2006.01)
	C07K 16/00	(2006.01)
	G01N 33/569	(2006.01)

- (52) U.S. Cl.
  - CPC ...... **C07K 16/10** (2013.01); **G01N 33/56983** (2013.01); *A61K 2039/505* (2013.01); *C07K 2317/52* (2013.01); *C07K 2317/56* (2013.01); *C07K 2317/565* (2013.01); *C07K 2317/567* (2013.01); *C07K 2317/72* (2013.01); *C07K 2317/732* (2013.01); *C07K 2317/76* (2013.01); *G01N 2333/08* (2013.01); *G01N 2800/26* (2013.01)

# (56) References Cited

#### U.S. PATENT DOCUMENTS

6,875,433	B2	4/2005	Hart et al.
8,513,391	B2	8/2013	Jones et al
9,097,713	B2	8/2015	Dye et al.

# FOREIGN PATENT DOCUMENTS

WO	WO 2004/018649	3/2004
WO	WO 2011/071574	6/201
WO	WO 2016/075546	5/201

## OTHER PUBLICATIONS

International Search Report and Written Opinion for PCT/IB2015/ 0023432, mailed by the European Patent Office acting as International Searching Authority dated Feb. 23, 2016 (13 pages).

Lee, et al. "Neutralizing ebolavirus: structural insights into the envelope glycoprotein and antibodies targeted against it." *Current Opinion in Structural Biology* 19, No. 4 (2009): 408-417.

Qiu, et al. "Characterization of Zaire ebolavirus glycoproteinspecific monoclonal antibodies." *Clinical Immunology* 141, No. 2 (2011): 218-227.

Qiu, et al. "Successful treatment of Ebola virus-infected cynomoigus macaques with monoclonal antibodies." *Science Translational Medicine* 4, No. 138 (2012): 138ra81-138ra81.

Primary Examiner - Barry A Chestnut

(74) Attorney, Agent, or Firm — Klarquist Sparkman, LLP

# (57) **ABSTRACT**

Neutralizing antibodies and antigen binding fragments that specifically bind to Ebola virus glycoprotein are disclosed. Nucleic acids encoding these antibodies, vectors and host cells are also provided. Methods for detecting Ebola virus using the antibodies and antigen binding fragments are disclosed. The antibodies, antigen binding fragments, nucleic acids, and vectors, can be used, for example, to prevent and/or treat Ebola virus infection in a subject.

#### 39 Claims, 33 Drawing Sheets

# Specification includes a Sequence Listing.



FIG. 1B







**FIG. 1C** 



FIG. 2C

	VH	Identity %	Ð	JH	VL	Identity %	JL	HCDR3 length	IgG subclass
EVB100	V4-59*01	94.74%	D3-16'02	J4*02	VL3-1*01	92.11%	J2*01	19	lgG3
EVB114	V3-13*01	87.37%	D6-19*01	J4*02	VK1-27*01	92.83%	J4*01	13	lgG1
EVB165	V3-7*01	89.93%	D2-2*02	J4*02	VK1-39*01	89.96%	J1*01	21	lgG1
EVB166	V1-69*04	92.01%	D3-22*01	J6*03	VK3-20*01	97.16%	J3*01	16	lgG3

FIG. 2D	EVB100 VH UCA	QVOLQESGPGLVKPSETLSLTCTVSGGSISSYYWSWIROPPGKGLEWIGYLYYSGSTNYNPSLKSRVTIS 70
	Hg	02
	gH-FR	
	gH-FR1-2-4	
	ЧS	
	EVB100 VH UCA	VDTSKNOFSLKLSSVTAADTAVYYCARASRSYVWGSYRPTAFDYWGOGTLVTVSS 125
	ВH	······································
	gH-FR	
	gH-FR1-2-4	TR I D
	84	TRID
FIG. 2E	EVB100 VL UCA	SYELTQPSVSVSPGQTASITCSGDKLGDKYACWYQOKPGQSPVLVIYQDSKRPSGIPERFSGSNSGNTA 70
	9L	20
	gl-FR	70 N
	ಶ	L
	EVB100 VL UCA	TLTISGTOAMDEADYYCOAWDSTVYFGGGTKLTVL 105
	ցե	
	gt-FR	
	st	······ST·····ST·······················

FIG. 2F	EVB114 VH UCA	EVOLVESGGGLVOPGGSLRLSCAASGETESSYDMHWVROATGKGLEWVSAJGTAGDTYYPGSVKGRFTIS 70
	Нĝ	0.2
	gH-FR	
	gH-FR3	
	gH-FR1-2-4	
	Hs	
	EVB114 VH UCA	RENAKNSLYLOMNSLRAGDTAVYYCARSDRGVAGLEDYWGOGTLVTVSS 119
	HÔ	
	gH-FR	· · · · · · · · · · · · · · · · · · ·
	gH-FR3	
	gH-FR1-2-4	STIVSS119
	Hs	sTIXSIX
FIG. 2G	EVB114 VK UCA	DIOMTOSPSSLSASVGDRVTITCRASOGISNYLAWYQOKPGKVPKLLIYAASTLOSGVPSRFSGSGSGTD 70
	<u></u> д	70
	glFR	
	¥	
	EVB114 VK UCA	FTLTISSLOPEDVATYYCOKYNSAPLTFGGGTKVEIK 107
	<u>_</u> 6	
	gL-FR	
	SK	











				Analyte		
		KZ52	EVB100	13c6	EVB114	Control
	KZ52	83.4%	73.1%	7.9%	-6.1%	NB
tito	EVB100	85.9%	92.0%	-2.7%	-4.4%	NB
mpe	13c6	3.8%	12.8%	104.0%	84.4%	NB
ပိ	EVB114	14.8%	8.1%	102.5%	97.4%	NB
	Control	28.1%	5.2%	-5.0%	-10.8%	NB













FIG. 6



































MPV

FIG. 10C

RDW



























FIG. 12 EBOV mAbs neutralization at IC50 concentration in

FIG. 13









FIG. 15





# FIG. 17A

IP/WB



# FIG. 17B

**Direct WB** 





5

30

# NEUTRALIZING ANTIBODIES TO EBOLA VIRUS GLYCOPROTEIN AND THEIR USE

### CROSS REFERENCE TO RELATED APPLICATIONS

This is the U.S. National Stage of International Application No. PCT/US2015/060733, filed Nov. 13, 2015, which was published in English under PCT Article 21(2), which in turn claims the benefit of U.S. Provisional Application No. 62/087,087, filed Dec. 3, 2014, and U.S. Provisional Application No. 62/080,094, filed Nov. 14, 2014; each of the provisional applications is incorporated by reference herein in its entirety.

### PARTIES TO A JOINT RESEARCH AGREEMENT

This invention was made under Research Collaboration Agreement No. 2007-0166 between the National Institutes<sup>20</sup> of Health National Institute of Allergy and Infectious Disease and Institute for Research in Biomedicine.

### FIELD OF THE DISCLOSURE

This relates to monoclonal antibodies and antigen binding fragments that specifically bind to Ebola virus (EBOV) glycoprotein (GP) and their use, for example, in methods of treating or preventing EBOV infection or EBOV disease (EVD) in a subject.

#### BACKGROUND

EVD is a disease in humans, chimpanzees, and monkeys, caused by infection with EBOV. This virus was first recog-<sup>35</sup> nized in Zaire, Africa in 1976. EBOV is a member of the Filoviridae family of RNA viruses and causes a severe hemorrhagic fever with a high mortality rate. For example, infection with the Ebola virus Zaire (ZEBOV) strain of the virus is associated with a mortality rate of up to 90% in <sup>40</sup> humans. Currently, there are no licensed vaccines or therapeutics approved for human use.

An enveloped virus, EBOV hides from humoral recognition behind a wide array of protective mechanisms. EBOV GP, the major envelope glycoprotein of EBOV is approxiter approximately 165 kD in size. During infection proteases of the host cell cleave a precursor of GP, termed  $GP_0$ , into  $GP_1$  and  $GP_2$ . GP<sub>1</sub> is an integral membrane protein, while GP<sub>1</sub> protrudes from the mature virus. Together GP<sub>1</sub> and GP<sub>2</sub> make up the EBOV envelope spike, which is a target for neutralizing antibodies. Although certain EBOV neutralizing antibodies that bind to the EBOV GP have been identified, there is a need to develop additional neutralizing antibodies for EBOV with varying EBOV GP recognition profiles and increased neutralization potency. 55

#### SUMMARY

Isolated monoclonal antibodies and antigen binding fragments that specifically bind to an epitope on EBOV GP are 60 provided herein. The antibodies and antigen binding fragments can neutralize EBOV infection.

In some embodiments, the antibody or antigen binding fragment comprises a heavy chain variable region  $(V_H)$  comprising a HCDR1, a HCDR2, and a HCDR3 of the  $V_H$  65 set forth as SEQ ID NO: 1 (EVB114 VH) and/or a light chain variable region  $(V_L)$  comprising a LCDR1, a LCDR2, and a

LCDR3 of the  $V_L$  set forth as SEQ ID NO: 2 (EVB114 VL) and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment comprises a  $V_H$  comprising a HCDR1, a HCDR2, and a HCDR3 of the  $V_H$  set forth as SEQ ID NO: 3 (EVB100 VH) and/or a  $V_L$  comprising a LCDR1, a LCDR2, and a LCDR3 of the  $V_L$  set forth as SEQ ID NO: 4 (EVB100 VL) and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment comprises a  $V_H$  and a  $V_L$  comprising the amino acid sequences set forth as SEQ ID NOs: 1 and 2, respectively, or SEQ ID NOs: 3 and 4, respectively.

In additional embodiments, the glycosylation (for example, fucosylation) or sequence of a disclosed antibody <sup>15</sup> or antigen binding fragment can be altered compared to that observed in nature. For example the glycosylation or sequence of a disclosed of the antibody or antigen binding fragment can be altered compared to that of native antibodies to increase half-life, antibody-dependent cell-mediated <sup>20</sup> cytotoxic activity, and/or EBOV neutralization or EBOV GP binding profile.

Also disclosed are compositions including the antibodies and antigen binding fragments, nucleic acids encoding the antibodies and antigen binding fragments, expression vectors comprising the nucleic acids, and isolated host cells that comprise the nucleic acids. In several embodiments, the nucleic acid molecule encoding a disclosed antibody or antigen binding fragment can be a cDNA molecule that encodes the antibody or antigen binding fragment. In additional embodiments, the nucleic acid molecule can be a bicistronic expression construct encoding the antibody or antigen binding fragment.

Surprisingly, the disclosed antibodies and antigen binding fragments potently neutralize EBOV infection in vitro and in vivo. Accordingly, a method is disclosed for treating or preventing an EBOV infection (e.g., ZEBOV infection) in a subject comprising administering a therapeutically effective amount of one or more of the disclosed antibodies or antigen binding fragments to the subject, for example to a subject at risk of or having an EBOV infection.

The antibodies, antigen binding fragments, nucleic acid molecules, vectors, and compositions disclosed herein can be used for a variety of additional purposes, such as for detecting an EBOV infection or diagnosing EVD in a subject, or detecting EBOV GP in a sample.

The foregoing and other features and advantages of this disclosure will become more apparent from the following detailed description of several embodiments which proceeds with reference to the accompanying figures.

#### BRIEF DESCRIPTION OF THE FIGURES

FIGS. 1A-1D are a set of graphs concerning isolation of antigen-specific monoclonal antibodies from an Ebola virus
55 disease survivor. (1A) Plasma obtained from two human survivors, an uninfected human donor and a non-human primate (NHP) vaccinated against EBOV GP were serially diluted and analyzed by GP ELISA, A450 (n=1). (1B) Lentivirus particles expressing luciferase and bearing
60 EBOV GP were incubated in the presence of heat inactivated serum for 1 hour prior to addition to HEK293T. Infection was determined by measuring relative luminescence (RLU) after 3 days. Infection %=(RLU with serum/RLU without serum)×100% (n=3). (1C) Immortalized B cell supernatants
65 isolated from Survivor 1 were screened by EBOV GP ELISA A450 (n=1). (1D) Immortalized B cell supernatants from (1C) were diluted 1:50, incubated with Lentivirus

particles pseudotyped with EBOV GP and infection determined as in (1B). Infection %=(RLU with supernatant/RLU without supernatant)×100% (n=1).

FIGS. 2A-2I are a set of graphs and tables concerning characterization of purified EBOV GP monoclonal antibod- 5 ies. (2A) EBOV GP ELISA in the presence of purified monoclonal antibodies as indicated, A450. (2B) Lentivirus particles pseudotyped EBOV GP particles were incubated with increasing amounts of purified monoclonal antibodies and infection measured as in FIG. 1B. Infection %=(RLU with antibody/RLU without antibody) $\times 100\%$  (n=3). (2C) V gene usage, sequence analysis and IgG subclass of antibodies from Survivor 1. (2D-2G) Amino acid sequence of EVB100, EVB114 and variants descended from a putative unmutated common ancestor (UCA) for heavy and light 15 chains. Shaded regions represent complementary determination regions 1-3. (2H) and (2I) Binding to EBOV GP expressed on the surface of MDCK-SIAT cells by different EVB100 (2H) and EVB114 (2I) versions in which all or subsets of somatic mutations in the wild type sH. sL 20 (EVB100) or sK (EVB114) chain were reverted to the germline sequence. Shown is the ratio between the EC50 values of the variants and EC50 values of the wild-type sH/sL (EVB100) or sH/sK (EVB114). UCA, unmutated common ancestor; gH or gL, germline V-gene revertants of 25 sH, sL, or sK in which the HCDR or LCDR3 are mature; gH-FR or gL-FR, germline V347 gene revertants of sH, sL or sK in which the HCDRs or LCDRs are mature; gH-FR1-2-4, germline V-gene revertants of sH in which the HCDRs and HFR3 are mature; gH-FR3, germline V-gene revertants 30 of sH in which the HCDRs and HFR1, HFR2 and HFR4 are mature; wild type, somatically mutated are sH, sL, or sK. EC50 ratio values above 100 indicate lack of detectable binding

FIGS. 3A-3D are a set of graphs and tables concerning the 35 binding region and effector function of EBOV GP specific antibodies. (3A) Inhibition of binding of biotinylated EVB114 (left) and EVB100 (right) to GP-expressing MDCK-SIAT cells by pre-incubation with increasing amounts of homologous or heterologous unlabeled antibod- 40 ies. Shown is the percentage of binding of biotinylated antibodies as measured by flow cytometry using fluorophore-conjugated streptavidin. (3B) and (3C) Biolayer interferometry competitive binding assay to soluble EBOV GP using EVB100, EVB114, KZ52, 13C6 and isotype negative 45 control. Biosensors were preloaded with GP followed by the competitor and analyte antibodies as indicated. Analyte binding curves (3B) and quantitated % inhibition (3C) are reported (n=3). (3D) Antibody-dependent cell-mediated cytotoxicity (ADCC) assay was determined at 31.6 ng/mL of 50 EVB100, EVB114 (n=3), control antibody or derivative antibodies with LALA mutations that abrogate Fc365 mediated killing (n=1).

FIGS. **4**A-**4**I are a set of graphs showing that passive transfer of EBOV GP-specific antibodies can inhibit EBOV 55 disease. **(4**A) Experimental challenge. Animals were challenged with a lethal dose of EBOV GP on Day 0 and given injections of antibody totaling 50 mg/kg at 24, 48 and 72 hours post-exposure. Surviving animals were euthanized at the conclusion of the study (Day 28). Challenge data from 60 monoclonal antibody EVB114/EVB100 mixture (**4**B-**4**E), or EVB114 monotherapy (**4**F-**4**I). Treatment animal in black, untreated control in grey. **(4B)** and **(4F)** Ebola GP specific ELISA titer (EC90). **(4C)** and **(4G)** Viremia in blood by qRT-PCR expressed as genome equivalents (ge) per mL. 65 (**4**D) and (**4**H) Survival. **(4**E) and **(4**I) Selected hematologic and chemistry data. Platelets (PLT), alanine transaminase

(ALT), creatinine (CRE). "nt" is used to indicate data concerning the no treatment (control) animal.

FIG. **5** is a graph illustrating inhibition of EBOV Makona variant by EVB100 and EVB114. Lentivirus particles bearing GPs from EBOV Makona variant were incubated with serially diluted EVB100, EVB114 or isotype control. Infection measured as in FIG. **2**B (n=3).

FIGS. **6-8** are a set of graphs showing clinical data from 10 the EBOV challenge study using passive transfer of a combination of EVB114 and EVB100. "nt" is used to indicate data concerning the no treatment (control) animal.

FIGS. **9-11** are a set of graphs showing clinical data from the EBOV challenge study using passive transfer of a monotherapy using EVB114. "nt" is used to indicate data concerning the no treatment (control) animal.

FIG. **12** shows a graph illustrating the neutralization properties of the EVB100, EVB114, EVB165, and EVB166 antibodies in the presence of soluble GP (sGP), which is believed to interfere with the natural immune response to EBOV in human subjects. sGP is a GP splice variant that lacks a transmembrane domain and is therefore secreted from infected cells. Pseudotyped lentiviral vectors expressing EBOV GP were incubated with the IC50 concentration of each antibody (as shown in FIG. **2**B) and sGP prior to infection of 293T cells, and infection inhibition was calculated as a percent of infection in the absence of antibody.

FIG. **13** shows a Western blot indicating that the EVB100, EVB114, EVB165, and EVB166 antibodies can immunoprecipitate several different forms of EBOV GP, including the GP<sub>0</sub>, GP<sub>1</sub>, GP<sub>2</sub>, pre-GP<sub>er</sub>, and GP<sub>CatL</sub> forms of GP. KZ52 was used as a positive control.

FIGS. **14**A and **14**B are a set of diagrams illustrating EBOV GP and regions thereof (FIG. **4**A) and several deletion mutants of EBOV GP used herein (FIG. **4**B).

FIGS. **15** and **16** are a set of Western blots illustrating the ability of the EVB100, EVB114, EVB165, and EVB166 antibodies to immunoprecipitate the GP dMUC and GP dGP2 deletion mutants illustrated in FIG. **4**B.

FIGS. **17**A and **17**B are a set of Western blots illustrating the ability of the EVB100, EVB114, EVB165, and EVB166 antibodies to immunoprecipitate the sGP form of EBOV GP (FIG. **7**A) and to recognize the sGP form by direct Western blot (FIG. **7**B).

FIG. **18** is a set of graphs illustrating the cross-species neutralization properties of the EVB100, EVB114, EVB165, and EVB166 antibodies. Pseudotyped lentiviral vectors expressing EBOV GP from the Bundibugyo or Sudan EBOV strains were incubated with antibody prior to infection of 293T cells, and infection inhibition was calculated as a percent of infection in the absence of antibody.

#### SEQUENCES

The nucleic and amino acid sequences are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. The Sequence Listing is submitted as an ASCII text file in the form of the file named "Sequence.txt" (~60 kb), which was created on May 10, 2017, which is incorporated by reference herein.
SEQ ID NO: 1 is the amino acid sequence of the V<sub>H</sub> of the EVB114 mAb. EVQLVESGGGLIQPGGSLRLSCAASgfalrmydMHWVRQTIDKRLEWVSAvgpsgdtYYADSVKGRFAVSR

ENAKNSLSLQMNSLTAGDTAIYYC vrsdrgvaglfds WGQGILVTVSS

SEQ ID NO: 2 is the amino acid sequence of the V\_L of the EVB114 mAb. DIQMTQSPSSLSASVGDRITITCRAS qafdnyVAWYQQRPGKVPKLLISaasALHAGVPSRFSGSGSGTHF

TLTISSLQPEDVATYYC**qnynsaplt**FGGGTKVEIK

SEQ ID NO: 3 is the amino acid sequence of the  $V_H$  of the EVB100 mAb.  $\rm QVQLQESGPGLVKPSDTLSLTCTVSggslssfyWSWIRQPPGKGLEWIGYiyysgspNYSPSLESRVTMSV$ 

DTTRNQISLKLDSVTAADTAVYYCvrasrsyywgsyrptafdsWGQGTLVTVSS

SEQ ID NO: 4 is the amino acid sequence of the  $V_L$  of the EVB100 mAb. SYELTQPLSVSVSPGQTAIFTCSGD**nlgdky**VCWFQQRPGQSPMLLIY**qdn**KRPSGIPERFSGSNSGNTAT

LTISGTQSTDEADYYCqtwdstvvFGGGTKLTVL

SEQ ID NO: 5 is the amino acid sequence of the  $\rm V_{H}$  of the EVB166 mAb. QVQLVQSGAEVKKPGSSVKVSCKTSggtlsnyaISWVRQAPGQGLEWMGGtiptlgmsTYAPNFQGRVAIT

ADKSTSTAYMELSSLRSDDTAVYYCatmgsadtsfyfymdvWGKGTTVTVSS

SEQ ID NO: 6 is the amino acid sequence of the V<sub>L</sub> of the EVB166 mAb. EIVLTQSPGTLSLSPGERATLSCRASqsvsssyLAWYQQKPGQAPRLLIYgtsSRATGIPDRFSGSASGTD

FTLTISRLEPEDFAVYYCqqyayspftFGPGTKVDIK

 $\verb+tctcagctGTGGGTCCTTCTGGTGACACCtactatgcagactccgtgaagggccgattcgccgtctccaga$ 

gagaatgccaagaactccttgtctcttcagatgaacagcctgacagccggggacacggctatatactattg

 ${\tt t} {\tt GTAAGGTCTGACCGAGGAGTGGCTGGCCTTTTTGACAGC} {\tt t} ggggccagggaatcctggtcaccgtctctt$ 

cag

SEQ ID NO: 8 is an exemplary nucleotide sequence encoding the  $\rm V_L$  of the EVB114 mAb. gacatccagatgacccagtctccatcatccctgtctgcatctgtgggagacagaatcaccatcacttgccg

 $ggcgagt {\tt CAGGCCTTTGACAATTAT} gtagcctggtatcaacagagaccagggaaggttcctaagctcctga$ 

 ${\tt tctct} {\tt GCTGCATCC} {\tt gctttgcacgcaggggtcccatctcgcttcagcggcagtggctctgggacacatttc}$ 

 $actctcaccatcagcagcctgcagcctgaagatgttgcaacttattactgt {\tt CAAAACTATAACAGTGCCCC}$ 

**GCTCACT**ttcggcggagggaccaaggtggagatcaaac

SEQ ID NO: 9 is an exemplary nucleotide sequence encoding the  ${\rm V}_H$  of the EVB100 mAb. caggtgcagctgcaggagtcggggcccaggactggtgaagccttcggataccctgtccctcacctgtactgt

ctct GGTGGCTCCCTCAGTAGTTTCTACtqqaqctqqatccqqcaqccccccaqqqaaqqqactqqaqtqqa

 ${\tt ttgggtat} {\tt ATCTATTACAGTGGGAGCCCC} {\tt actacagcccctccctcgagagtcgagtcaccatgtcagta}$ 

gacacgaccaggaaccagatctccctgaagttggactctgtgaccgcggcggacacggccgtgtattactg

t GTGAGAGCCTCCCGAAGTTACTATTGGGGGGAGTTATCGCCCAACGGCTTTTGACTCCtggggccagggaa

ccctggtcaccgtctcctcag

SEQ ID NO: 10 is an exemplary nucleotide sequence encoding the  $\rm V_L$  of the EVB100 mAb. tcctatgagctgactcagccactctcagtgtccctggtgtccccaggccagacagccatcttcacctgctctgg

agat AATTTGGGGGGATAAGTAT gtttgctggtttcaacagaggccaggccagtcccctatgctgctcatct

atCAAGACAAT aagcggccctcggggatccctgagcgattctctggctccaactctgggaacacagccact

ctgactatcagcggggacccagtctaccagatgaggctgactattactgt CAGACGTGGGACAGCACCGTGGT

 ${\tt G} {\tt ttcggcggagggagccaaactgaccgtcctgg}$ 

SEQ ID NO: 11 is an exemplary nucleotide sequence encoding the  ${\rm V}_H$  of the EVB166 mAb. caggtccagctggtgcagtctggggctgaggtgaagaagcctgggtcctcggtgaaagtctcctgcaagac

-continued

ttct**GGAGGCACCCTCAGCAACTATGCT**atcagctgggtgcgacaggcccctggacaagggcttgagtgga

tgggaggcACCATTCCTACCCTTGGTATGTCCacctacgcaccgaacttccagggcagagtcgcgattacc

 ${\tt gcggacaaatccacgagcacagcctacatggagttgagtagtctgaggtctgacgacacggccgtttatta$ 

tgt GCGACTATGGGCAGTGCGGACACTAGTTTCTACTTCTACATGGACGTC tggggcaaagggaccacgg

tcaccgtctcctcag

SEQ ID NO: 12 is an exemplary nucleotide sequence encoding a variant  $V_L$  of the EVB166 mAb that includes a K104T substitution. Gaaattgtgttgacgcagtctccaggcaccctgtctttgtctccaggggagagagccaccctctcctgcag

ggccagt**CAGAGTGTTAGTAGCAGCTAC**ttagcctggtaccagcagaaacctggccaggctcccagactcc

 $tt cactet caccat caq caq a ctq q a q ctt q a q a ttt tq caq tq t a tt a ctq t { {\it CAGCAGTATGCTTACTC}}$ 

ACCATTCACTttcqqccctqqqaccacaqtqqatatcaaac

SEQ ID NO: 13 is an exemplary amino acid sequence of a precursor of the GP from Bundibugyo EBOV (GENBANK Acc. No. ACI28624.1, which is incorporated by reference herein in its entirety). MVTSGILQLPRERFRKTSFFVWVIILFHKVFPIPLGVVHNNTLQVSDIDKLVCRDKLSSTSQLKSVGLNLE

GNGVATDVPTATKRWGFRAGVPPKVVNYEAGEWAENCYNLDIKKADGSECLPEAPEGVRGFPRCRYVHKVS

GTGPCPEGYAFHKEGAFFLYDRLASTIIYRSTTFSEGVVAFLILPETKKDFFQSPPLHEPANMTTDPSSYY

 ${\tt VGEWAFWENKKNFTKTLSSEELSVIFVPRAQDPGSNQKTKVTPTSFANNQTSKNHEDLVPEDPASVVQVRD}$ 

LORENTVPTPPPDTVPTTLIPDTMEEOTTSHYEPPNISRNHOERNNTAHPETLANNPPDNTTPSTPPODGE

RTSSHTTPSPRPVPTSTIHPTTRETHIPTTMTTSHDTDSNRPNPIDISESTEPGPLTNTTRGAANLLTGSR

LQLFLRATTELRTFSILNRKAIDFLLQRWGGTCHILGPDCCIEPHDWTKNITDKIDQIIHDFIDKPLPDQT

DNDNWWTGWRQWVPAGIGITGVIIAVIALLCICKFLL

SEQ ID NO: 14 is an exemplary amino acid sequence of a precursor of the GP from Sudan EBOV (GENBANK Acc. No. ACR33190.1, which is incorporated by reference herein in its entirety). MEGLSLLQLPRDKFRKSSFFVWVIILFQKAFSMPLGVVTNSTLEVTEIDQLVCKDHLASTDQLKSVGLNLE

GSGVSTDIPSATKRWGFRSGVPPKVFSYEAGEWAENCYNLEIKKPDGSECLPPPPDGVRGFPRCRYVHKAQ

GTGPCPGDYAFHKDGAFFLYDRLASTVIYRGVNFAEGVIAFLILAKPKETFLQSPPIREAVNYTENTSSYY

 ${\tt ATSYLEYEIENFGAQHSTTLFKINNNTFVLLDRPHTPQFLFQLNDTIHLHQQLSNTTGKLIWTLDANINAD$ 

 ${\tt IGEWAFWENKKNLSEQLRGEELSFETLSLNETEDDDATSSRTTKGRISDRATRKYSDLVPKDSPGMVSLHV}$ 

PEGETTLPSQNSTEGRRVDVNTQETITETTATIIGTNGNNMQISTIGTGLSSSQILSSSPTMAPSPETQTS

 ${\tt TTYTPKLPVMTTEESTTPPRNSPGSTTEAPTLTTPENITTAVKTVLPQESTSNGLITSTVTGILGSLGLRK$ 

RSRRQVNTRATGKCNPNLHYWTAQEQHNAAGIAWIPYFGPGAEGIYTEGLMHNQNALVCGLRQLANETTQA

LQLFLRATTELRTYTILNRKAIDFLLRRWGGTCRILGPDCCIEPHDWTKNITDKINQIIHDFIDNPLPNQD

NDDNWWTGWRQWIPAGIGITGIIIAIIALLCVCKLLC

SEQ ID NO: 15 is an exemplary amino acid sequence of a precursor of the GP from Zaire EBOV (GENBANK Acc. No. AIO11753.1, which is incorporated by reference herein in its entirety). MGVTGILQLPRDRFKKTSFFLWVIILFQRTFSIPLGVIHNSTLQVSDVDKLVCRDKLSSTNQLRSVGLNLE

GNGVATDVPSATKRWGFRSGVPPKVVNYEAGEWAENCYNLEIKKPDGSECLPAAPDGIRGFPRCRYVHKVS

GTGPCAGDFAFHKEGAFFLYDRLASTVIYRGTTFAEGVVAFLILPOAKKDFFSSHPLREPVNATEDPSSGY

YSTTIRYOATGFGTNETEYLFEVDNLTYVOLESRFTPOFLLOLNETIYTSGKRSNTTGKLIWKVNPEIDTT

 ${\tt IGEWAFWETKKNLTRKIRSEELSFTAVSNRAKNISGQSPARTSSDPGTNTTTEDHKIMASENSSAMVQVHS}$ 

OGREAAVSHLTTLATISTSPOPPTTKPGPDNSTHNTPVYKLDISEATOAEOHHRRTDNDSTTSDTPPAMTA

 ${\tt AGPPKAENTNTSKGTDLPDPATTTSPQNHSETAGNNNTHHQDTGEESASSGKLGLITNTIAGVAGLITGGR}$ 

#### -continued

RTRREAIVNAQPKCNPNLHYWTTQDEGAAIGLAWIPYFGPAAEGIYTEGLMHNQDGLICGLRQLANETTQA

DNDNWWTGWRQWIPAGIGVTGVIIAVIALFCICKFVF

SEQ ID NO: 16 is an exemplary amino acid sequence of a precursor of the GP from Reston EBOV (GENBANK Acc. No. AAC54891.1, which is incorporated by reference herein in its entirety). MGSGYQLLQLPRERFRKTSFLVWVIILFQRAISMPLGIVTNSTLKATEIDQLVCRDKLSSTSQLKSVGLNL

EGNGIATDVPSATKRWGFRSGVPPKVVSYEAGEWAENCYNLEIKKSDGSECLPLPPDGVRGFPRCRYVHKV

QGTGPCPGDLAFHKNGAFFLYDRLASTVIYRGTTFTEGVVAFLILSEPKKHFWKATPAHEPVNTTDDSTSY

YMTLTLSYEMSNFGGKESNTLFKVDNHTYVQLDRPHTPQFLVQLNETLRRNNRLSNSTGRLTWTLDPKIEP

DVGEWAFWETKKNFSQQLHGENLHFQILSTHTNNSSDQSPAGTVQGKISYHPPTNNSELVPTDSPPVVSVL

TAGR TEEMS TOGLTNGET I TGFTAN PMTTTIAPS PTMTSEVDNNVPSEOPNNTAS I EDSPPSASNET I DHS

EMNPIOGSNNSAOSPOTKTTPAPTASPMTODPOETANSSKLGTSPGSAAEPSOPGFTINTVSKVADSLSPT

RKOKRSVRONTANKCNPDLHYWTAVDEGAAVGLAWI PYFGPAAEGIYI EGVMHNONGLI CGLROLANETTO

ALQLFLRATTELRTYSLLNRKAIDFLLQRWGGTCRILGPSCCIEPHDWTKNITDEINQIKHDFIDNPLPDH

GDDLNLWTGWRQWIPAGIGIIGVIIAIIALLCICKILC

SEQ ID NO: 17 is an exemplary amino acid sequence of a precursor of the GP from Taï Forest EBOV (GENBANK Acc. No. ACI28632.1, which is incorporated by reference herein in its entirety). MGASGILQLPRERFRKTSFFVWVIILFHKVFSIPLGVVHNNTLQVSDIDKFVCRDKLSSTSQLKSVGLNLE

GNGVATDVPTATKRWGFRAGVPPKVVNCEAGEWAENCYNLAIKKVDGSECLPEAPEGVRDFPRCRYVHKVS

GTGPCPGGLAFHKEGAFFLYDRLASTIIYRGTTFAEGVIAFLILPKARKDFFOSPPLHEPANMTTDPSSYY

 ${\tt HTTTINYVVD} {\tt NFGT} {\tt NTTEFLF} {\tt QVD} {\tt LTYVQ} {\tt LEARFTP} {\tt QFLVLL} {\tt NETIYSD} {\tt NRRSNTTGKLIWKINPTVDTS}$ 

 ${\tt MGEWAFWENKKNFTKTLSSEELSFVPVPETQNQVLDTTATVSPPISAHNHAAEDHKELVSEDSTPVVQMQN}$ 

 ${\tt IKGKDTMPTTVTGVPTTTPSPFPINARNTDHTKSFIGLEGPQEDHSTTQPAKTTSQPTNSTESTTLNPTSE}$ 

 ${\tt PSSRGTGPSSPTVPNTTESHAELGKTTPTTLPEQHTAASAIPRAVHPDELSGPGFLTNTIRGVTNLLTGSR}$ 

RKRRDVTPNTQPKCNPNLHYWTALDEGAAIGLAWIPYFGPAAEGIYTEGIMENQNGLICGLRQLANETTQA

 ${\tt LQLFLRATTELRTFSILNRKAIDFLLQRWGGTCHILGPDCCIEPQDWTKNITDKIDQIIHDFVDNNLPNQN}$ 

DGSNWWTGWKQWVPAGIGITGVIIAIIALLCICKFML

SEQ ID NO: 18 is an exemplary amino acid sequence of a precursor of the soluble form of GP from Zaire EBOV (GENBANK Acc. No. AAD14584.1, which is incorporated by reference herein in its entirety).

 ${\tt MGVTGILQLPRDRFKRTSFFLWVIILFQRTFSIPLGVIHNSTLQVSDVDKLVCRDKLSSTNQLRSVGLNLE}$ 

 ${\tt GNGVATDVPSATKRWGFRSGVPPKVVNYEAGEWAENCYNLEIKKPDGSECLPAAPDGIRGFPRCRYVHKVS}$ 

 ${\tt GTGPCAGDFAFHKEGAFFLYDRLASTVIYRGTTFAEGVVAFLILPQAKKDFFSSHPLREPVNATEDPSSGY}$ 

YSTTIRYQATGFGTNETEYLFEVDNLTYVQLESRFTPQFLLQLNETIYTSGKRSNTTGKLIWKVNPEIDTT

IGEWAFWETKKTSLEKFAVKSCLSQLYQTEPKTSVVRVRRELLPTQGPTQQLKTTKSWLQKIPLQWFKCTV

#### KEGKLQCRI

SEQ ID NO: 19 is the amino acid sequence of the  $V_H$  of the EVB165 mAb. DVQLVESGGGVVQPGGSLKLACVVS**gfrfsdyw**MSWVRQAPGKGLEWVAN**ikqdgsgk**YYVDSVKGRFTVS

 ${\tt RDNAKNSLYLHMTSLGAEDTAVYFC} {\tt araaptgsytnilvdnvhfdy} {\tt WGQGILVAVSS}$ 

SEQ ID NO: 20 is the amino acid sequence of the  $\rm V_L$  of the EVB165 mAb. GIQLTQSPGSLSASVGDSVTITCRPNqniatyINWYQQTPGKAPKLLIYaasILQSGVPSRFSGAGSGTHF

### TLIISTLQPEDSATYYC qqsystpwt FGQGTKVEIK

 $\verb+ctct} \textbf{GGATTCAGGTTTAGTGACTACTGG} \texttt{atgagttgggtccgccaggccccagggaaggggctggaatggg}$ 

## -continued

tggccaacATAAAACAAGATGGAAGTGGGAAGtactatgtggactccgtgaagggccgattcaccgtctcc

agagacaacgccaagaactcactgtatctacacatgaccagcctgggagccgaggacacggccgtatactt

### ctgcGCGAGAGCAGCCCCCACCGGCTCCTACACTAATATCCTAGTCGACAACGTCCACTTCGACTACtggg

gccagggaatcctggtcgccgtctcctcag

SEQ ID NO: 22 is an exemplary nucleotide sequence encoding the  $\rm V_L$  of the EVB165 mAb. ggcatccagctgacccagtctcccaggctccctgtctgcatctgtaggagacagtgtcaccatcacttgccg

 $\verb|gccaaatCAGAACATCGCCACCTATataaattggtatcagcagacaccagggaaagcccctaagctcctga||$ 

tctat GCCGCATCC attttgcagagtggggtcccatcaaggttcagtggcgctggatctgggacacatttc

GTGGACAttcggccaagggaccaaagtggaaatcaaac

SEQ ID NO: 23 is the amino acid sequence of the  $V_H$  of the EVB167 mAb. AVQLVQSGAEVKKPGTTVKISCKVS**gytfiqey**IHWVQQAPGKGLVWMGL**gdpennet**LYSEDFQGRVTMT

ADTSSDTAYLELRSLTFADTAVYFC**tsrksw**WGQGTLVTVAS

SEQ ID NO: 24 is the amino acid sequence of the  $\rm V_L$  of the EVB167 mAb. ELVLTQSPGTLSLSPGESATLSCRASqslssdsVSWFQQKPGQAPRLVIHgtsKRATGIPDRFSGGGSGTD

### FTLTIARLEPEDFAVYYCqrsgygmsvtwtFGQGTTVEIK

SEQ ID NO: 25 is an exemplary nucleotide sequence encoding the  ${\rm V}_H$  of the EVB167 mAb. gcggtccagttggtacaatctggggctgaggtgaagaagcctgggaccaccgtcaaaatctcctgcaaagt

 $\verb+tct \texttt{GGATACACCTTCATTCAAGAATAC} a \verb+tacactgggtgcaacaggcccctggaaaagggcttgtgtgga$ 

 ${\tt tgggactt} {\tt GGTGACCCTGAAAATAATGAGACT} {\tt ctatattcagaggatttccaaggcagagtcaccatgacc}$ 

gcggacacatcctcagacacagcctatctggaactgcgcagcctgacatttgcagacacggccgtctattt

ctgtACATCACGAAAGTCCTGGtgggggccagggaaccctggtcaccgtcgcctcag

SEQ ID NO: 26 is an exemplary nucleotide sequence encoding the  $\rm V_L$  of the EVB167 mAb. gaacttgtgttgacgcagtctcccaggcaccctgtctttgtctccaggggaaagcgccaccctctc

 $\tt ctgtagggccagt {\tt CAGAGTCTTAGCAGCGACTCT} gtatcttggttccagcagaaacctggccagg$ 

ctcccaggctcgtcatccatGGTACATCAaagagggccactggcatcccagacaggttcagtggc

ggtgggtctgggacagacttcactctcaccatcgccagactggagcctgaggattttgcagtcta

ttattgt**CAGCGGTCTGGGTATGGTATGTCAGTCACGTGGACG**ttcggccaagggaccacggtgg

### agatcaaac

SEQ ID NO: 27 is an exemplary nucleotide sequence encoding the  $\rm V_L$  of the EVB167 mAb. gaacttgtgttgacgcagtctcccaggcaccctgtctttgtctccaggggaaagcgccaccctctc

 $\tt ctgtagggccagt {\tt CAGAGTCTTAGCAGCGACTCT} gtatcttggttccagcagaaacctggccagg$ 

 $\verb+ctcccaggctcgtcatccatggtAcATCA+ a agagggccactggcatcccagacaggttcagtggc+ agagggccactggcatcccagacaggttcagtggc+ agagggccactggcatcccagacaggttcagtggc+ agagggccactggcatcccagacaggttcagtggc+ agagggccactggcatcccagacaggttcagtggc+ agagggccactggcatcccagacaggttcagtggc+ agagggccactggcatcccagacaggttcagtggc+ agagggccactggcatcccagacaggttcagtggc+ agagggccactggcatcccagacaggttcagtggc+ agagggc+ agagggc+ agagggccactggc+ agagggc+ agaggg+ agagggc+ agagggc+ agagggc+ agaggg+ agagggc+ agagggc+ agaggg+ agagg+ agaggg+ agaggg+ agaggg+ agagg+ agagg+ agagg+ agaggg+ agagg+ agaggg+ agagg+ aga$ 

ggtgggtctgggacagacttcactctcaccatcgccagactggagcctgaggattttgcagtcta

ttattgt CAGCGGTCTGGGTATGGTATGTCAGTCACGTGGACG ttggccaagggaccacggtgg

### agatcaaac

SEQ ID NO: 28 is the amino acid sequence of the  $\rm V_{\it H}$  of the EVB114 version 2 mAb. EVQLVESGGGLIQPGGSLRLSCAAsgfalrsydMHWVRQTIDKRLEWVSAvgpsgdtYYADSVKGRFAVSR

ENAKNSLSLQMNSLTAGDTAIYYC vrsdrgvaglfds WGQGILVTVSS

SEQ ID NO: 29 is the amino acid sequence of the  $\rm V_L$  of the EVB114 version 2 mAb. DIQMTQSPSSLSASVGDRITITCRASqafsnyVAWYQQRPGKVPKLLISaasALHAGVPSRFSGSGSGTHF

#### TLTISSLQPEDVATYYCqnynsapltFGGGTKVEI

SEQ ID NO: 30 is an exemplary nucleotide sequence encoding the  $\mathrm{V}_H$  of the EVB114 version 2 mAb.

gaagtgcagctggtggagtctggaggaggtctgattcagcccgggggttccctgcgtctgagttgtgccgc

atct GGATTTGCTCTGCGAAGCTACGAC atgcactgggtgagacagactatcgataagcgcctggagtggg

### -continued

 ${\tt tgtctgct} {\tt GTCGGCCCCAGTGGAGACACC} {\tt tgtctgcagattcagtgaagggaggttcgcagtctcccgg}$ 

gaaaacgccaaaaattccctgagcctgcagatgaactctctgaccgccggcgacacagctatctactattg

c GTCAGGAGCGATAGAGGGGTCGCAGGACTGTTTGATTCA t ggggtcagggtattctggtcaccgtgtctt

са

SEQ ID NO: 31 is an exemplary nucleotide sequence encoding the  $\mathrm{V}_L$  of the EVB114 version 2 mAb.

gatattcagatgactcagagcccttcctcactgtccgcatccgtggggagaccgtattactattacttgtag

 $agcttct {\tt CAGGCTTTTTCTAACTAC} gtggcttggtatcagcagaggcccggcaaggtccctaaactgctgaaggtcccggcaaggtcccctaaactgctgaaggtccctaaactgctgaaggtcccggcaaggtcccctaaactgctgaaggtcccggcaaggtcccctaaactgctgaaggtcccqgcaaggtcccctaaactgctgaaggtcccqgcaaggtcccctaaactgctgaaggtcccqgcaaggtccccqgcaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccqgcaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaaggtccccqgcaaggtcccctaaactgctgaaggtcccctaaactgctgaaggtcccctaaaggtcccctaaaggtcccctaaaggtcccctaaaggtcccctaaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtccccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtcccctaaggtccccctaaggtcccccctaaggtcccccta$ 

accctgacaatttccagcctgcagcccgaggatgtcgcaacctactattgc CAGAACTACAACAGTGCTCC

**CCTGACA**ttcggtggtggaacaaaggtcgagatc

 ${\tt SEQ}$  ID NOs: 32-61 are amino acid sequences of antibody heavy and light chain CDRs by IMGT positioning.

SEQ ID NO: 62 is the amino acid sequence of a variant  $V_L$  of the EVB166 mAb that includes a K104T substitution.

25

EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPRLLIYGTSSRATGIPDRFSGSASGTD

FTLTISRLEPEDFAVYYCQQYAYSPFTFGPGTTVDIK

SEQ ID NOs: 63-65 are primer and probe sequences. SEQ ID NO: 66 is the amino acid sequence of a modified fragment of EBOV GP.

For SEQ ID NOs: 1-6, 19-20, 23-24, and 28-29 the amino acid sequence of the IMGT CDRs are shown in bold and <sup>30</sup> lower case letters. For SEQ ID NOs: 7-12, 21-22, 25-27, and 30-31, the nucleotide sequences encoding IMGT CDRs are shown in bold and upper case letters.

## DETAILED DESCRIPTION

## I. Summary of Terms

Unless otherwise noted, technical terms are used according to conventional usage. Definitions of common terms in molecular biology may be found in Benjamin Lewin, Genes X, published by Jones & Bartlett Publishers, 2009; and Meyers et al. (eds.), *The Encyclopedia of Cell Biology and Molecular Medicine*, published by Wiley-VCH in 16 vol-45 umes, 2008; and other similar references.

As used herein, the singular forms "a," "an," and "the," refer to both the singular as well as plural, unless the context clearly indicates otherwise. For example, the term "an antigen" includes single or plural antigens and can be 50 considered equivalent to the phrase "at least one antigen." As used herein, the term "comprises" means "includes." It is further to be understood that any and all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approxi-55 mate, and are provided for descriptive purposes, unless otherwise indicated. Although many methods and materials similar or equivalent to those described herein can be used, particular suitable methods and materials are described herein. In case of conflict, the present specification, includ- 60 ing explanations of terms, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting. To facilitate review of the various embodiments, the following explanations of terms are provided: 65

Administration: The introduction of a composition into a subject by a chosen route.

Administration can be local or systemic. For example, if the chosen route is intravenous, the composition is administered by introducing the composition into a vein of the subject. Exemplary routes of administration include, but are not limited to, oral, injection (such as subcutaneous, intramuscular, intradermal, intraperitoneal, and intravenous), sublingual, rectal, transdermal (for example, topical), intranasal, vaginal, and inhalation routes.

Agent: Any substance or any combination of substances that is useful for achieving an end or result; for example, a substance or combination of substances useful for inhibiting EBOV infection in a subject. Agents include proteins, antibodies, nucleic acid molecules, compounds, small molecules, organic compounds, inorganic compounds, or other molecules of interest. An agent can include a therapeutic agent, a diagnostic agent or a pharmaceutical agent. In some embodiments, the agent is a polypeptide agent (such as an EBOV-neutralizing antibody), or an anti-viral agent. Some agents may be useful to achieve more than one result.

Amino acid substitution: The replacement of one amino acid in peptide with a different amino acid.

Antibody: An immunoglobulin, antigen-binding fragment, or derivative thereof, that specifically binds and recognizes an analyte (antigen) such as EBOV GP. The term "antibody" is used herein in the broadest sense and encompasses various antibody structures, including but not limited to monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments, so long as they exhibit the desired antigenbinding activity.

Non-limiting examples of antibodies include, for example, intact immunoglobulins and variants and fragments thereof known in the art that retain binding affinity for the antigen. Examples of antibody fragments include but are not limited to Fv, Fab, Fab', Fab'-SH,  $F(ab')_2$ ; diabodies; linear antibodies; single-chain antibody molecules (e.g. scFv); and multispecific antibodies formed from antibody fragments. Antibody fragments include antigen binding fragments either produced by the modification of whole antibodies or those synthesized de novo using recombinant DNA methodologies (see, e.g., Kontermann and Dubel (Ed), Antibody Engineering, Vols. 1-2,  $2^{nd}$  Ed., Springer Press, 2010).

14

A single-chain antibody (scFv) is a genetically engineered molecule containing the  $V_H$  and  $V_L$  domains of one or more antibody(ies) linked by a suitable polypeptide linker as a genetically fused single chain molecule (see, for example, Bird et al., Science, 242:423-426, 1988; Huston et al., Proc. Natl. Acad. Sci., 85:5879-5883, 1988; Ahmad et al., Clin. Dev. Immunol., 2012, doi:10.1155/2012/980250; Marbry, IDrugs, 13:543-549, 2010). The intramolecular orientation of the  $V_H$ -domain and the  $V_L$ -domain in a scFv, is typically not decisive for scFvs. Thus, scFvs with both possible 10 arrangements (V<sub>H</sub>-domain-linker domain-V<sub>L</sub>-domain; V<sub>L</sub>domain-linker domain- $V_H$ -domain) may be used.

In a dsFv the  $V_H$  and  $V_L$  have been mutated to introduce a disulfide bond to stabilize the association of the chains. Diabodies also are included, which are bivalent, bispecific 15 antibodies in which  $V_H$  and  $V_L$  domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen 20 binding sites (see, for example, Holliger et al., Proc. Natl. Acad. Sci., 90:6444-6448, 1993; Poljak et al., Structure, 2:1121-1123, 1994).

Antibodies also include genetically engineered forms such as chimeric antibodies (such as humanized murine 25 antibodies) and heteroconjugate antibodies (such as bispecific antibodies). See also, Pierce Catalog and Handbook, 1994-1995 (Pierce Chemical Co., Rockford, Ill.); Kuby, J., Immunology, 3rd Ed., W.H. Freeman & Co., New York, 1997.

An "antibody that binds to the same epitope" as a reference antibody refers to an antibody that blocks binding of the reference antibody to its antigen in a competition assay by 50% or more, and conversely, the reference antibody blocks binding of the antibody to its antigen in a competition 35 assay by 50% or more. Antibody competition assays are known, and an exemplary competition assay is provided herein.

An antibody may have one or more binding sites. If there is more than one binding site, the binding sites may be 40 identical to one another or may be different. For instance, a naturally-occurring immunoglobulin has two identical binding sites, a single-chain antibody or Fab fragment has one binding site, while a bispecific or bifunctional antibody has two different binding sites. 45

Typically, a naturally occurring immunoglobulin has heavy (H) chains and light (L) chains interconnected by disulfide bonds. Immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as the myriad immunoglobulin vari- 50 able domain genes. There are two types of light chain, lambda ( $\lambda$ ) and kappa ( $\kappa$ ). There are five main heavy chain classes (or isotypes) which determine the functional activity of an antibody molecule: IgM, IgD, IgG, IgA and IgE.

Each heavy and light chain contains a constant region (or 55 constant domain) and a variable region (or variable domain; see, e.g., Kindt et al. Kuby Immunology, 6<sup>th</sup> ed., W.H. Freeman and Co., page 91 (2007).) In several embodiments, the  $V_H$  and  $V_L$  combine to specifically bind the antigen. In additional embodiments, only the  $V_H$  is required. For 60 example, naturally occurring camelid antibodies consisting of a heavy chain only are functional and stable in the absence of light chain (see, e.g., Hamers-Casterman et al., Nature, 363:446-448, 1993; Sheriff et al., Nat. Struct. Biol., 3:733-736, 1996). Any of the disclosed antibodies can 65 include a heterologous constant domain. For example the antibody can include constant domain that is different from

16

a native constant domain, such as a constant domain including one or more modifications (such as the "LS" mutations) to increase half-life.

References to " $V_H$ " or "VH" refer to the variable region of an antibody heavy chain, including that of an antigen binding fragment, such as Fv, scFv, dsFv or Fab. References to " $V_L$ " or "VL" refer to the variable domain of an antibody light chain, including that of an Fv, scFv, dsFv or Fab.

The  $V_H$  and  $V_L$  contain a "framework" region interrupted by three hypervariable regions, also called "complementarity-determining regions" or "CDRs" (see, e.g., Kabat et al., Sequences of Proteins of Immunological Interest, U.S. Department of Health and Human Services, 1991). The sequences of the framework regions of different light or heavy chains are relatively conserved within a species. The framework region of an antibody, that is the combined framework regions of the constituent light and heavy chains, serves to position and align the CDRs in three-dimensional space.

The CDRs are primarily responsible for binding to an epitope of an antigen. The amino acid sequence boundaries of a given CDR can be readily determined using any of a number of well-known schemes, including those described by Kabat et al. ("Sequences of Proteins of Immunological Interest," 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md., 1991; "Kabat" numbering scheme), Al-Lazikani et al., (JMB 273,927-948, 1997; "Chothia" numbering scheme), and Lefranc et al. ("IMGT unique numbering for immunoglobulin and T cell receptor variable domains and Ig superfamily V-like domains," Dev. Comp. Immunol., 27:55-77, 2003; "IMGT" numbering scheme). The CDRs of each chain are typically referred to as CDR1, CDR2, and CDR3 (from the N-terminus to C-terminus), and are also typically identified by the chain in which the particular CDR is located. Thus, a  $V_H$  CDR3 is the CDR3 from the  $V_H$  of the antibody in which it is found, whereas a  $V_L$  CDR1 is the CDR1 from the  $V_L$  of the antibody in which it is found. Light chain CDRs are sometimes referred to as LCDR1, LCDR2, and LCDR3. Heavy chain CDRs are sometimes referred to as HCDR1, HCDR2, and HCDR3.

A "monoclonal antibody" is an antibody obtained from a population of substantially homogeneous antibodies, that is, the individual antibodies comprising the population are identical and/or bind the same epitope, except for possible variant antibodies, for example, containing naturally occurring mutations or arising during production of a monoclonal antibody preparation, such variants generally being present in minor amounts. In contrast to polyclonal antibody preparations, which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody of a monoclonal antibody preparation is directed against a single determinant on an antigen. Thus, the modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies may be made by a variety of techniques, including but not limited to the hybridoma method, recombinant DNA methods, phagedisplay methods, and methods utilizing transgenic animals containing all or part of the human immunoglobulin loci, such methods and other exemplary methods for making monoclonal antibodies being described herein. In some examples monoclonal antibodies are isolated from a subject. Monoclonal antibodies can have conservative amino acid substitutions which have substantially no effect on antigen

binding or other immunoglobulin functions. (See, for example, Harlow & Lane, *Antibodies, A Laboratory Manual*,  $2^{nd}$  ed. Cold Spring Harbor Publications, New York (2013).)

A "humanized" antibody or antigen binding fragment 5 includes a human framework region and one or more CDRs from a non-human (such as a mouse, rat, or synthetic) antibody or antigen binding fragment. The non-human antibody or antigen binding fragment providing the CDRs is termed a "donor," and the human antibody or antigen 10 binding fragment providing the framework is termed an "acceptor." In one embodiment, all the CDRs are from the donor immunoglobulin in a humanized immunoglobulin. Constant regions need not be present, but if they are, they can be substantially identical to human immunoglobulin 15 constant regions, such as at least about 85-90%, such as about 95% or more identical. Hence, all parts of a humanized antibody or antigen binding fragment, except possibly the CDRs, are substantially identical to corresponding parts of natural human antibody sequences.

A "chimeric antibody" is an antibody which includes sequences derived from two different antibodies, which typically are of different species. In some examples, a chimeric antibody includes one or more CDRs and/or framework regions from one human antibody and CDRs and/or 25 framework regions from another human antibody.

A "fully human antibody" or "human antibody" is an antibody which includes sequences from (or derived from) the human genome, and does not include sequence from another species. In some embodiments, a human antibody 30 includes CDRs, framework regions, and (if present) an Fc region from (or derived from) the human genome. Human antibodies can be identified and isolated using technologies for creating antibodies based on sequences derived from the human genome, for example by phage display or using 35 transgenic animals (see, e.g., Barbas et al. *Phage display: A Laboratory Manuel.* 1<sup>st</sup> Ed. New York: Cold Spring Harbor Laboratory Press, 2004. Print.; Lonberg, Nat. Biotech., 23: 1117-1125, 2005; Lonenberg, Curr. Opin. Immunol., 20:450-459, 2008)

Antibody or antigen binding fragment that neutralizes EBOV: An antibody or antigen binding fragment that specifically binds to EBOV GP (such as ZEBOV GP) in such a way as to inhibit a biological function associated with EBOV GP (such as binding to its target receptor). In several 45 embodiments, an antibody or antigen binding fragment that neutralizes EBOV reduces the infectious titer of EBOV. In some embodiments, an antibody or antigen binding fragment that specifically binds to EBOV GP can neutralize two or more (such as 3, 4, 5, 6, 7, 8, 9, 10, or more) strains of 50 EBOV.

Biological sample: A sample obtained from a subject. Biological samples include all clinical samples useful for detection of disease or infection (for example, EVD or EBOV infection) in subjects, including, but not limited to, 55 cells, tissues, and bodily fluids, such as blood, derivatives and fractions of blood (such as serum), cerebrospinal fluid; as well as biopsied or surgically removed tissue, for example tissues that are unfixed, frozen, or fixed in formalin or paraffin. In a particular example, a biological sample is 60 obtained from a subject having or suspected of having an Ebola infection.

Bispecific antibody: A recombinant molecule composed of two different antigen binding domains that consequently binds to two different antigenic epitopes. Bispecific antibodies include chemically or genetically linked molecules of two antigen-binding domains. The antigen binding domains

can be linked using a linker. The antigen binding domains can be monoclonal antibodies, antigen-binding fragments (e.g., Fab, scFv), or combinations thereof. A bispecific antibody can include one or more constant domains, but does not necessarily include a constant domain.

Conditions sufficient to form an immune complex: Conditions which allow an antibody or antigen binding fragment to bind to its cognate epitope to a detectably greater degree than, and/or to the substantial exclusion of, binding to substantially all other epitopes. Conditions sufficient to form an immune complex are dependent upon the format of the binding reaction and typically are those utilized in immunoassay protocols or those conditions encountered in vivo. See Harlow & Lane, Antibodies, A Laboratory Manual, 2nd ed. Cold Spring Harbor Publications, New York (2013) for a description of immunoassay formats and conditions. The conditions employed in the methods are "physiological conditions" which include reference to conditions (e.g., 20 temperature, osmolarity, pH) that are typical inside a living mammal or a mammalian cell. While it is recognized that some organs are subject to extreme conditions, the intraorganismal and intracellular environment normally lies around pH 7 (e.g., from pH 6.0 to pH 8.0, more typically pH 6.5 to 7.5), contains water as the predominant solvent, and exists at a temperature above 0° C. and below 50° C. Osmolarity is within the range that is supportive of cell viability and proliferation.

The formation of an immune complex can be detected through conventional methods, for instance immunohistochemistry, immunoprecipitation, flow cytometry, immunofluorescence microscopy, ELISA, immunoblotting (for example, Western blot), magnetic resonance imaging, CT scans, X-ray and affinity chromatography. Immunological binding properties of selected antibodies may be quantified using methods well known in the art.

Conjugate: A complex of two molecules linked together, for example, linked together by a covalent bond. In one embodiment, an antibody is linked to an effector molecule; for example, an antibody that specifically binds to EBOV GP covalently linked to an effector molecule. The linkage can be by chemical or recombinant means. In one embodiment, the linkage is chemical, wherein a reaction between the antibody moiety and the effector molecule has produced a covalent bond formed between the two molecules to form one molecule. A peptide linker (short peptide sequence) can optionally be included between the antibody and the effector molecule. Because conjugates can be prepared from two molecules with separate functionalities, such as an antibody and an effector molecule, they are also sometimes referred to as "chimeric molecules."

Conservative variants: "Conservative" amino acid substitutions are those substitutions that do not substantially affect or decrease a function of a protein, such as the ability of the protein to interact with a target protein. For example, an EBOV-specific antibody can include up to 1, 2, 3, 4, 5, 6, 7, 8, 9, or up to 10 conservative substitutions compared to a reference antibody sequence and retain specific binding activity for EBOV antigen, and/or EBOV neutralization activity. The term conservative variation also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid.

Furthermore, individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids (for instance less than 5%, in some embodiments less than 1%) in an encoded sequence are conservative variations where the alterations result in the substitution of an amino acid with a chemically similar amino acid.

Conservative amino acid substitution tables providing functionally similar amino acids are known. The following 5 six groups are examples of amino acids that are considered to be conservative substitutions for one another:

1) Alanine (A), Serine (S), Threonine (T);

2) Aspartic acid (D), Glutamic acid (E);

3) Asparagine (N), Glutamine (Q);

4) Arginine (R), Lysine (K);

5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and

6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

Non-conservative substitutions are those that reduce an 15 activity or function of the EBOV-specific antibody, such as the ability to specifically bind to EBOV GP. For instance, if an amino acid residue is essential for a function of the protein, even an otherwise conservative substitution may disrupt that activity. Thus, a conservative substitution does 20 not alter the basic function of a protein of interest.

Contacting: Placement in direct physical association; includes both in solid and liquid form, which can take place either in vivo or in vitro. Contacting includes contact between one molecule and another molecule, for example 25 the amino acid on the surface of one polypeptide, such as an antigen, that contacts another polypeptide, such as an antibody. Contacting can also include contacting a cell for example by placing an antibody in direct physical association with a cell.

Control: A reference standard. In some embodiments, the control is a negative control sample obtained from a healthy patient. In other embodiments, the control is a positive control sample obtained from a patient diagnosed with EBOV infection. In still other embodiments, the control is a 35 historical control or standard reference value or range of values (such as a previously tested control sample, such as a group of EBOV patients with known prognosis or outcome, or group of samples that represent baseline or normal values). 40

A difference between a test sample and a control can be an increase or conversely a decrease. The difference can be a qualitative difference or a quantitative difference, for example a statistically significant difference. In some examples, a difference is an increase or decrease, relative to 45 a control, of at least about 5%, such as at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 100%, at least about 150%, at least about 200%, at least about 50 250%, at least about 300%, at least about 350%, at least about 400%, at least about 500%, or greater than 500%.

Degenerate variant: In the context of the present disclosure, a "degenerate variant" refers to a polynucleotide encoding a protein (for example, an antibody that specifi- 55 cally binds EBOV GP) that includes a sequence that is degenerate as a result of the genetic code. There are twenty natural amino acids, most of which are specified by more than one codon. Therefore, all degenerate nucleotide sequences are included as long as the amino acid sequence 60 of the antibody that binds EBOV GP encoded by the nucleotide sequence is unchanged.

Detectable marker: A detectable molecule (also known as a label) that is conjugated directly or indirectly to a second molecule, such as an antibody, to facilitate detection of the 65 second molecule. For example, the detectable marker can be capable of detection by ELISA, spectrophotometry, flow

20

cytometry, microscopy or diagnostic imaging techniques (such as CT scans, MRIs, ultrasound, fiberoptic examination, and laparoscopic examination). Specific, non-limiting examples of detectable markers include fluorophores, chemiluminescent agents, enzymatic linkages, radioactive isotopes and heavy metals or compounds (for example super paramagnetic iron oxide nanocrystals for detection by MRI). In one example, a "labeled antibody" refers to incorporation of another molecule in the antibody. For example, the label 10 is a detectable marker, such as the incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (for example, streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). Various methods of labeling polypeptides and glycoproteins are known in the art and may be used. Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (such as <sup>35</sup>S or <sup>131</sup>I), fluorescent labels (such as fluorescein isothiocvanate (FITC), rhodamine, lanthanide phosphors), enzymatic labels (such as horseradish peroxidase, betagalactosidase, luciferase, alkaline phosphatase), chemiluminescent markers, biotinyl groups, predetermined polypeptide epitopes recognized by a secondary reporter (such as a leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags), or magnetic agents, such as gadolinium chelates. In some embodiments, labels are attached by spacer arms of various lengths to reduce potential steric hindrance. Methods for using detectable markers and guidance in the choice of detectable markers appropriate for various purposes are discussed for example in Sambrook et al. (Molecular Cloning: A Laboratory Manual, 4th ed, Cold Spring Harbor, N.Y., 2012) and Ausubel et al. (In Current Protocols in Molecular Biology, John Wiley & Sons, New York, through supplement 104, 2013).

Detecting: To identify the existence, presence, or fact of something. General methods of detecting are known and may be supplemented with the protocols and reagents disclosed herein. For example, included herein are methods of detecting a cell that expresses EBOV GP in a subject.

Ebola Virus (EBOV): An enveloped, non-segmented, negative, single-stranded RNA virus that causes Ebola virus disease (EVD), formerly known as Ebola hemorrhagic fever (EHF), in humans. EBOV spreads through human-to-human transmission, with infection resulting from direct contact with blood, secretions, organs or other bodily fluids of infected people, and indirect contact with environments contaminated by such fluids (see, e.g., Baize et al., N Engl J Med., 371, 1418-1425, 2014, which is incorporated by reference herein).

The symptoms of EBOV infection and disease are wellknown. Briefly, in humans, EBOV has an initial incubation period of 2 to 21 days (7 days on average, depending on the strain) followed by a rapid onset of non-specific symptoms such as fever, extreme fatigue, gastrointestinal complaints, abdominal pain, anorexia, headache, myalgias and/or arthralgias. These initial symptoms last for about 2 to 7 days after which more severe symptoms related to hemorrhagic fever occur, including hemorrhagic rash, epistaxis, mucosal bleeding, hematuria, hemoptysis, hematemesis, melena, conjunctival hemorrhage, tachypnea, confusion, somnolence, and hearing loss. In general, the symptoms last for about 7 to 14 days after which recovery may occur. Death can occur 6 to 16 days after the onset of symptoms (Geisbert and Jahrling, Nat Med., 10, S110-21. 2004; Hensley et al., Curr Mol Med, 5, 761-72, 2005). People are infectious as

long as their blood and secretions contain the virus; the virus was isolated from semen 61 days after onset of illness in a man who was infected in a laboratory (Baize et al., N Engl J Med., 371, 1418-1425, 2014).

Immunoglobulin M (IgM) antibodies to the virus appear 5 2 to 9 days after infection whereas immunoglobulin G (IgG) antibodies appear approximately 17 to 25 days after infection, which coincides with the recovery phase. In survivors of EVD, both humoral and cellular immunity are detected, however, their relative contribution to protection is unknown 10 (Sullivan, Yang, and Nabel, J Virol, 77, 9733-7, 2003).

Five distinct EBOV species are known, including Bundibugyo (BDBV), Reston (RESTV), Sudan (SUDV), Tai Forest (TAFV), and Zaire (ZEBOV) (Kuhn, J. H., et al., Arch Virol, 2013. 158(1): p. 301-11). BDBV, EBOV, and 15 SUDV have been associated with large outbreaks of EVD in Africa and reported case fatality rates of up to 90%. Exemplary amino acid sequences of EBOV GP from the BDBV, RESTV, SUDV, TAFV, and ZEBOV strains are set forth as SEO ID NOs: 13-17.

The EBOV genome includes about 19K nucleotides, which encode seven structural proteins including NP (a nucleoprotein), VP35 (a polymerase cofactor), VP30 (a transcription activator), VP24, L (a RNA polymerase), and GP (a glycoprotein).

EBOV glycoprotein (GP): The virion-associated transmembrane glycoprotein of EBOV is initially synthesized as a precursor protein of about 675 amino acids in size, designated GP<sub>0</sub>. Individual GP<sub>0</sub> polypeptides form a homotrimer and undergo glycosylation within the Golgi 30 apparatus as well as processing to remove the signal peptide, and cleavage by a cellular protease between approximately positions 500/501 to generate separate  $\mathrm{GP}_1$  and  $\mathrm{GP}_2$  polypeptide chains, which remain associated as GP1/GP2 protomers within the homotrimer. The extracellular  $GP_1$  poly- 35 peptide (approx. 140 kDa) is derived from the aminoterminal portion of the GP<sub>0</sub> precursor, and the GP<sub>2</sub> polypeptide (approx. 26 kDa), which includes extracellular, transmembrane, and cytosolic domains, is derived from the carboxyl-terminal portion of the  $\text{GP}_0$  precursor.  $\text{GP}_1$  is 40 responsible for attachment to new host cells while  $GP_2$ mediates fusion with those cells.

A splice variant of the gene encoding EBOV GP encodes a soluble glycoprotein (sGP) that is secreted from the viral host cell. (Volchkov et al., Virology, 245, 110-119, 1998). 45 sGP and GP1 are identical in their first 295 N-terminal amino acids, whereas the remaining 69 C-terminal amino acids of sGP and 206 amino acids of GP1 are encoded by different reading frames. It has been suggested that secreted sGP may effectively bind antibodies that might otherwise be protec- 50 tive (see, e.g., Sanchez el al., Proc. Natl. Acad. Sci. U.S.A., 93, 3602-3607, 1996; and Volchkov et al., Virology, 245, 110-119, 1998, each of which is incorporated by reference herein in its entirety).

Comparisons of the predicted amino acid sequences for 55 the GPs of the different EBOV strains show conservation of amino acids in the amino-terminal and carboxy-terminal regions with a highly variable region in the middle of the protein (Feldmann el al., Virus Res. 24: 1-19, 1992). The GP of Ebola viruses are highly glycosylaled and contain both 60 N-linked and O-linked carbohydrates that contribute up to 50% of the molecular weight of the protein. Most of the glycosylation sites are found in the central variable region of GP.

The numbering used in the disclosed EBOV GPs and 65 fragments thereof is relative to the EBOV GP protein from the Zaire strain set forth as SEQ ID NO: 15, unless context

indicates otherwise. Effector molecule: A molecule intended to have or produce a desired effect; for example, a desired effect on a cell to which the effector molecule is targeted. Effector molecules can include, for example, polypeptides and small molecules. In one non-limiting example, the effector molecule is a toxin. Some effector molecules may have or produce more than one desired effect.

Epitope: An antigenic determinant. These are particular chemical groups or peptide sequences on a molecule that are antigenic, i.e. that elicit a specific immune response. An antibody specifically binds a particular antigenic epitope on a polypeptide. In some examples a disclosed antibody specifically binds to an epitope on EBOV GP.

Expression: Transcription or translation of a nucleic acid sequence. For example, an encoding nucleic acid sequence (such as a gene) can be expressed when its DNA is transcribed into an RNA or RNA fragment, which in some examples is processed to become mRNA. An encoding nucleic acid sequence (such as a gene) may also be 20 expressed when its mRNA is translated into an amino acid sequence, such as a protein or a protein fragment. In a particular example, a heterologous gene is expressed when it is transcribed into an RNA. In another example, a heterologous gene is expressed when its RNA is translated into an amino acid sequence. Regulation of expression can include controls on transcription, translation, RNA transport and processing, degradation of intermediary molecules such as mRNA, or through activation, inactivation, compartmentalization or degradation of specific protein molecules after they are produced.

Expression Control Sequences: Nucleic acid sequences that regulate the expression of a heterologous nucleic acid sequence to which it is operatively linked. Expression control sequences are operatively linked to a nucleic acid sequence when the expression control sequences control and regulate the transcription and, as appropriate, translation of the nucleic acid sequence. Thus expression control sequences can include appropriate promoters, enhancers, transcription terminators, a start codon (ATG) in front of a protein-encoding gene, splicing signal for introns, maintenance of the correct reading frame of that gene to permit proper translation of mRNA, and stop codons. The term 'control sequences" is intended to include, at a minimum, components whose presence can influence expression, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences. Expression control sequences can include a promoter.

A promoter is a minimal sequence sufficient to direct transcription. Also included are those promoter elements which are sufficient to render promoter-dependent gene expression controllable for cell-type specific, tissue-specific, or inducible by external signals or agents; such elements may be located in the 5' or 3' regions of the gene. Both constitutive and inducible promoters are included (see for example, Bitter et al., Methods in Enzymology 153:516-544, 1987). For example, when cloning in bacterial systems, inducible promoters such as pL of bacteriophage lambda, plac, ptrp, ptac (ptrp-lac hybrid promoter) and the like may be used. In one embodiment, when cloning in mammalian cell systems, promoters derived from the genome of mammalian cells (such as metallothionein promoter) or from mammalian viruses (such as the retrovirus long terminal repeat; the adenovirus late promoter; the vaccinia virus 7.5K promoter) can be used. Promoters produced by recombinant DNA or synthetic techniques may also be used to provide for transcription of the nucleic acid sequences.

A polynucleotide can be inserted into an expression vector that contains a promoter sequence which facilitates the efficient transcription of the inserted genetic sequence of the host. The expression vector typically contains an origin of replication, a promoter, as well as specific nucleic acid <sup>5</sup> sequences that allow phenotypic selection of the transformed cells.

Expression vector: A vector comprising a recombinant polynucleotide comprising expression control sequences operatively linked to a nucleotide sequence to be expressed.<sup>10</sup> An expression vector comprises sufficient cis-acting elements for expression; other elements for expression can be supplied by the host cell or in an in vitro expression system. Expression vectors include all those known in the art, such as cosmids, plasmids (e.g., naked or contained in liposomes) and viruses (e.g., lentiviruses, retroviruses, adenoviruses, and adeno-associated viruses) that incorporate the recombinant polynucleotide.

Fc polypeptide: The polypeptide including the constant 20 region of an antibody excluding the first constant region immunoglobulin domain. Fc region generally refers to the last two constant region immunoglobulin domains of IgA, IgD, and IgG, and the last three constant region immunoglobulin domains of IgE and IgM. An Fc region may also 25 include part or all of the flexible hinge N-terminal to these domains. For IgA and IgM, an Fc region may or may not include the tailpiece, and may or may not be bound by the J chain. For IgG, the Fc region includes immunoglobulin domains Cgamma2 and Cgamma3 (Cy2 and Cy3) and the 30 lower part of the hinge between Cgamma1 (C $\gamma$ 1) and C $\gamma$ 2. Although the boundaries of the Fc region may vary, the human IgG heavy chain Fc region is usually defined to include residues C226 or P230 to its carboxyl-terminus, wherein the numbering is according to the EU index as in 35 Kabat. For IgA, the Fc region includes immunoglobulin domains Calpha2 and Calpha3 (C $\alpha$ 2 and C $\alpha$ 3) and the lower part of the hinge between Calpha1 (C $\alpha$ 1) and C $\alpha$ 2.

Heterologous: Originating from a different genetic source. A nucleic acid molecule that is heterologous to a cell 40 otide or ribonucleotide polymer or combination thereof originated from a genetic source other than the cell in which it is expressed. Nucleic acid (molecule or sequence): A deoxyribonucle-otide polymer or combination thereof including without limitation, cDNA, mRNA, genomic DNA, and synthetic (such as chemically synthesized) DNA

IgA: A polypeptide belonging to the class of antibodies that are substantially encoded by a recognized immunoglobulin alpha gene. In humans, this class or isotype com-5 prises IgA<sub>1</sub> and IgA<sub>2</sub>. IgA antibodies can exist as monomers, polymers (referred to as pIgA) of predominantly dimeric form, and secretory IgA. The constant chain of wild-type IgA contains an 18-amino-acid extension at its C-terminus called the tail piece (tp). Polymeric IgA is secreted by 50 plasma cells with a 15-kDa peptide called the J chain linking two monomers of IgA through the conserved cysteine residue in the tail piece. IgG: A polypeptide belonging to the class or isotype of antibodies that are substantially encoded by a recognized immunoglobulin gamma gene. In humans, 55 this class comprises IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>3</sub>, and IgG<sub>4</sub>. In mice, this class comprises IgG<sub>1</sub>, IgG<sub>2</sub>a, IgG<sub>2</sub>b, IgG<sub>3</sub>.

Immune complex: The binding of antibody or antigen binding fragment (such as a scFv) to a soluble antigen forms an immune complex. The formation of an immune complex <sup>60</sup> can be detected through conventional methods, for instance immunohistochemistry, immunoprecipitation, flow cytometry, immunofluorescence microscopy, ELISA, immunoblotting (for example, Western blot), magnetic resonance imaging, CT scans, X-ray and affinity chromatography. <sup>65</sup> Immunological binding properties of selected antibodies may be quantified using methods well known in the art. 24

Isolated: A biological component (such as a nucleic acid, peptide, protein or protein complex, for example an antibody) that has been substantially separated, produced apart from, or purified away from other biological components in the cell of the organism in which the component naturally occurs, that is, other chromosomal and extra-chromosomal DNA and RNA, and proteins. Thus, isolated nucleic acids, peptides and proteins include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids, peptides and proteins prepared by recombinant expression in a host cell, as well as, chemically synthesized nucleic acids. A isolated nucleic acid, peptide or protein, for example an antibody, can be at least 50%, at least 96%, at least 97%, at least 98%, or at least 99% pure.

Linker: A bi-functional molecule that can be used to link two molecules into one contiguous molecule, for example, to link an effector molecule to an antibody. In some embodiments, the provided conjugates include a linker between the effector molecule or detectable marker and an antibody. In some cases, a linker is a peptide within an antigen binding fragment (such as an Fv fragment) which serves to indirectly bond the variable heavy chain to the variable light chain. Non-limiting examples of peptide linkers include glycine, serine, and glycine-serine linkers.

The terms "conjugating," "joining," "bonding," or "linking" can refer to making two molecules into one contiguous molecule; for example, linking two polypeptides into one contiguous polypeptide, or covalently attaching an effector molecule or detectable marker radionuclide or other molecule to a polypeptide, such as an scFv. In the specific context, the terms include reference to joining a ligand, such as an antibody moiety, to an effector molecule. The linkage can be either by chemical or recombinant means. "Chemical means" refers to a reaction between the antibody moiety and the effector molecule such that there is a covalent bond formed between the two molecules to form one molecule.

Nucleic acid (molecule or sequence): A deoxyribonucleincluding without limitation, cDNA, mRNA, genomic DNA, and synthetic (such as chemically synthesized) DNA or RNA. The nucleic acid can be double stranded (ds) or single stranded (ss). Where single stranded, the nucleic acid can be the sense strand or the antisense strand. Nucleic acids can include natural nucleotides (such as A, T/U, C, and G), and can include analogs of natural nucleotides, such as labeled nucleotides. "Encoding" refers to the inherent property of specific sequences of nucleotides in a polynucleotide, such as a gene, a cDNA, or an mRNA, to serve as templates for synthesis of other polymers and macromolecules in biological processes having either a defined sequence of nucleotides (i.e., rRNA, tRNA and mRNA) or a defined sequence of amino acids and the biological properties resulting therefrom. Thus, a gene encodes a protein if transcription and translation of mRNA produced by that gene produces the protein in a cell or other biological system. Both the coding strand, the nucleotide sequence of which is identical to the mRNA sequence and is usually provided in sequence listings, and non-coding strand, used as the template for transcription, of a gene or cDNA can be referred to as encoding the protein or other product of that gene or cDNA. Unless otherwise specified, a "nucleotide sequence encoding an amino acid sequence" includes all nucleotide sequences that are degenerate versions of each other and that encode the same amino acid sequence. Nucleotide sequences that encode proteins and RNA may include introns. "cDNA"

65

refers to a DNA that is complementary or identical to an mRNA, in either single stranded or double stranded form.

Operably linked: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter, such as the CMV promoter, is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein-coding regions, in the same reading frame.

Pharmaceutically acceptable carriers: The pharmaceutically acceptable carriers of use are conventional. Remington's Pharmaceutical Science, 22th ed., Pharmaceutical 15 Press, London, UK (2012), describes compositions and formulations suitable for pharmaceutical delivery of the disclosed agents.

In general, the nature of the carrier will depend on the particular mode of administration being employed. For 20 instance, parenteral formulations usually include injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced salt solutions, aqueous dextrose, glycerol or the like as a vehicle. For solid compositions (e.g., powder, pill, tablet, 25 or capsule forms), conventional non-toxic solid carriers can include, for example, pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. In addition to biologically neutral carriers, pharmaceutical compositions to be administered can contain minor amounts of non-toxic aux- 30 iliary substances, such as wetting or emulsifying agents, added preservatives (such as on-natural preservatives), and pH buffering agents and the like, for example sodium acetate or sorbitan monolaurate. In particular examples, the pharmaceutically acceptable carrier is sterile and suitable for 35 parenteral administration to a subject for example, by injection. In some embodiments, the active agent and pharmaceutically acceptable carrier are provided in a unit dosage form such as a pill or in a selected quantity in a vial. Unit dosage forms can include one dosage or multiple dosages 40 (for example, in a vial from which metered dosages of the agents can selectively be dispensed).

Polypeptide: A polymer in which the monomers are amino acid residues that are joined together through amide bonds. When the amino acids are alpha-amino acids, either the 45 L-optical isomer or the D-optical isomer can be used, the L-isomers being preferred. The terms "polypeptide" or "protein" as used herein are intended to encompass any amino acid sequence and include modified sequences such as glycoproteins. A polypeptide includes both naturally occur- 50 ring proteins, as well as those that are recombinantly or synthetically produced. A polypeptide has an amino terminal (N-terminal) end and a carboxy-terminal end. In some embodiments, the polypeptide is a disclosed antibody or a fragment thereof.

Purified: The term purified does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified peptide preparation is one in which the peptide or protein (such as an antibody) is more enriched than the peptide or protein is in its natural environment 60 within a cell. In one embodiment, a preparation is purified such that the protein or peptide represents at least 50% of the total peptide or protein content of the preparation, such as at least 80%, at least 90%, at least 95% or greater of the total peptide or protein content.

Recombinant: A recombinant nucleic acid is one that has a sequence that is not naturally occurring or has a sequence 26

that is made by an artificial combination of two otherwise separated segments of sequence. This artificial combination can be accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, for example, by genetic engineering techniques. A recombinant protein is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. In several embodiments, a recombinant protein is encoded by a heterologous (for example, recombinant) nucleic acid that has been introduced into a host cell, such as a bacterial or eukaryotic cell. The nucleic acid can be introduced, for example, on an expression vector having signals capable of expressing the protein encoded by the introduced nucleic acid or the nucleic acid can be integrated into the host cell chromosome.

Sequence identity: The similarity between amino acid or nucleic acid sequences is expressed in terms of the similarity between the sequences, otherwise referred to as sequence identity. Sequence identity is frequently measured in terms of percentage identity (or similarity or homology); the higher the percentage, the more similar the two sequences are. Homologs or variants of a polypeptide or nucleic acid molecule will possess a relatively high degree of sequence identity when aligned using standard methods.

Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith and Waterman, Adv. Appl. Math. 2:482, 1981; Needleman and Wunsch, J. Mol. Biol. 48:443, 1970; Pearson and Lipman, Proc. Natl. Acad. Sci. U.S.A. 85:2444, 1988; Higgins and Sharp, Gene 73:237, 1988; Higgins and Sharp, CABIOS 5:151, 1989; Corpet et al., Nucleic Acids Research 16:10881, 1988; and Pearson and Lipman, Proc. Natl. Acad. Sci. U.S.A. 85:2444, 1988. Altschul et al., Nature Genet. 6:119, 1994, presents a detailed consideration of sequence alignment methods and homology calculations.

The NCBI Basic Local Alignment Search Tool (BLAST) (Altschul et al., J. Mol. Biol. 215:403, 1990) is available from several sources, including the National Center for Biotechnology Information (NCBI, Bethesda, Md.) and on the internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx. A description of how to determine sequence identity using this program is available on the NCBI website on the internet.

Homologs and variants of a polypeptide (such as an insect ferritin heavy or light chain) are typically characterized by possession of at least about 75%, for example at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity counted over the full length alignment with the amino acid sequence of interest. Proteins with even greater similarity to the reference sequences will show increasing percentage identities when assessed by this method, such as at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or at least 99% sequence identity. When less than the entire sequence is being compared for sequence identity, homologs and variants will typically possess at least 80% sequence identity over short windows of 10-20 amino acids, and may possess sequence identities of at least 85% or at least 90% or 95% depending on their similarity to the reference sequence. Methods for determining sequence identity over such short windows are available at the NCBI website on the internet. One of skill in the art will appreciate that these sequence identity ranges are provided for guidance only; it is entirely possible that strongly significant homologs could be obtained that fall outside of the ranges provided.

As used herein, reference to "at least 90% identity" refers to "at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or even 100% identity" to a specified reference sequence.

Specifically bind: When referring to an antibody or antigen binding fragment, refers to a binding reaction which determines the presence of a target protein, peptide, or polysaccharide in the presence of a heterogeneous population of proteins and other biologics. Thus, under designated 10 conditions, an antibody binds preferentially to a particular target protein, peptide or polysaccharide (such as an antigen present on the surface of a pathogen, for example EBOV GP) and does not bind in a significant amount to other proteins or polysaccharides present in the sample or subject. 15 With reference to an antibody-antigen complex, specific binding of the antigen and antibody has a K<sub>d</sub> of less than about  $10^{-7}$  Molar, such as less than about  $10^{-8}$  Molar,  $10^{-9}$ , or even less than about  $10^{-10}$  Molar.

 $K_d$  refers to the dissociation constant for a given interaction, such as a polypeptide ligand interaction or an antibody antigen interaction. For example, for the bimolecular interaction of an antibody or antigen binding fragment (such as EVB114 or an antigen binding fragment thereof) and an antigen (such as EBOV GP) it is the concentration of the 25 individual components of the bimolecular interaction divided by the concentration of the complex.

The antibodies disclosed herein specifically bind to a defined target (or multiple targets, in the case of a bispecific antibody). Thus, an antibody that specifically binds to an 30 epitope on EBOV GP is an antibody that binds substantially to EBOV GP, including cells or tissue expressing EBOV GP, substrate to which the EBOV GP is attached, or EBOV GP in a biological specimen. It is, of course, recognized that a certain degree of non-specific interaction may occur 35 between an antibody or conjugate including an antibody (such as an antibody that specifically binds EBOV GP or conjugate including such antibody) and a non-target (such as a cell that does not express EBOV GP). Typically, specific binding results in a much stronger association between the 40 antibody and protein or cells bearing the antigen than between the antibody and protein or cells lacking the antigen. Specific binding typically results in greater than 2-fold, such as greater than 5-fold, greater than 10-fold, or greater than 100-fold increase in amount of bound antibody (per 45 unit time) to a protein including the epitope or cell or tissue expressing the target epitope as compared to a protein or cell or tissue lacking this epitope. Specific binding to a protein under such conditions requires an antibody that is selected for its specificity for a particular protein. A variety of 50 immunoassay formats are appropriate for selecting antibodies or other ligands specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See Harlow & 55 Lane, Antibodies, A Laboratory Manual, 2<sup>nd</sup> ed., Cold Spring Harbor Publications, New York (2013), for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity.

Subject: Living multi-cellular vertebrate organisms, a 60 category that includes human and non-human mammals. In an example, a subject is a human. In an additional example, a subject is selected that is in need of inhibiting of an EBOV infection. For example, the subject is either uninfected and at risk of EBOV infection or is infected in need of treatment. 65

Therapeutically effective amount: The amount of agent, such as a disclosed EBOV GP specific antibody or antigen

binding fragment that is sufficient to prevent, treat (including prophylaxis), reduce and/or ameliorate the symptoms and/or underlying causes of a disorder or disease, for example to prevent, inhibit, and/or treat EBOV infection. In some embodiments, a therapeutically effective amount is sufficient to reduce or eliminate a symptom of a disease, such as EVD. For instance, this can be the amount necessary to inhibit or prevent EBOV replication or to measurably alter outward symptoms of the EBOV infection. Ideally, a therapeutically effective amount provides a therapeutic effect without causing a substantial cytotoxic effect in the subject.

In some embodiments, a desired response is to inhibit or reduce or prevent EBOV infection. The EBOV infection does not need to be completely eliminated or reduced or prevented for the method to be effective. For example, administration of a therapeutically effective amount of the agent can reduce or inhibit the EBOV infection (for example, as measured by infection of cells, or by number or percentage of subjects infected by EBOV, or by an increase in the survival time of infected subjects) by a desired amount, for example by at least 10%, at least 20%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 98%, or even at least 100% (elimination or prevention of detectable EBOV infection, as compared to a suitable control.

A therapeutically effective amount of an antibody or antigen binding fragment that specifically binds EBOV GP that is administered to a subject will vary depending upon a number of factors associated with that subject, for example the overall health and/or weight of the subject. A therapeutically effective amount encompasses a fractional dose that contributes in combination with previous or subsequent administrations to attaining a therapeutic response. For example, a therapeutically effective amount of an agent can be administered in a single dose, or in several doses, for example daily, during a course of treatment lasting several days or weeks. However, the therapeutically effective amount can depend on the subject being treated, the severity and type of the condition being treated, and the manner of administration. A unit dosage form of the agent can be packaged in a therapeutic amount, or in multiples of the therapeutic amount, for example, in a vial (e.g., with a pierceable lid) or syringe having sterile components.

Transformed: A transformed cell is a cell into which a nucleic acid molecule has been introduced by molecular biology techniques. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of DNA by electroporation, lipofection, and particle gun acceleration.

Treating or preventing a disease: Inhibiting the full development of a disease or condition, for example, in a subject who is at risk of or has an EBOV infection. "Treatment" refers to a therapeutic intervention that ameliorates a sign or symptom of a disease or pathological condition after it has begun to develop. The term "ameliorating," with reference to a disease or pathological condition, refers to any observable beneficial effect of the treatment. The beneficial effect can be evidenced, for example, by a delayed onset of clinical symptoms of the disease in a susceptible subject, a reduction in severity of some or all clinical symptoms of the disease, a slower progression of the disease, a reduction in the viral load, an improvement in the overall health or well-being of the subject, or by other parameters well known in the art that are specific to the particular disease. A "prophylactic" treatment is a treatment administered to a subject who does not exhibit signs of a disease for the purpose of reducing the risk of developing pathology.

The term "reduces" is a relative term, such that an agent reduces a disease or condition (or a symptom of a disease or condition) if the disease or condition is quantitatively diminished following administration of the agent, or if it is diminished following administration of the agent, as compared to a reference agent. Similarly, the term "prevents" does not necessarily mean that an agent completely eliminates the disease or condition, so long as at least one characteristic of the disease or condition is eliminated. Thus, an antibody that reduces or prevents an infection, can, but does not necessarily completely, eliminate such an infection, so long as the infection is measurably diminished, for example, by at least about 50%, such as by at least about 70%, or about 80%, or even by about 90% the infection in the absence of the agent, or in comparison to a reference agent.

Under conditions sufficient for: A phrase that is used to describe any environment that permits a desired activity. In <sup>20</sup> one example the desired activity is formation of an immune complex. In particular examples the desired activity is treatment of EBOV infection.

Vector: Recombinant DNA vectors are vectors having recombinant DNA. A vector can include nucleic acid 25 sequences that permit it to replicate in a host cell, such as an origin of replication. A vector can also include one or more selectable marker genes and other genetic elements known in the art. Viral vectors are recombinant nucleic acid vectors having at least some nucleic acid sequences derived from 30 one or more viruses. In some embodiments, a viral vector is provided that comprises one or more nucleic acid molecules encoding a disclosed antibody or antigen binding fragment that specifically binds to EBOV GP and neutralizes EBOV. In some embodiments, the viral vector can be an adeno-35 associated virus (AAV) vector. A replication deficient viral vector is a vector that requires complementation of one or more regions of the viral genome required for replication due to a deficiency in at least one replication-essential gene function. For example, such that the viral vector does not replicate in typical host cells, especially those in a human 40patient that could be infected by the viral vector in the course of a therapeutic method.

## II. Description of Several Embodiments

Isolated monoclonal antibodies and antigen binding fragments that specifically bind an epitope on EBOV GP protein are provided. The antibodies and antigen binding fragments can be fully human. In several embodiments, the antibodies and antigen binding fragments can be used to neutralize EBOV infection. Also disclosed herein are compositions including the antibodies and antigen binding fragments and a pharmaceutically acceptable carrier. Nucleic acids encoding the antibodies or antigen binding fragments, expression vectors including these nucleic acids, and isolated host cells that express the nucleic acids are also provided.

The antibodies, antigen binding fragments, nucleic acid molecules, host cells, and compositions can be used for research, diagnostic and therapeutic purposes. For example, the monoclonal antibodies and antigen binding fragments can be used to diagnose or treat a subject with an EBOV, or can be administered prophylactically to prevent EBOV infection in a subject. In some embodiments, the antibodies can be used to determine EBOV titer in a subject.

A. Neutralizing Monoclonal Antibodies and Antigen Binding Fragments

This disclosure provides the novel EVB114, EVB114 version 2, EVB100, EVB165, EVB166, and EVB167 antibodies and variants thereof (including antigen binding fragments), which specifically bind to EBOV GP. The disclosed antibodies and antigen binding fragments are surprisingly effective for neutralization of EBOV. For example, as discussed in Example 1, the EVB114 antibody or a combination of EVB114 and EVB100 neutralize EBOV in in vitro assays and were 100% effective in preventing lethality in a primate model of human EBOV infection.

In some embodiments, the antibodies and antigen binding fragments include a  $V_H$  and a  $V_L$  and specifically bind to EBOV GP and neutralize EBOV infection. In several embodiments, the antibody or antigen binding fragment can include a  $V_H$  comprising a HCDR1, a HCDR2 and a HCDR3, and a  $V_L$  comprising a LCDR1, a LCDR2, and a LCDR3, and specifically bind to EBOV GB and neutralize EBOV infection. In several embodiments, the antibody or antigen binding fragment includes a  $V_H$  and a  $V_L$  including the HCDR1, HCDR2, HCDR3, LCDR1, LCDR2, and LCDR3, respectively, of one of the EVB114, EVB114 version 2, EVB100, EVB165, EVB166, or EVB167 antibodies, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, a HCDR3, a LCDR1, a LCDR2, and a LCDR3 comprising amino acid sequences that are at least 90% (for example, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to the amino acid sequences of the CDRs of one of the EVB114, EVB114 version 2, EVB100, EVB165, EVB166, or EVB167 antibodies, and can specifically bind to EBOV GP and neutralize EBOV.

Various CDR numbering schemes (such as the Kabat, Chothia or IMGT numbering schemes) can be used to determine CDR positions. The discussion of monoclonal antibodies below refers to monoclonal antibodies that include a  $V_H$  and a  $V_L$  including CDRs with reference to the IMGT numbering scheme (unless the context indicates otherwise). The amino acid sequence and the CDR positions of the heavy and light chain of the EVB114, EVB114 version 2, EVB100, EVB165, EVB166, or EVB167 antibodies according to the IMGT numbering scheme are shown in Table 1. In several embodiments, an antibody or antigen binding fragment is provided that includes the IMGT CDRs of an antibody listed in Table 1, and can specifically bind to EBOV GP and neutralize EBOV.

TABLE 2

IMGT CDR sequences of EBOV GP specific antibodies						
EVB114 $V_H$						
$\overline{V_H}$	SEQ ID NO: 1 positions	A.A. Sequence	CDR SEQ ID NO			
HCDR1	26-33	GFALRMYD	32			
HCDR2	51-57	VGPSGDT	33			
HCDR3	96-108	VRSDRGVAGLFDS	34			

	T	ABLE 2-continued					
	IMGT CDR sequ	ences of EBOV GP specific antibod	ies				
EVB114 $V_L$							
V <sub>L</sub> LCDR1 LCDR2 LCDR3	SEQ ID NO: 2 positions 27-32 50-52 89-97	A.A. Sequence QAFDNY AAS QNYNSAPLT	CDR SEQ ID NO 35 36 37				
	]	EVB114 version 2 V <sub>H</sub>					
V <sub>H</sub> HCDR1 HCDR2 HCDR3	SEQ ID NO: 28 positions 26-33 51-57 96-108	A.A. Sequence GFALRSYD VGPSGDT VRSDRGVAGLFDS	CDR SEQ ID NO 38 33 34				
	]	EVB114 version 2 $V_L$					
V <sub>L</sub> LCDR1 LCDR2 LCDR3	SEQ ID NO: 29 positions 27-32 50-52 89-97	A.A. Sequence QAFSNY AAS QNYNSAPLT	CDR SEQ ID NO 39 36 37				
		EVB100 $V_H$					
V <sub>H</sub> HCDR1 HCDR2 HCDR3	SEQ ID NO: 3 positions 26-33 51-57 96-114	A.A. Sequence GGSLSSFY IYYSGSP VRASRSYYWGSYRPTAFDS	CDR SEQ ID NO 40 41 42				
		EVBb100 $V_L$					
V <sub>L</sub> LCDR1 LCDR2 LCDR3	SEQ ID NO: 4 positions 26-31 49-51 88-95	A.A. Sequence NLGDKY QDN QTWDSTVV	CDR SEQ ID NO 43 44 45				
		EVB165 $V_H$					
V <sub>H</sub> HCDR1 HCDR2 HCDR3	SEQ ID NO: 19 positions 26-33 51-58 97-117	A.A. Sequence GFRFSDYW IKQDGSGK ARAAPTGSYTNILVDNVHFDY	CDR SEQ ID NO 46 47 48				
		EVB165 $V_L$					
V <sub>L</sub> LCDR1 LCDR2 LCDR3	SEQ ID NO: 20 positions 27-32 50-52 89-97	A.A. Sequence QNIATY AAS QQSYSTPWT	CDR SEQ ID NO 49 36 50				
		EVB166 $V_H$					
V <sub>H</sub> HCDR1 HCDR2 HCDR3	SEQ ID NO: 5 positions 26-33 51-58 97-112	A.A. Sequence GGTLSNYA TIPTLGMS ATMGSADTSFYFYMDV	CDR SEQ ID NO 51 52 53				
		EVB166 $V_L$					
V <sub>L</sub> LCDR1 LCDR2 LCDR3	SEQ ID NO: 6 positions 27-33 51-53 90-98	A.A. Sequence QSVSSSY GTS QQYAYSPFT	CDR SEQ ID NO 54 55 56				
		EVDIO/ V <sub>H</sub>	000 000 00 000				
V <sub>H</sub> HCDR1 HCDR2 HCDR3	SEQ ID NO: 23 positions 26-33 51-58 97-102	A.A. sequence GYTFIQEY GDPENNET TSRKSW	CDR SEQ ID NO 57 58 59				
		EVB167 V <sub>L</sub>					
V <sub>L</sub> LCDR1 LCDR2 LCDR3	SEQ ID NO: 24 positions 27-33 51-53 90-101	A.A. Sequence QSLSSDS GTS QRSGYGMSVTWT	CDR SEQ ID NO 60 55 61				

In several embodiments, the antibody or antigen binding fragment includes IMGT CDRs, such as those listed in Table 1, and can specifically bind to EBOV GP and neutralize EBOV.

**EVB114** 

In some embodiments, the antibody or antigen binding fragment can be based on or derived from the EVB114 antibody, and can specifically bind to EBOV GP and neutralize EBOV. For example, in some embodiments, the antibody or antigen binding fragment can comprise a  $V_H$  10 comprising a HCDR1, a HCDR2, and a HCDR3 of the  $V_H$ set forth as SEQ ID NO: 1 (EVB114  $V_H$ ), and a  $V_L$ comprising a LCDR1, a LCDR2, and a LCDR3 of the  $V_L$  set forth as SEQ ID NO: 2 (EVB114  $V_L$ ), and can specifically bind to EBOV GP and neutralize EBOV. In some embodi- 15 ments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-57, and 96-108 of SEQ ID NO: 1, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In further embodiments, the antibody or 20 antigen binding fragment includes a  $V_r$  including a LCDR1, a LCDR2, and a LCDR3 including amino acids 27-32, 50-52, and 89-97 of SEQ ID NO: 2, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding 25 fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-57, and 96-108 of SEQ ID NO: 1, respectively, and a  $\mathbf{V}_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acids 27-32, 50-52, and 89-97 of SEQ ID NO: 2, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  including a HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 35 98%, or at least 99%) identical to amino acids 26-33, 51-57, and 96-108, respectively, of SEQ ID NO: 1, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a  $V_L$  including a LCDR1, a LCDR2, and a 40 LCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 27-32, 50-52, and 89-97, respectively, of SEQ ID NO: 2, and can specifically bind to EBOV GP and neutralize EBOV. In 45 additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-57, 50 and 96-108, respectively, of SEQ ID NO: 1, and a  $\mathrm{V}_{L}$ including a LCDR1, a LCDR2, and a LCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 27-32, 50-52, and 55 89-97, respectively, of SEQ ID NO: 2, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a  $V_{H}$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to 60 the amino acid sequence set forth as SEQ ID NO: 1, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_L$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the 65 amino acid sequence set forth as SEQ ID NO: 2, and can specifically bind to EBOV GP and neutralize EBOV. In

additional embodiments, the antibody or antigen binding fragment includes a  $V_{H}$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 1, and a  $V_{L}$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 2, and can specifically bind to EBOV GP and neutralize EBOV.

In additional embodiments, the antibody or antigen binding fragment includes a  $V_H$  including the amino acid sequence set forth as one of SEQ ID NO: 1, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_L$  including the amino acid sequence set forth as SEQ ID NO: 2, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  and a  $V_L$  including the amino acid sequences set forth as SEQ ID NOs: 1 and 2, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, and a HCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 32, 33, and 34, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 35, 36, and 37, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, a HCDR3, a LCDR1, a LCDR2, and a LCDR2, and a LCDR2, a HCDR3, a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 32, 33, 34, 35, 36, and 37, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the  $V_H$  of any of the disclosed antibodies or antigen binding fragments based on the EVB114 antibody can further include a M31S substitution (kabat numbering) and/or the  $V_L$  of any of the disclosed antibodies or antigen binding fragments based on the EVB114 antibody can further include a D30S substitution (kabat numbering), and can specifically bind to EBOV GP and neutralize

EBOV. For example, the antibody or antigen binding fragment can includes a HCDR1, a HCDR2, a HCDR3, a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 38, 33, 34, 35, 36, and 37, respectively, SEQ ID NOs: 32, 33, 34, 39, 36, and 37, respectively, or SEQ ID NOs: 38, 33, 34, 39, 36, and 37, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment can include a  $V_H$  comprising the amino acid sequence set forth as SEQ ID NO: 1 further including a M31S substitution, and a  $V_L$  comprising the amino acid sequence set forth as SEQ ID NO: 2 further comprising a D30S substitution.

In some embodiments, the  $V_H$  of any of the disclosed antibodies or antigen binding fragments based on the EVB114 antibody can comprise a value residue at kabat position 96 and a serine residue at kabat position 108. EVB114 Version 2

In some embodiments, the antibody or antigen binding fragment can be based on or derived from the EVB114 version 2 antibody, and can specifically bind to EBOV GP and neutralize EBOV. For example, in some embodiments, the antibody or antigen binding fragment can comprise a  $V_H$  comprising a HCDR1, a HCDR2, and a HCDR3 of the  $V_H$ 

set forth as SEQ ID NO: 28 (EVB114 version 2  $V_H$ ), and a V<sub>L</sub> comprising a LCDR1, a LCDR2, and a LCDR3 of the V<sub>L</sub> set forth as SEQ ID NO: 29 (EVB114 version  $2 V_{\tau}$ ). In some embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-57, and 96-108 of SEQ ID NO: 28, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In further embodiments, the antibody or antigen binding fragment includes a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acids 10 27-32, 50-52, and 89-97 of SEQ ID NO: 29, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a  $V_H$  including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-57, and 96-108 15 of SEQ ID NO: 28, respectively, and a  $\mathrm{V}_{L}$  including a LCDR1, a LCDR2, and a LCDR3 including amino acids 27-32, 50-52, and 89-97 of SEQ ID NO: 29, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding 20 fragment includes a  $V_{FF}$  including a

HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-57, and 96-108, respectively, of SEQ ID 25 NO: 28, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a  $\mathbf{V}_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, 30 at least 98%, or at least 99%) identical to amino acids amino acids 27-32, 50-52, and 89-97, respectively, of SEQ ID NO: 29, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a  $\mathrm{V}_H$  including a HCDR1, a  $_{\rm 35}$ HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-57, and 96-108, respectively, of SEQ ID NO: 28, and a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including 40 amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 27-32, 50-52, and 89-97, respectively, of SEQ ID NO: 29, and can specifically bind to EBOV GP and neutralize EBOV. 45

In some embodiments, the antibody or antigen binding fragment includes a  $\mathrm{V}_{H}$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 28, and can specifically bind to EBOV GP and neutralize EBOV. In 50 more embodiments, the antibody or antigen binding fragment includes a V<sub>L</sub> including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 29, and can specifically bind to EBOV GP and neutralize EBOV. In 55 additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 28, and a  $V_L$  including an amino acid sequence at least 80%, 90%, 60 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 29, and can specifically bind to EBOV GP and neutralize EBOV.

In additional embodiments, the antibody or antigen binding fragment includes a  $V_H$  including the amino acid sequence set forth as one of SEQ ID NO: 28, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_L$  including the amino acid sequence set forth as SEQ ID NO: 29, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  and a  $V_L$  including the amino acid sequences set forth as SEQ ID NOs: 28 and 29, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, and a HCDR3, comprising the amino acid sequences set forth as SEQ ID NOS: 38, 33, and 34, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOS: 39, 36, and 37, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOS: 39, 36, and 37, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, a HCDR3, a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOS: 38, 33, 34, 39, 36, and 37, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the V<sub>H</sub> of any of the disclosed antibodies or antigen binding fragments based on the EVB114 version 2 antibody can further include a S31M substitution (kabat numbering) and/or the V<sub>L</sub> of any of the disclosed antibodies or antigen binding fragments based on the EVB114 antibody can further include a S30D substitution (kabat numbering). For example, in some embodiments, the antibody or antigen binding fragment can include a V<sub>H</sub> comprising the amino acid sequence set forth as SEQ ID NO: 28 further including a S31M substitution, and a V<sub>L</sub> comprising the amino acid sequence set forth as SEQ ID NO: 29 further comprising a S30D substitution.

EVB100

In some embodiments, the antibody or antigen binding fragment can be based on or derived from the EVB100 antibody, and can specifically bind to EBOV GP and neutralize EBOV. For example, in some embodiments, the antibody or antigen binding fragment can comprise a  $V_H$ comprising a HCDR1, a HCDR2, and a HCDR3 of the  $V_H$ set forth as SEQ ID NO: 3 (EVB100  $V_H$ ), and a  $V_L$ comprising a LCDR1, a LCDR2, and a LCDR3 of the  $V_L$  set forth as SEQ ID NO: 4 (EVB100  $V_L$ ). In some embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-57, and 96-114 of SEQ ID NO: 3, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In further embodiments, the antibody or antigen binding fragment includes a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acids 26-31, 49-51, and 88-95 of SEQ ID NO: 4, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a  $\mathbf{V}_{H}$  including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-57, and 96-114 of SEQ ID NO: 3, respectively, and a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acids 26-31, 49-51, and 88-95 of SEQ ID NO: 4, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  including a HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-57, and 96-114, respectively, of SEQ ID NO: 3, and can spe-

cifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a V<sub>1</sub> including a LCDR1, a LCDR2, and a LCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 5 99%) identical to amino acids amino acids 26-31, 49-51, and 88-95, respectively, of SEQ ID NO: 4, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-57, and 96-114, respectively, of SEQ ID NO: 3, and a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acid 15 sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 26-31, 49-51, and 88-95, respectively, of SEQ ID NO: 4, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 3, and can specifically bind to EBOV GP and neutralize EBOV. In more 25 embodiments, the antibody or antigen binding fragment includes a  $V_L$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 4, and can specifically bind to EBOV GP and neutralize EBOV. In 30 additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 3, and a  $V_{L}$  including an amino acid sequence at least 80%, 90%, 35 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 4, and can specifically bind to EBOV GP and neutralize EBOV.

In additional embodiments, the antibody or antigen binding fragment includes a  $V_{H}$  including the amino acid 40 sequence set forth as one of SEQ ID NO: 3, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_{L}$  including the amino acid sequence set forth as SEQ ID NO: 4, and can specifically bind to EBOV GP and 45 neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a  $V_{H}$  and a  $V_{L}$  including the amino acid sequences set forth as SEQ ID NOs: 3 and 4, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, and a HCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 40, 41, and 42, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, 55 the antibody or antigen binding fragment includes a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 43, 44, and 45, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or 60 antigen binding fragment includes a HCDR1, a HCDR2, a HCDR3, a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 40, 41, 42, 43, 44, and 45, respectively, and can specifically bind to EBOV GP and neutralize EBOV. 65

In some embodiments, the  $V_H$  of any of the disclosed antibodies or antigen binding fragments based on the

EVB100 antibody can comprise a value residue at kabat position 96, a tyrosine residue at kabat position 103 and a serine residue at kabat position 114. EVB165

In some embodiments, the antibody or antigen binding fragment can be based on or derived from the EVB165 antibody, and can specifically bind to EBOV GP and neutralize EBOV. For example, in some embodiments, the antibody or antigen binding fragment can comprise a  $V_H$  comprising a HCDR1, a

HCDR2, and a HCDR3 of the  $V_H$  set forth as SEQ ID NO: 19 (EVB165  $V_H$ ), and a  $V_L$  comprising a LCDR1, a LCDR2, and a LCDR3 of the  $V_L$  set forth as SEQ ID NO: 20 (EVB165  $V_L$ ). In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-58, and 97-117 of SEQ ID NO: 19, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In further embodiments, the antibody or antigen binding frag-20 ment includes a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acids 27-32, 50-52, and 89-97 of SEQ ID NO: 20, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-58, and 97-117 of SEQ ID NO: 19, respectively, and a V<sub>L</sub> including a LCDR1, a LCDR2, and a LCDR3 including amino acids 27-32, 50-52, and 89-97 of SEQ ID NO: 20, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-58, and 97-117, respectively, of SEQ ID NO: 19, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 27-32, 50-52, and 89-97, respectively, of SEQ ID NO: 20, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-58, and 97-117, respectively, of SEQ ID NO: 19, and a  $V_L$ including a LCDR1, a LCDR2, and a LCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 27-32, 50-52, and 89-97, respectively, of SEQ ID NO: 20, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a  $V_{H}$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 19, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_{L}$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 20, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a  $V_{L}$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 20, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding

-5

fragment includes a V<sub>H</sub> including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 19, and a V<sub>L</sub> including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 20, and can specifically bind to EBOV GP and neutralize EBOV.

In additional embodiments, the antibody or antigen binding fragment includes a  $V_H$  including the amino acid sequence set forth as one of SEQ ID NO: 19, and can 10 specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_L$  including the amino acid sequence set forth as SEQ ID NO: 20, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or 15 antigen binding fragment includes a  $V_H$  and a  $V_L$  including the amino acid sequences set forth as SEQ ID NOs: 19 and 20, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding 20 fragment includes a HCDR1, a HCDR2, and a HCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 46, 47, and 48, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a LCDR1, 25 a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 49, 36, and 50, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, a 30 HCDR3, a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 46, 47, 48, 49, 36, and 50, respectively, and can specifically bind to EBOV GP and neutralize EBOV. **EVB166** 35

In some embodiments, the antibody or antigen binding fragment can be based on or derived from the EVB166 antibody, and can specifically bind to EBOV GP and neutralize EBOV. For example, in some embodiments, the antibody or antigen binding fragment can comprise a  $V_H$  40 comprising a HCDR1, a HCDR2, and a HCDR3 of the  $V_H$ set forth as SEQ ID NO: 5 (EVB166  $V_H$ ), and a  $V_L$ comprising a LCDR1, a LCDR2, and a LCDR3 of the  $V_L$  set forth as SEQ ID NO: 6 (EVB166 V<sub>L</sub>). In some embodiments, the antibody or antigen binding fragment includes a 45 V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-58, and 97-112 of SEO ID NO: 5, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In further embodiments, the antibody or antigen binding fragment includes a  $V_L$  including a LCDR1, 50 a LCDR2, and a LCDR3 including amino acids 27-33, 51-53, and 90-98 of SEQ ID NO: 6, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and 55 a HCDR3 including amino acids 26-33, 51-58, and 97-112 of SEQ ID NO: 5, respectively, and a  $\mathrm{V}_{L}$  including a LCDR1, a LCDR2, and a LCDR3 including amino acids 27-33, 51-53, and 90-98 of SEQ ID NO: 6, respectively, and can specifically bind to EBOV GP and neutralize EBOV. 60

In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  including a HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-58, 65 and 97-112, respectively, of SEQ ID NO: 5, and can specifically bind to EBOV GP and neutralize EBOV. In some

embodiments, the antibody or antigen binding fragment includes a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 27-33, 51-53, and 90-98, respectively, of SEQ ID NO: 6, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-58, and 97-112, respectively, of SEQ ID NO: 5, and a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 27-33, 51-53, and 90-98, respectively, of SEQ ID NO: 6, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 5, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_{L}$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 6, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_L$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 62, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 5, and a  $V_L$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 6, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 5, and a  $V_L$ including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 62, and can specifically bind to EBOV GP and neutralize EBOV.

In additional embodiments, the antibody or antigen binding fragment includes a  $\mathbf{V}_{H}$  including the amino acid sequence set forth as one of SEQ ID NO: 5, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a V<sub>L</sub> including the amino acid sequence set forth as SEQ ID NO: 6, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_L$  including the amino acid sequence set forth as SEQ ID NO: 62, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  and a  $V_L$  including the amino acid sequences set forth as SEQ ID NOs: 5 and 6, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  and a  $V_L$  including the amino acid sequences set forth as SEQ ID NOs: 5 and 62, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, and a HCDR3, comprising the amino acid sequences set forth as SEQ ID -5 NOs: 51, 52, and 53, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 54, 55, and 56, 10 respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, a HCDR3, a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 51, 52, 15 53, 54, 55, and 56, respectively, and can specifically bind to EBOV GP and neutralize EBOV. **EVB167** 

In some embodiments, the antibody or antigen binding fragment can be based on or derived from the EVB167 20 antibody, and can specifically bind to EBOV GP and neutralize EBOV. For example, in some embodiments, the antibody or antigen binding fragment can comprise a  $\mathbf{V}_{\!H}$ comprising a HCDR1, a HCDR2, and a HCDR3 of the  $\mathrm{V}_{H}$ set forth as SEQ ID NO: 23 (EVB167  $\mathrm{V}_{H}\mathrm{)},$  and a  $\mathrm{V}_{L}$  25 comprising a LCDR1, a LCDR2, and a LCDR3 of the  $V_L$  set forth as SEQ ID NO: 24 (EVB167 V<sub>L</sub>). In some embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-58, and 97-102 of SEQ ID NO: 23, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In further embodiments, the antibody or antigen binding fragment includes a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acids 27-33, 51-53, and 90-101 of SEQ ID NO: 24, respectively, and can 35 specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acids 26-33, 51-58, and 97-102 of SEQ ID NO: 23, respectively, and a  $V_L$  including a 40 LCDR1, a LCDR2, and a LCDR3 including amino acids 27-33, 51-53, and 90-101 of SEQ ID NO: 24, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and 45 a HCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-58, and 97-102, respectively, of SEQ ID NO: 23, and can specifically bind to EBOV GP and neutralize EBOV. In 50 some embodiments, the antibody or antigen binding fragment includes a  $V_L$  including a LCDR1, a LCDR2, and a LCDR3 including amino acid sequences at least 90% (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 27-33, 55 51-53, and 90-101, respectively, of SEQ ID NO: 24, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a V<sub>H</sub> including a HCDR1, a HCDR2, and a HCDR3 including amino acid sequences at least 90% 60 (such as at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids 26-33, 51-58, and 97-102, respectively, of SEQ ID NO: 23, and a  $V_{T}$ including a LCDR1, a LCDR2, and a LCDR3 including amino acid sequences at least 90% (such as at least 95%, at 65 least 96%, at least 97%, at least 98%, or at least 99%) identical to amino acids amino acids 27-33, 51-53, and

90-101, respectively, of SEQ ID NO: 24, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEO ID NO: 23, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_{T}$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 24, and can specifically bind to EBOV GP and neutralize EBOV. In additional embodiments, the antibody or antigen binding fragment includes a  $V_H$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 23, and a  $V_L$  including an amino acid sequence at least 80%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence set forth as SEQ ID NO: 24, and can specifically bind to EBOV GP and neutralize EBOV.

In additional embodiments, the antibody or antigen binding fragment includes a  $V_H$  including the amino acid sequence set forth as one of SEQ ID NO: 23, and can specifically bind to EBOV GP and neutralize EBOV. In more embodiments, the antibody or antigen binding fragment includes a  $V_L$  including the amino acid sequence set forth as SEQ ID NO: 24, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a  $V_H$  and a  $V_L$  including the amino acid sequences set forth as SEQ ID NOs: 23 and 24, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, and a HCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 57, 58, and 59, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a LCDR1, a LCDR2, and a LCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 60, 55, and 61, respectively, and can specifically bind to EBOV GP and neutralize EBOV. In some embodiments, the antibody or antigen binding fragment includes a HCDR1, a HCDR2, a HCDR3, a LCDR1, a LCDR2, and a LCDR2, and a LCDR2, and a HCDR3, comprising the amino acid sequences set forth as SEQ ID NOs: 57, 58, 59, 60, 55, and 61, respectively, and can specifically bind to EBOV GP and neutralize EBOV.

1. Additional Description of Antibodies and Antigen Binding Fragments

The antibody or antigen binding fragment can be a human antibody or fragment thereof. Chimeric antibodies are also provided. The antibody or antigen binding fragment can include any suitable framework region, such as (but not limited to) a human framework region. Human framework regions, and mutations that can be made in a human antibody framework regions, are known in the art (see, for example, in U.S. Pat. No. 5,585,089, which is incorporated herein by reference). Alternatively, a heterologous framework region, such as, but not limited to a mouse or monkey framework region, can be included in the heavy or light chain of the antibodies. (See, for example, Jones et al., Nature 321:522, 1986; Riechmann et al., Nature 332:323, 1988; Verhoeyen et al., Science 239:1534, 1988; Carter et al., Proc. Natl. Acad. Sci. U.S.A. 89:4285, 1992; Sandhu, Crit. Rev. Biotech. 12:437, 1992; and Singer et al., J. Immunol. 150:2844, 1993.)

The antibody can be of any isotype. The antibody can be, for example, an IgM or an IgG antibody, such as IgG<sub>1</sub>, IgG<sub>2</sub>,  $IgG_3$ , or  $IgG_4$ . The class of an antibody that specifically binds EBOV GP can be switched with another. In one aspect, a nucleic acid molecule encoding  $V_L$  or  $V_H$  is isolated using 5 methods well-known in the art, such that it does not include any nucleic acid sequences encoding the constant region of the light or heavy chain, respectively. A nucleic acid molecule encoding  $V_L$  or  $V_H$  is then operatively linked to a nucleic acid sequence encoding a  $C_L$  or  $C_H$  from a different 10 class of immunoglobulin molecule. This can be achieved using a vector or nucleic acid molecule that comprises a  $C_L$ or  $C_H$  chain, as known in the art. For example, an antibody that specifically binds EBOV GP, that was originally IgM may be class switched to an IgG. Class switching can be 15 used to convert one IgG subclass to another, such as from  $IgG_1$  to  $IgG_2$ ,  $IgG_3$ , or  $IgG_4$ .

In some examples, the disclosed antibodies are oligomers of antibodies, such as dimers, trimers, tetramers, pentamers, hexamers, septamers, octomers and so on. (a) Binding Affinity

In several embodiments, the antibody or antigen binding fragment can specifically bind EBOV GP with an affinity (e.g., measured by  $K_d$ ) of no more than  $1.0 \times 10^{-8}$  M, no more than  $5.0 \times 10^{-8}$  M, no more than  $1.0 \times 10^{-11}$  M, no more than 25  $5.0 \times 10^{-9}$  M, no more than  $1.0 \times 10^{-10}$  M, no more than  $5.0 \times 10^{-10}$  M, or no more than  $1.0 \times 10^{-11}$  M. K<sub>d</sub> can be measured, for example, by a radiolabeled antigen binding assay (RIA) performed with the Fab version of an antibody of interest and its antigen using known methods. In one 30 assay, solution binding affinity of Fabs for antigen is measured by equilibrating Fab with a minimal concentration of (<sup>125</sup>I)-labeled antigen in the presence of a titration series of unlabeled antigen, then capturing bound antigen with an anti-Fab antibody-coated plate (see, e.g., Chen et al., J. Mol. 35 Biol. 293:865-881, 1999, which is incorporated by reference herein in its entirety). To establish conditions for the assay, MICROTITER® multi-well plates (Thermo Scientific) are coated overnight with 5 µg/ml of a capturing anti-Fab antibody (Cappel Labs) in 50 mM sodium carbonate (pH 40 9.6), and subsequently blocked with 2% (w/v) bovine serum albumin in PBS for two to five hours at room temperature (approximately 23° C.). In a non-adsorbent plate (Nunc #269620), 100  $\mu$ M or 26 pM [<sup>125</sup>I]-antigen are mixed with serial dilutions of a Fab of interest (e.g., consistent with 45 assessment of the anti-VEGF antibody, Fab-12, in Presta et al., Cancer Res. 57:4593-4599 (1997)). The Fab of interest is then incubated overnight; however, the incubation may continue for a longer period (e.g., about 65 hours) to ensure that equilibrium is reached. Thereafter, the mixtures are 50 transferred to the capture plate for incubation at room temperature (e.g., for one hour). The solution is then removed and the plate washed eight times with 0.1% polysorbate 20 (TWEEN-20®) in PBS. When the plates have dried, 150 µl/well of scintillant (MICROSCINT-20™; Pack- 55 ard) is added, and the plates are counted on a TOP-COUNT<sup>™</sup> gamma counter (Packard) for ten minutes. Concentrations of each Fab that give less than or equal to 20% of maximal binding are chosen for use in competitive binding assays. 60

In another assay,  $K_d$  can be measured using surface plasmon resonance assays using a BIACORE®-2000 or a BIACORE®-3000 (BIAcore, Inc., Piscataway, N.J.) at 25° C. with immobilized antigen CMS chips at ~10 response units (RU). Briefly, carboxymethylated dextran biosensor 65 chips (CMS, BIACORE®, Inc.) are activated with N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide hydrochloride

44

(EDC) and N-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 µg/ml (~0.2 µM) before injection at a flow rate of 5 l/minute to achieve approximately 10 response units (RU) of coupled protein. Following the injection of antigen, 1 M ethanolamine is injected to block unreacted groups. For kinetics measurements, twofold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with 0.05% polysorbate 20 (TWEEN-20™) surfactant (PBST) at 25° C. at a flow rate of approximately 25 l/min. Association rates  $(k_{on})$  and dissociation rates  $(k_{off})$  are calculated using a simple one-to-one Langmuir binding model (BIACORE® Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant  $(K_d)$  is calculated as the ratio koff/kon. See, e.g., Chen et al., J. Mol. *Biol.* 293:865-881 (1999). If the on-rate exceeds  $10^6 \text{ M}^{-1} \text{ s}^{-1}$ by the surface plasmon resonance assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase or decrease in fluorescence emission intensity (excitation=295 nm; emission=340 nm, 16 nm band-pass) at 25° C. of a 20 nM anti-antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such as a stop-flow equipped spectrophometer (Aviv Instruments) or a 8000-series SLM-AMINCO<sup>™</sup> spectrophotometer (ThermoSpectronic) with a stirred cuvette. (b) Neutralization

In several embodiments, the antibodies and antigen binding fragments disclosed herein can neutralize EBOV infection by at least two, at least three, at least four, or at least five strains of EBOV, such as the Bundibugyo (BDBV), Reston (RESTV), Sudan (SUDV), Tai Forest (TAFV), and Zaire (ZEBOV), with an IC50 of less than 50 µg/ml. In more embodiments, the antibodies and antigen binding fragments disclosed herein can neutralize EBOV infection by at least two, at least three, at least four, or at least five strains of EBOV, such as the BDBV, RESTV, SUDV, TAFV, and ZEBOV, with an IC50 of less than 10 µg/ml. In several embodiments the antibodies and antigen binding fragments disclosed herein can neutralize infection by ZEBOV, with an IC50 of less than 50 µg/ml or less than 10 µg/ml. Exemplary methods of assaying EBOV neutralization are provided in the Examples. In some embodiments, neutralization assays can be performed using a single-round EBOV GP-pseudoviruses infection of 293-T cells. In some embodiments, methods to assay for neutralization activity includes a singlecycle infection assay as described in Martin et al. (2003) Nature Biotechnology 21:71-76. In this assay, the level of viral activity is measured via a selectable marker whose activity is reflective of the amount of viable virus in the sample, and the  $IC_{50}$  is determined.

(c) Multispecific Antibodies

In some embodiments, the antibody or antigen binding fragment is included on a multispecific antibody, such as a bi-specific antibody. Such multispecific antibodies can be produced by known methods, such as crosslinking two or more antibodies, antigen binding fragments (such as scFvs) of the same type or of different types. Exemplary methods of making multispecific antibodies include those described in PCT Pub. No. WO2013/163427, which is incorporated by reference herein in its entirety. Suitable crosslinkers include those that are heterobifunctional, having two distinctly reactive groups separated by an appropriate spacer (such as m-maleimidobenzoyl-N-hydroxysuccinimide ester) or

25

50

homobifunctional (such as disuccinimidyl suberate). Such linkers are available from Pierce Chemical Company, Rockford, Ill.

Various types of multi-specific antibodies are known. Bispecific single chain antibodies can be encoded by a single nucleic acid molecule. Examples of bispecific single chain antibodies, as well as methods of constructing such antibodies are known in the art (see, e.g., U.S. Pat. Nos. 8,076,459, 8,017,748, 8,007,796, 7,919,089, 7,820,166, 7,635,472, 7,575,923, 7,435,549, 7,332,168, 7,323,440, 7,235,641, 7,229,760, 7,112,324, 6,723,538, incorporated by reference herein). Additional examples of bispecific single chain antibodies can be found in PCT application No. WO 99/54440; Mack, J. Immunol., 158:3965-3970, 1997; Mack, PNAS, 92:7021-7025, 1995; Kufer, Cancer Immunol. Immunother., 45:193-197, 1997; Loffler, Blood, 95:2098-2103, 2000; and Bruhl, J. Immunol., 166:2420-2426, 2001. Production of bispecific Fab-scFv ("bibody") molecules are described, for example, in Schoonjans et al. (J. Immunol. 20 and Vr chains connected by a peptide linker. These single-165:7050-57, 2000) and Willems et al. (J Chromatogr B Analyt Technol Biomed Life Sci. 786:161-76, 2003). For bibodies, a scFv molecule can be fused to one of the VL-CL (L) or VH-CH1 chains, e.g., to produce a bibody one scFv is fused to the C-term of a Fab chain.

(d) Fragments

Antigen binding fragments are encompassed by the present disclosure, such as Fab,  $F(ab')_2$ , and Fv which include a heavy chain and light chain variable region and specifically bind EBOV GP. These antibody fragments retain the ability to selectively bind with the antigen and are "antigen-binding" fragments. Non-limiting examples of such fragments include:

(1) Fab, the fragment which contains a monovalent antigen-binding fragment of an antibody molecule, can be produced by digestion of whole antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain;

(2) Fab', the fragment of an antibody molecule can be  $_{40}$ obtained by treating whole antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain; two Fab' fragments are obtained per antibody molecule;

(3)  $(Fab')_2$ , the fragment of the antibody that can be 45 obtained by treating whole antibody with the enzyme pepsin without subsequent reduction;  $F(ab')_2$  is a dimer of two Fab' fragments held together by two disulfide bonds;

(4) Fv, a genetically engineered fragment containing the  $V_H$  and  $V_L$  expressed as two chains; and

(5) Single chain antibody (such as scFv), defined as a genetically engineered molecule containing the variable region of the light chain, the variable region of the heavy chain, linked by a suitable polypeptide linker as a genetically fused single chain molecule. A scFv is a fusion protein in 55 which a  $\tilde{V}_L$  of an immunoglobulin and a  $V_H$  of an immunoglobulin are bound by a linker (see, e.g., Ahmad et al., Clin. Dev. Immunol., 2012, doi:10.1155/2012/980250; Marbry, IDrugs, 13:543-549, 2010). The intramolecular orientation of the  $V_H$ -domain and the  $V_L$ -domain in a scFv, is not 60 decisive for the provided antibodies (e.g., for the provided multispecific antibodies). Thus, scFvs with both possible arrangements ( $V_H$ -domain-linker domain- $V_L$ -domain;  $V_L$ domain-linker domain- $V_H$ -domain) may be used.

(6) A dimer of a single chain antibody (scFV<sub>2</sub>), defined as 65 a dimer of a scFV. This has also been termed a "miniantibody."

Methods of making these fragments are known in the art (see for example, Harlow and Lane, Antibodies: A Laboratory Manual, 2<sup>nd</sup>, Cold Spring Harbor Laboratory, New York. 2013).

In some embodiments, the antigen binding fragment can be an Fv antibody, which is typically about 25 kDa and contain a complete antigen-binding site with three CDRs per each heavy chain and each light chain. To produce Fv antibodies, the  $V_H$  and the  $V_L$  can be expressed from two individual nucleic acid constructs in a host cell.

If the  $V_H$  and the  $V_L$  are expressed non-contiguously, the chains of the Fv antibody are typically held together by noncovalent interactions. However, these chains tend to dissociate upon dilution, so methods have been developed to crosslink the chains through glutaraldehyde, intermolecular disulfides, or a peptide linker. Thus, in one example, the Fv can be a disulfide stabilized Fv (dsFv), wherein the  $V_H$  and the  $V_L$  are chemically linked by disulfide bonds.

In an additional example, the Fv fragments comprise  $V_H$ chain antigen binding proteins (scFv) are prepared by constructing a nucleic acid molecule encoding the  $V_H$  and  $V_L$ domains connected by an oligonucleotide. The nucleic acid molecule is inserted into an expression vector, which is subsequently introduced into a host cell such as a mammalian cell. The recombinant host cells synthesize a single polypeptide chain with a linker peptide bridging the two V domains. Methods for producing scFvs are known in the art (see Whitlow et al., Methods: a Companion to Methods in Enzymology, Vol. 2, page 97, 1991; Bird et al., Science 242:423, 1988; U.S. Pat. No. 4,946,778; Pack et al., Bio/ Technology 11:1271, 1993; Ahmad et al., Clin. Dev. Immunol., 2012, doi:10.1155/2012/980250; Marbry, IDrugs, 13:543-549, 2010). Dimers of a single chain antibody (scFV<sub>2</sub>), are also contemplated.

Antigen binding fragments can be prepared by proteolytic hydrolysis of the antibody or by expression in a host cell (such as an E. coli cell) of DNA encoding the fragment. Antigen binding fragments can also be obtained by pepsin or papain digestion of whole antibodies by conventional methods. For example, antigen binding fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a 5S fragment denoted  $F(ab')_2$ . This fragment can be further cleaved using a thiol reducing agent, and optionally a blocking group for the sulfhydryl groups resulting from cleavage of disulfide linkages, to produce 3.58 Fab' monovalent fragments. Alternatively, an enzymatic cleavage using pepsin produces two monovalent Fab' fragments and an Fc fragment directly (see U.S. Pat. No. 4,036,945 and U.S. Pat. No. 4,331,647, and references contained therein; Nisonhoff et al., Arch. Biochem. Biophys. 89:230, 1960; Porter, Biochem. J. 73:119, 1959; Edelman et al., Methods in Enzymology, Vol. 1, page 422, Academic Press, 1967; and Coligan et al. at sections 2.8.1-2.8.10 and 2.10.1-2.10.4).

Other methods of cleaving antibodies, such as separation of heavy chains to form monovalent light-heavy chain fragments, further cleavage of fragments, or other enzymatic, chemical, or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody.

Antigen binding single  $\mathrm{V}_{H}$  domains, called domain antibodies (dAb), have also been identified from a library of murine  $V_H$  genes amplified from genomic DNA of immunized mice (Ward et al. Nature 341:544-546, 1989). Human single immunoglobulin variable domain polypeptides capable of binding antigen with high affinity have also been described (see, for example, PCT Publication Nos. WO

2005/035572 and WO 2003/002609). The CDRs disclosed herein can also be included in a dAb.

In some embodiments, one or more of the heavy and/or light chain complementarity determining regions (CDRs) from a disclosed antibody (such as the EVB100, EVB114, or 5 EVB166 antibody) is expressed on the surface of another protein, such as a scaffold protein. The expression of domains of antibodies on the surface of a scaffolding protein are known in the art (see e.g. Liu et al., *J. Virology* 85(17): 8467-8476, 2011). Such expression creates a chimeric pro-10 tein that retains the binding for EBOV GP. In some specific embodiments, one or more of the heavy chain CDRs is grafted onto a scaffold protein, such as one or more of heavy chain CDR1, CDR2, and/or CDR3. One or more CDRs can also be included in a diabody or another type of single chain 15 antibody molecule.

(e) Additional Antibodies that Bind to the EVB114, EVB100, EVB165, EVB166, or EVB167 Epitope on EBOV GP.

Also included are antibodies that bind to the same epitope 20 on EBOV GP to which the EVB114, EVB100, EVB165, EVB166, or EVB167 antibody binds. Antibodies that bind to such an epitope can be identified based on their ability to cross-compete (for example, to competitively inhibit the binding of, in a statistically significant manner) with the 25 EVB114, EVB100, EVB165, EVB166, or EVB167 antibodies provided herein in EBOV GP binding assays (such as those described in the Examples). An antibody "competes" for binding when the competing antibody inhibits EBOV GP binding of the EVB114, EVB100, EVB165, EVB166, or 30 EVB167 antibody by more than 50%, in the presence of competing antibody concentrations higher than  $10^6 \times K_D$  of the competing antibody. In a certain embodiment, the antibody that binds to the same epitope on EBOV GP as the EVB114, EVB100, EVB165, EVB166, or EVB167 antibody 35 is a human monoclonal antibody. Human antibodies that bind to the same epitope on EBOV GP to which the EVB114, EVB100, EVB165, EVB166, or EVB167 antibody binds can be produced using various techniques known in the art. Human antibodies are described generally in van 40 Dijk and van de Winkel, Curr. Opin. Pharmacol. 5: 368-74 (2001) and Lonberg, Curr. Opin. Immunol. 20:450-459 (2008). Such antibodies may be prepared, for example, by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact 45 antibodies with human variable regions in response to antigenic challenge. Such animals typically contain all or a portion of the human immunoglobulin loci, which replace the endogenous immunoglobulin loci, or which are present extrachromosomally or integrated randomly into the ani- 50 mal's chromosomes. In such transgenic mice, the endogenous immunoglobulin loci have generally been inactivated. For review of methods for obtaining human antibodies from transgenic animals, see Lonberg, Nat. Biotech. 23:1117-1125 (2005). See also, e.g., U.S. Pat. Nos. 6,075,181 and 55 6,150,584 describing XENOMOUSE<sup>™</sup> technology; U.S. Pat. No. 5,770,429 describing HUMAB® technology; U.S. Pat. No. 7,041,870 describing K-M MOUSE® technology, and U.S. Patent Application Publication No. US 2007/ 0061900, describing VELOCIMOUSE® technology). 60 Human variable regions from intact antibodies generated by such animals may be further modified, e.g., by combining with a different human constant region.

Human antibodies that bind to the same epitope on EBOV GP to which the EVB114, EVB100, EVB165, EVB166, or 65 EVB167 antibody binds can also be made by hybridomabased methods. Human myeloma and mouse-human hetero-

myeloma cell lines for the production of human monoclonal antibodies have been described. (See, e.g., Kozbor J. Immunol., 133: 3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, pp. 51-63 (Marcel Dekker, Inc., New York, 1987); and Boerner et al., J. Immunol., 147: 86 (1991).) Human antibodies generated via human B-cell hybridoma technology are also described in Li et al., Proc. Natl. Acad. Sci. USA, 103:3557-3562 (2006). Additional methods include those described, for example, in U.S. Pat. No. 7,189,826 (describing production of monoclonal human IgM antibodies from hybridoma cell lines) and Ni, Xiandai Mianyixue, 26(4):265-268 (2006) (describing human-human hybridomas). Human hybridoma technology (Trioma technology) is also described in Vollmers and Brandlein, Histology and Histopathology, 20(3):927-937 (2005) and Vollmers and Brandlein, Methods and Findings in Experimental and Clinical Pharmacology, 27(3): 185-91 (2005). Human antibodies may also be generated by isolating Fv clone variable domain sequences selected from human-derived phage display libraries. Such variable domain sequences may then be combined with a desired human constant domain.

Antibodies and antigen binding fragments that specifically bind to the same epitope on EBOV GP as EVB114, EVB100, EVB165, EVB166, or EVB167 can also be isolated by screening combinatorial libraries for antibodies with the desired binding characteristics. For example, a variety of methods are known in the art for generating phage display libraries and screening such libraries for antibodies possessing the desired binding characteristics. Such methods are reviewed, e.g., in Hoogenboom et al. in Methods in Molecular Biology 178:1-37 (O'Brien et al., ed., Human Press, Totowa, N.J., 2001) and further described, e.g., in the McCafferty et al., Nature 348:552-554; Clackson et al., Nature 352: 624-628 (1991); Marks et al., J. Mol. Biol. 222: 581-597 (1992); Marks and Bradbury, in Methods in Molecular Biology 248:161-175 (Lo, ed., Human Press, Totowa, N.J., 2003); Sidhu et al., J. Mol. Biol. 338(2): 299-310 (2004); Lee et al., J. Mol. Biol. 340(5): 1073-1093 (2004); Fellouse, Proc. Natl. Acad. Sci. USA 101(34): 12467-12472 (2004); and Lee et al., J. Immunol. Methods 284(1-2): 119-132 (2004).

In certain phage display methods, repertoires of  $V_H$  and V<sub>L</sub> genes are separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be screened for antigen-binding phage as described in Winter et al., Ann. Rev. Immunol., 12: 433-455 (1994). Phage typically display antibody fragments, either as singlechain Fv (scFv) fragments or as Fab fragments. Libraries from immunized sources provide high-affinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned (e.g., from human) to provide a single source of antibodies to a wide range of non-self and also self antigens without any immunization as described by Griffiths et al., EMBO J, 12: 725-734 (1993). Finally, naive libraries can also be made synthetically by cloning unrearranged V-gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to accomplish rearrangement in vitro, as described by Hoogenboom and Winter, J. Mol. Biol., 227: 381-388 (1992). Patent publications describing human antibody phage libraries include, for example: U.S. Pat. No. 5,750,373, and US Patent Publication Nos. 2005/0079574, 2005/0119455, 2005/0266000, 2007/0117126, 2007/ 0160598, 2007/0237764, 2007/0292936, and 2009/ 0002360.

(f) Variants

In certain embodiments, amino acid sequence variants of the antibodies provided herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. Amino 5 acid sequence variants of an antibody may be prepared by introducing appropriate modifications into the nucleotide sequence encoding the antibody, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or substitutions of residues within 10 the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution can be made to arrive at the final construct, provided that the final construct possesses the desired characteristics, e.g., antigen-binding.

In certain embodiments, antibody variants having one or 15 more amino acid substitutions are provided. Sites of interest for substitutional mutagenesis include the CDRs and the framework regions. Amino acid substitutions may be introduced into an antibody of interest and the products screened for a desired activity, e.g., retained/improved antigen bind-20 ing, decreased immunogenicity, or improved ADCC or CDC.

The variants typically retain amino acid residues necessary for correct folding and stabilizing between the  $V_H$  and the  $V_L$  regions, and will retain the charge characteristics of 25 the residues in order to preserve the low pI and low toxicity of the molecules. Amino acid substitutions can be made in the  $V_H$  and the  $V_L$  regions to increase yield. or removed. Where the antibody comprises an Fc region, the carbohydrate attached thereto may be altered. Native antibodies produced by mammalian cells typically comprise a branched, biantennary oligosaccharide that is generally attached by an N-linkage to Asn297 of the CH<sub>2</sub> domain of

In some embodiments, the heavy chain of the antibody includes up to 10 (such as up to 1, up to 2, up to 3, up to 4, 30 up to 5, up to 6, up to 7, up to 8, or up to 9) amino acid substitutions (such as conservative amino acid substitutions) compared to the amino acid sequence set forth as one of SEQ ID NOS: 1, 3, 5, 19, 23, or 28. In some embodiments, the light chain of the antibody includes up to 10 (such as up 35 to 1, up to 2, up to 3, up to 4, up to 5, up to 6, up to 7, up to 8, or up to 9) amino acid substitutions (such as conservative amino acid substitutions) compared to the amino acid sequence set forth as one of SEQ ID NOs: 2, 4, 6, 20, 24, or 29. 40

In some embodiments, the antibody or antigen binding fragment can include up to 10 (such as up to 1, up to 2, up to 3, up to 4, up to 5, up to 6, up to 7, up to 8, or up to 9) amino acid substitutions (such as conservative amino acid substitutions) in the framework regions of the heavy chain of 45 the antibody, or the light chain of the antibody, or the heavy and light chains of the antibody, compared to a known framework region, or compared to the framework regions of the EVB114, EVB100, EVB165, EVB166, or EVB167 antibody, and maintain the specific binding activity for 50 EBOV GP.

In certain embodiments, substitutions, insertions, or deletions may occur within one or more CDRs so long as such alterations do not substantially reduce the ability of the antibody to bind antigen. For example, conservative altera-55 tions (e.g., conservative substitutions as provided herein) that do not substantially reduce binding affinity may be made in CDRs. In certain embodiments of the variant  $V_H$  and  $V_L$ sequences provided above, each CDR either is unaltered, or contains no more than one, two or three amino acid substi-60 tutions.

To increase binding affinity of the antibody, the  $V_L$  and  $V_H$  segments can be randomly mutated, such as within HCDR3 region or the LCDR3 region, in a process analogous to the in vivo somatic mutation process responsible for affinity 65 maturation of antibodies during a natural immune response. Thus in vitro affinity maturation can be accomplished by

amplifying  $V_H$  and  $V_L$  regions using PCR primers complementary to the HCDR3 or LCDR3, respectively. In this process, the primers have been "spiked" with a random mixture of the four nucleotide bases at certain positions such that the resultant PCR products encode  $V_H$  and  $V_L$  segments into which random mutations have been introduced into the  $\mathrm{V}_{H}$  and/or  $\mathrm{V}_{L}$  CDR3 regions. These randomly mutated  $\mathrm{V}_{H}$ and  $V_L$  segments can be tested to determine the binding affinity for EBOV GP. In particular examples, the  $\mathrm{V}_{H}$  amino acid sequence is one of SEQ ID NOs: 1, 3, 5, 19, 23, or 28. In other examples, the  $V_L$  amino acid sequence is one of SEQ ID NOs: 2, 4, 6, 20, 24, or 29. Methods of in vitro affinity maturation are known (see, e.g., Chowdhury, Methods Mol. Biol. 207:179-196 (2008)), and Hoogenboom et al. in Methods in Molecular Biology 178:1-37 (O'Brien et al., ed., Human Press, Totowa, N.J., (2001).)

In certain embodiments, an antibody or antigen binding fragment is altered to increase or decrease the extent to which the antibody or antigen binding fragment is glycosylated. Addition or deletion of glycosylation sites may be conveniently accomplished by altering the amino acid sequence such that one or more glycosylation sites is created or removed.

Where the antibody comprises an Fc region, the carbohydrate attached thereto may be altered. Native antibodies produced by mammalian cells typically comprise a branched, biantennary oligosaccharide that is generally attached by an N-linkage to Asn297 of the CH<sub>2</sub> domain of the Fc region. See, e.g., Wright et al. *TIBTECH* 15:26-32 (1997). The oligosaccharide may include various carbohydrates, e.g., mannose, N-acetyl glucosamine (GlcNAc), galactose, and sialic acid, as well as a fucose attached to a GlcNAc in the "stem" of the biantennary oligosaccharide structure. In some embodiments, modifications of the oligosaccharide in an antibody may be made in order to create antibody variants with certain improved properties.

In one embodiment, antibody variants are provided having a carbohydrate structure that lacks fucose attached (directly or indirectly) to an Fc region. For example, the amount of fucose in such antibody may be from 1% to 80%, from 1% to 65%, from 5% to 65% or from 20% to 40%. The amount of fucose is determined by calculating the average amount of fucose within the sugar chain at Asn297, relative to the sum of all glycostructures attached to Asn 297 (e.g. complex, hybrid and high mannose structures) as measured by MALDI-TOF mass spectrometry, as described in WO 2008/077546, for example. Asn297 refers to the asparagine residue located at about position 297 in the Fc region; however, Asn297 may also be located about  $\pm 3$  amino acids upstream or downstream of position 297, i.e., between positions 294 and 300, due to minor sequence variations in antibodies. Such fucosylation variants may have improved ADCC function. See, e.g., US Patent Publication Nos. US 2003/0157108 (Presta, L.); US 2004/0093621 (Kyowa Hakko Kogyo Co., Ltd). Examples of publications related to "defucosylated" or "fucose-deficient" antibody variants include: US 2003/0157108; WO 2000/61739; WO 2001/ 29246; US 2003/0115614; US 2002/0164328; US 2004/ 0093621; US 2004/0132140; US 2004/0110704; US 2004/ 0110282; US 2004/0109865; WO 2003/085119; WO 2003/ 084570; WO 2005/035586; WO 2005/035778; WO2005/ 053742; WO2002/031140; Okazaki et al. J. Mol. Biol. 336:1239-1249 (2004); Yamane-Ohnuki et al. Biotech. Bioeng. 87: 614 (2004). Examples of cell lines capable of producing defucosylated antibodies include Lec 13 CHO cells deficient in protein fucosylation (Ripka et al. Arch. Biochem. Biophys. 249:533-545 (1986); US Pat Appl No US

.

2003/0157108 A1, Presta, L; and WO 2004/056312 A1, Adams et al., especially at Example 11), and knockout cell lines, such as alpha-1,6-fucosyltransferase gene, FUT8, knockout CHO cells (see, e.g., Yamane-Ohnuki et al. Biotech. Bioeng. 87: 614 (2004); Kanda, Y. et al., Biotechnol. 5 Bioeng., 94(4):680-688 (2006); and WO2003/085107).

Antibodies variants are further provided with bisected oligosaccharides, e.g., in which a biantennary oligosaccharide attached to the Fc region of the antibody is bisected by GlcNAc. Such antibody variants may have reduced fucosylation and/or improved ADCC function. Examples of such antibody variants are described, e.g., in WO 2003/011878 (Jean-Mairet et al.); U.S. Pat. No. 6,602,684 (Umana et al.); and US 2005/0123546 (Umana et al.). Antibody variants with at least one galactose residue in the oligosaccharide attached to the Fc region are also provided. Such antibody variants may have improved CDC function. Such antibody variants are described, e.g., in WO 1997/30087 (Patel et al.); WO 1998/58964 (Raju, S.); and WO 1999/22764 (Raju, S.). 20

In several embodiments, the constant region of the antibody includes one or more amino acid substitutions to optimize in vivo half-life of the antibody. The serum halflife of IgG Abs is regulated by the neonatal Fc receptor (FcRn). Thus, in several embodiments, the antibody 25 includes an amino acid substitution that increases binding to the FcRn. Several such substitutions are known, such as substitutions at IgG constant regions T250Q and M428L (see, e.g., Hinton et al., J Immunol., 176:346-356, 2006); M428L and N434S (the "LS" mutation, see, e.g., Zalevsky, 30 et al., Nature Biotechnology, 28:157-159, 2010); N434A (see, e.g., Petkova et al., Int. Immunol., 18:1759-1769, 2006); T307A, E380A, and N434A (see, e.g., Petkova et al., Int. Immunol., 18:1759-1769, 2006); and M252Y, S254T, and T256E (see, e.g., Dall'Acqua et al., J. Biol. Chem., 35 tized or linked to another molecule (such as another peptide 281:23514-23524, 2006). The disclosed antibodies and antigen binding fragments can be linked to a Fc polypeptide including any of the substitutions listed above, for example, the Fc polypeptide can include the M428L and N434S substitutions.

In some embodiments, the constant region of the antibody includes one of more amino acid substitutions to optimize antibody-dependent cell-mediated cytotoxicity (ADCC). ADCC is mediated primarily through a set of closely related Fcy receptors. In some embodiments, the antibody includes 45 one or more amino acid substitutions that increase binding to FcyRIIIa. Several such substitutions are known, such as substitutions at IgG constant regions S239D and I332E (see, e.g., Lazar et al., Proc. Natl., Acad. Sci. U.S.A., 103:4005-4010, 2006); and S239D, A330L, and I332E (see, e.g., Lazar 50 et al., Proc. Natl., Acad. Sci. U.S.A., 103:4005-4010, 2006).

Combinations of the above substitutions are also included, to generate an IgG constant region with increased binding to FcRn and FcyRIIIa. The combinations increase antibody half-life and ADCC. For example, such combina- 55 tion include antibodies with the following amino acid substitution in the Fc region:

(1) S239D/I332E and T250Q/M428L;

- (2) S239D/I332E and M428L/N434S;
- (3) S239D/I332E and N434A;
- (4) S239D/I332E and T307A/E380A/N434A;
- (5) S239D/I332E and M252Y/S254T/T256E;
- (6) S239D/A330L/I332E and T250Q/M428L;
- (7) S239D/A330L/I332E and M428L/N434S;
- (8) S239D/A330L/I332E and N434A;
- (9) S239D/A330L/I332E and T307A/E380A/N434A; or
- (10) S239D/A330L/I332E and M252Y/S254T/T256E.

In some examples, the antibodies, or an antigen binding fragment thereof is modified such that it is directly cytotoxic to infected cells, or uses natural defenses such as complement, antibody dependent cellular cytotoxicity (ADCC), or phagocytosis by macrophages.

In certain embodiments, an antibody provided herein may be further modified to contain additional nonproteinaceous moieties that are known in the art and readily available. The moieties suitable for derivatization of the antibody include but are not limited to water soluble polymers. Non-limiting examples of water soluble polymers include, but are not limited to, polyethylene glycol (PEG), copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1,3dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), and dextran or poly(n-vinyl pyrrolidone) polyethylene glycol, propropylene glycol homopolymers, prolypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols (e.g., glycerol), polyvinyl alcohol, and mixtures thereof. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its stability in water. The polymer may be of any molecular weight, and may be branched or unbranched. The number of polymers attached to the antibody may vary, and if more than one polymer are attached, they can be the same or different molecules. In general, the number and/or type of polymers used for derivatization can be determined based on considerations including, but not limited to, the particular properties or functions of the antibody to be improved, whether the antibody derivative will be used in a therapy under defined conditions, etc.

The antibody or antigen binding fragment can be derivaor protein). In general, the antibody or antigen binding fragment is derivatized such that the binding to EBOV GP is not affected adversely by the derivatization or labeling. For example, the antibody or antigen binding fragment can be functionally linked (by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as another antibody (for example, a bi-specific antibody or a diabody), a detectable marker, an effector molecule, or a protein or peptide that can mediate association of the antibody or antibody portion with another molecule (such as a streptavidin core region or a polyhistidine tag).

B. Conjugates

40

65

The monoclonal antibodies and antigen binding fragments that specifically bind to an epitope on EBOV GP can be conjugated to an agent, such as an effector molecule or detectable marker, using any number of means known to those of skill in the art. Both covalent and noncovalent attachment means may be used. One of skill in the art will appreciate that various effector molecules and detectable markers can be used, including (but not limited to) toxins and radioactive agents such as <sup>125</sup>I, <sup>32</sup>P, <sup>14</sup>C, <sup>3</sup>H and <sup>35</sup>S and other labels, target moieties and ligands, etc. The choice of a particular effector molecule or detectable marker depends 60 on the particular target molecule or cell, and the desired biological effect.

The choice of a particular effector molecule or detectable marker depends on the particular target molecule or cell, and the desired biological effect. Thus, for example, the effector molecule can be a cytotoxin that is used to bring about the death of a particular target cell (such as an EBOV infected cell). In other embodiments, the effector molecule can be a

cytokine, such as IL-15; conjugates including the cytokine can be used, e.g., to stimulate immune cells locally.

The procedure for attaching an effector molecule or detectable marker to an antibody or antigen binding fragment varies according to the chemical structure of the 5 effector. Polypeptides typically contain a variety of functional groups; such as carboxylic acid (COOH), free amine (-NH<sub>2</sub>) or sulfhydryl (-SH) groups, which are available for reaction with a suitable functional group on a polypeptide to result in the binding of the effector molecule or 10 detectable marker. Alternatively, the antibody or antigen binding fragment is derivatized to expose or attach additional reactive functional groups. The derivatization may involve attachment of any of a number of known linker molecules such as those available from Pierce Chemical 15 Company, Rockford, Ill. The linker can be any molecule used to join the antibody or antigen binding fragment to the effector molecule or detectable marker. The linker is capable of forming covalent bonds to both the antibody or antigen binding fragment and to the effector molecule or detectable 20 marker. Suitable linkers are well known to those of skill in the art and include, but are not limited to, straight or branched-chain carbon linkers, heterocyclic carbon linkers, or peptide linkers. Where the antibody or antigen binding fragment and the effector molecule or detectable marker are 25 polypeptides, the linkers may be joined to the constituent amino acids through their side groups (such as through a disulfide linkage to cysteine) or to the alpha carbon amino and carboxyl groups of the terminal amino acids.

In view of the large number of methods that have been 30 reported for attaching a variety of radiodiagnostic compounds, radiotherapeutic compounds, labels (such as enzymes or fluorescent molecules), toxins, and other agents to antibodies one skilled in the art will be able to determine a suitable method for attaching a given agent to an antibody 35 or antigen binding fragment or other polypeptide. For example, the antibody or antigen binding fragment can be conjugated with effector molecules such as small molecular weight drugs such as Monomethyl Auristatin E (MMAE), Monomethyl Auristatin F (MMAF), maytansine, maytansine 40 derivatives, including the derivative of maytansine known as DM1 (also known as mertansine), or other agents to make an antibody drug conjugate (ADC). In several embodiments, conjugates of an antibody or antigen binding fragment and one or more small molecule toxins, such as a calicheamicin, 45 maytansinoids, dolastatins, auristatins, a trichothecene, and CCl065, and the derivatives of these toxins that have toxin activity, are provided.

The antibody or antigen binding fragment can be conjugated with a detectable marker; for example, a detectable 50 marker capable of detection by ELISA, spectrophotometry, flow cytometry, microscopy or diagnostic imaging techniques (such as computed tomography (CT), computed axial tomography (CAT) scans, magnetic resonance imaging (MRI), nuclear magnetic resonance imaging NMRI), mag- 55 netic resonance tomography (MTR), ultrasound, fiberoptic examination, and laparoscopic examination). Specific, nonlimiting examples of detectable markers include fluorophores, chemiluminescent agents, enzymatic linkages, radioactive isotopes and heavy metals or compounds (for 60 example super paramagnetic iron oxide nanocrystals for detection by MRI). For example, useful detectable markers include fluorescent compounds, including fluorescein, fluorescein isothiocyanate, rhodamine, 5-dimethylamine-1napthalenesulfonyl chloride, phycoerythrin, lanthanide 65 phosphors and the like. Bioluminescent markers are also of use, such as luciferase, Green fluorescent protein (GFP),

54

Yellow fluorescent protein (YFP). An antibody or antigen binding fragment can also be conjugated with enzymes that are useful for detection, such as horseradish peroxidase, β-galactosidase, luciferase, alkaline phosphatase, glucose oxidase and the like. When an antibody or antigen binding fragment is conjugated with a detectable enzyme, it can be detected by adding additional reagents that the enzyme uses to produce a reaction product that can be discerned. For example, when the agent horseradish peroxidase is present the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is visually detectable. An antibody or antigen binding fragment may also be conjugated with biotin, and detected through indirect measurement of avidin or streptavidin binding. It should be noted that the avidin itself can be conjugated with an enzyme or a fluorescent label.

The antibody or antigen binding fragment can be conjugated with a paramagnetic agent, such as gadolinium. Paramagnetic agents such as superparamagnetic iron oxide are also of use as labels. Antibodies can also be conjugated with lanthanides (such as europium and dysprosium), and manganese. An antibody or antigen binding fragment may also be labeled with a predetermined polypeptide epitopes recognized by a secondary reporter (such as leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags).

The antibody or antigen binding fragment can also be conjugated with a radiolabeled amino acid. The radiolabel may be used for both diagnostic and therapeutic purposes. For instance, the radiolabel may be used to detect EBOV GP and EBOV GP expressing cells by x-ray, emission spectra, or other diagnostic techniques. Examples of labels for polypeptides include, but are not limited to, the following radioisotopes or radionucleotides: <sup>3</sup>H, <sup>14</sup>C, <sup>15</sup>N, <sup>35</sup>S, <sup>90</sup>Y, <sup>99</sup>Te, <sup>111</sup>In, <sup>125</sup>I, <sup>131</sup>I.

Means of detecting such detectable markers are well known to those of skill in the art. Thus, for example, radiolabels may be detected using photographic film or scintillation counters, fluorescent markers may be detected using a photodetector to detect emitted illumination. Enzymatic labels are typically detected by providing the enzyme with a substrate and detecting the reaction product produced by the action of the enzyme on the substrate, and colorimetric labels are detected by simply visualizing the colored label.

The average number of effector molecule or detectable marker moieties per antibody or antigen binding fragment in a conjugate can range, for example, from 1 to 20 moieties per antibody or antigen binding fragment. In certain embodiments, the average number of effector molecule or detectable marker moieties per antibody or antigen binding fragment in a conjugate range from about 1 to about 2, from about 1 to about 3, about 1 to about 8; from about 2 to about 6; from about 3 to about 5; or from about 3 to about 4. The loading (for example, effector molecule/antibody ratio) of an conjugate may be controlled in different ways, for example, by: (i) limiting the molar excess of effector molecule-linker intermediate or linker reagent relative to antibody, (ii) limiting the conjugation reaction time or temperature, (iii) partial or limiting reductive conditions for cysteine thiol modification, (iv) engineering by recombinant techniques the amino acid sequence of the antibody such that the number and position of cysteine residues is modified for control of the number or position of linker-effector molecule attachments.

C. Polynucleotides and Expression

Nucleic acids molecules (for example, cDNA molecules) encoding the amino acid sequences of antibodies, antigen binding fragments, and conjugates that specifically bind EBOV GP are provided. Nucleic acids encoding these 5 molecules can readily be produced by one of skill in the art, using the amino acid sequences provided herein (such as the CDR sequences and  $V_H$  and  $V_L$  sequences), sequences available in the art (such as framework or constant region sequences), and the genetic code. In several embodiments, a 10 nucleic acid molecules can encode the  $V_H$ , the  $V_L$ , or both the  $V_H$  and  $V_L$  (for example in a bicistronic expression vector) of a disclosed antibody or antigen binding fragment. In several embodiments, the nucleic acid molecules can be expressed in a host cell (such as a mammalian cell) to 15 produce a disclosed antibody or antigen binding fragment.

One of skill in the art can readily use the genetic code to construct a variety of functionally equivalent nucleic acids, such as nucleic acids which differ in sequence but which encode the same antibody sequence, or encode a conjugate 20 or fusion protein including the  $V_L$  and/or  $V_H$  nucleic acid sequence.

In a non-limiting example, an isolated nucleic acid molecule encodes the  $V_H$  of a disclosed antibody or antigen binding fragment and includes the nucleic acid sequence set 25 forth as any one of SEQ ID NOs: 7, 9, 11, 21, 25, or 30. In a non-limiting example, an isolated nucleic acid molecule encodes the  $V_L$  of a disclosed antibody or antigen binding fragment and includes the nucleic acid sequence set forth as any one of SEQ ID NOs: 8, 10, 12, 22, 26, 27, or 31. In a 30 non-limiting example, an isolated nucleic acid molecule encodes the  $V_H$  and  $V_L$  of a disclosed antibody or antigen binding fragment and includes the nucleic acid sequences set forth as any one of SEQ ID NOs: 7 and 8, respectively, 9 and 11, respectively, 11 and 12, respectively, 21 and 22, respec-35 tively, 25 and 26, respectively, 25 and 27, respectively, or 30 and 31, respectively.

Nucleic acid sequences encoding the of antibodies, antigen binding fragments, and conjugates that specifically bind EBOV GP can be prepared by any suitable method includ- 40 ing, for example, cloning of appropriate sequences or by direct chemical synthesis by methods such as the phosphotriester method of Narang et al., Meth. Enzymol. 68:90-99, 1979; the phosphodiester method of Brown et al., Meth. Enzymol. 68:109-151, 1979; the diethylphosphoramidite 45 method of Beaucage et al., Tetra. Lett. 22:1859-1862, 1981; the solid phase phosphoramidite triester method described by Beaucage & Caruthers, Tetra. Letts. 22(20):1859-1862, 1981, for example, using an automated synthesizer as described in, for example, Needham-VanDevanter et al., 50 Fc. Nucl. Acids Res. 12:6159-6168, 1984; and, the solid support method of U.S. Pat. No. 4,458,066. Chemical synthesis produces a single stranded oligonucleotide. This can be converted into double stranded DNA by hybridization with a complementary sequence or by polymerization with a 55 DNA polymerase using the single strand as a template.

Exemplary nucleic acids can be prepared by cloning techniques. Examples of appropriate cloning and sequencing techniques, and instructions sufficient to direct persons of skill through many cloning exercises are known (see, e.g., 60 Sambrook et al. (Molecular Cloning: A Laboratory Manual, 4<sup>th</sup> ed, Cold Spring Harbor, N.Y., 2012) and Ausubel et al. (In Current Protocols in Molecular Biology, John Wiley & Sons, New York, through supplement 104, 2013). Product information from manufacturers of biological reagents and 65 experimental equipment also provide useful information. Such manufacturers include the SIGMA Chemical Company

(Saint Louis, Mo.), R&D Systems (Minneapolis, Minn.), Pharmacia Amersham (Piscataway, N.J.), CLONTECH Laboratories, Inc. (Palo Alto, Calif.), Chem Genes Corp., Aldrich Chemical Company (Milwaukee, Wis.), Glen Research, Inc., GIBCO BRL Life Technologies, Inc. (Gaithersburg, Md.), Fluka Chemica-Biochemika Analytika (Fluka Chemie AG, Buchs, Switzerland), Invitrogen (Carlsbad, Calif.), and Applied Biosystems (Foster City, Calif.), as well as many other commercial sources known to one of skill.

Nucleic acids can also be prepared by amplification methods. Amplification methods include polymerase chain reaction (PCR), the ligase chain reaction (LCR), the transcription-based amplification system (TAS), the self-sustained sequence replication system (3SR). A wide variety of cloning methods, host cells, and in vitro amplification methodologies are well known to persons of skill.

The nucleic acid molecules can be expressed in a recombinantly engineered cell such as bacteria, plant, yeast, insect and mammalian cells. The antibodies, antigen binding fragments, and conjugates can be expressed as individual  $V_H$  and/or  $V_L$  chain (linked to an effector molecule or detectable marker as needed), or can be expressed as a fusion protein. Methods of expressing and purifying antibodies and antigen binding fragments are known and further described herein (see, e.g., Al-Rubeai (ed), Antibody Expression and Production, Springer Press, 2011). An immunoadhesin can also be expressed. Thus, in some examples, nucleic acids encoding a  $V_H$  and  $V_L$ , and immunoadhesin are provided. The nucleic acid sequences can optionally encode a leader sequence.

To create a scFv the  $V_{H^-}$  and  $V_L$ -encoding DNA fragments can be operatively linked to another fragment encoding a flexible linker, e.g., encoding the amino acid sequence (Gly<sub>4</sub>-Ser)<sub>3</sub>, such that the  $V_H$  and  $V_L$  sequences can be expressed as a contiguous single-chain protein, with the  $V_L$ and  $V_H$  domains joined by the flexible linker (see, e.g., Bird et al., *Science* 242:423-426, 1988; Huston et al., *Proc. Natl. Acad. Sci. USA* 85:5879-5883, 1988; McCafferty et al., *Nature* 348:552-554, 1990; Kontermann and Dubel (Ed), Antibody Engineering, Vols. 1-2,  $2^{nd}$  Ed., Springer Press, 2010; Harlow and Lane, *Antibodies: A Laboratory Manual*,  $2^{nd}$ , Cold Spring Harbor Laboratory, New York, 2013,). Optionally, a cleavage site can be included in a linker, such as a furin cleavage site.

The nucleic acid encoding a  $V_H$  and/or the  $V_L$  optionally can encode an Fc domain (immunoadhesin). The Fc domain can be an IgA, IgM or IgG Fc domain. The Fc domain can be an optimized Fc domain, as described in U.S. Published Patent Application No. 20100/093979, incorporated herein by reference. In one example, the immunoadhesin is an IgG<sub>1</sub> Fc.

The single chain antibody may be monovalent, if only a single  $V_H$  and  $V_L$  are used, bivalent, if two  $V_H$  and  $V_L$  are used, or polyvalent, if more than two  $V_H$  and  $V_L$  are used. Bispecific or polyvalent antibodies may be generated that bind specifically to EBOV GP and another antigen. The encoded  $V_H$  and  $V_L$  optionally can include a furin cleavage site between the  $V_H$  and  $V_L$  domains.

Those of skill in the art are knowledgeable in the numerous expression systems available for expression of proteins including *E. coli*, other bacterial hosts, yeast, and various higher eukaryotic cells such as the COS, CHO, HeLa and myeloma cell lines.

One or more DNA sequences encoding the antibodies, antigen binding fragments, or conjugates can be expressed in vitro by DNA transfer into a suitable host cell. The cell may be prokaryotic or eukaryotic. The term also includes any progeny of the subject host cell. It is understood that all progeny may not be identical to the parental cell since there may be mutations that occur during replication. Methods of stable transfer, meaning that the foreign DNA is continuously maintained in the host, are known in the art. Hybridomas expressing the antibodies of interest are also encompassed by this disclosure.

The expression of nucleic acids encoding the antibodies and antigen binding fragments described herein can be achieved by operably linking the DNA or cDNA to a promoter (which is either constitutive or inducible), fol- 10 lowed by incorporation into an expression cassette. The promoter can be any promoter of interest, including a cytomegalovirus promoter and a human T cell lymphotrophic virus promoter (HTLV)-1. Optionally, an enhancer, such as a cytomegalovirus enhancer, is included in the 15 construct. The cassettes can be suitable for replication and integration in either prokaryotes or eukaryotes. Typical expression cassettes contain specific sequences useful for regulation of the expression of the DNA encoding the protein. For example, the expression cassettes can include 20 appropriate promoters, enhancers, transcription and translation terminators, initiation sequences, a start codon (i.e., ATG) in front of a protein-encoding gene, splicing signal for introns, sequences for the maintenance of the correct reading frame of that gene to permit proper translation of mRNA, 25 and stop codons. The vector can encode a selectable marker, such as a marker encoding drug resistance (for example, ampicillin or tetracycline resistance).

To obtain high level expression of a cloned gene, it is desirable to construct expression cassettes which contain, at 30 the minimum, a strong promoter to direct transcription, a ribosome binding site for translational initiation (internal ribosomal binding sequences), and a transcription/translation terminator. For E. coli, this includes a promoter such as the T7, trp, lac, or lambda promoters, a ribosome binding 35 site, and preferably a transcription termination signal. For eukaryotic cells, the control sequences can include a promoter and/or an enhancer derived from, for example, an immunoglobulin gene, HTLV, SV40 or cytomegalovirus, and a polyadenylation sequence, and can further include 40 splice donor and/or acceptor sequences (for example, CMV and/or HTLV splice acceptor and donor sequences). The cassettes can be transferred into the chosen host cell by well-known methods such as transformation or electroporation for E. coli and calcium phosphate treatment, elec- 45 troporation or lipofection for mammalian cells. Cells transformed by the cassettes can be selected by resistance to antibiotics conferred by genes contained in the cassettes, such as the amp, gpt, neo and hyg genes.

When the host is a eukaryote, such methods of transfec- 50 tion of DNA as calcium phosphate coprecipitates, conventional mechanical procedures such as microinjection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors may be used. Eukaryotic cells can also be cotransformed with polynucleotide sequences encoding the 55 antibody, labeled antibody, or antigen biding fragment, and a second foreign DNA molecule encoding a selectable phenotype, such as the herpes simplex thymidine kinase gene. Another method is to use a eukaryotic viral vector, such as simian virus 40 (SV40) or bovine papilloma virus, 60 to transiently infect or transform eukaryotic cells and express the protein (see for example, Viral Expression Vectors, Springer press, Muzyczka ed., 2011). One of skill in the art can readily use an expression systems such as plasmids and vectors of use in producing proteins in cells 65 including higher eukaryotic cells such as the COS, CHO, HeLa and myeloma cell lines.

Also provided is a population of cells comprising at least one host cell described herein. The population of cells can be a heterogeneous population comprising the host cell comprising any of the recombinant expression vectors described, in addition to at least one other cell, e.g., a host cell (e.g., a T cell), which does not comprise any of the recombinant expression vectors, or a cell other than a T cell, e.g., a B cell, a macrophage, a neutrophil, an erythrocyte, a hepatocyte, an endothelial cell, an epithelial cell, a muscle cell, a brain cell, etc. Alternatively, the population of cells can be a substantially homogeneous population, in which the population comprises mainly host cells (e.g., consisting essentially of) comprising the recombinant expression vector. The population also can be a clonal population of cells, in which all cells of the population are clones of a single host cell comprising a recombinant expression vector, such that all cells of the population comprise the recombinant expression vector. In one embodiment, the population of cells is a clonal population comprising host cells comprising a recombinant expression vector as described herein

Modifications can be made to a nucleic acid encoding a polypeptide described herein without diminishing its biological activity. Some modifications can be made to facilitate the cloning, expression, or incorporation of the targeting molecule into a fusion protein. Such modifications are well known to those of skill in the art and include, for example, termination codons, a methionine added at the amino terminus to provide an initiation, site, additional amino acids placed on either terminus to create conveniently located restriction sites, or additional amino acids (such as poly His) to aid in purification steps. In addition to recombinant methods, the immunoconjugates, effector moieties, and antibodies of the present disclosure can also be constructed in whole or in part using standard peptide synthesis well known in the art.

Once expressed, the antibodies, antigen binding fragments, and conjugates can be purified according to standard procedures in the art, including ammonium sulfate precipitation, affinity columns, column chromatography, and the like (see, generally, Simpson ed., Basic methods in Protein Purification and Analysis: A laboratory Manual, Cold Harbor Press, 2008). The antibodies, antigen binding fragment, and conjugates need not be 100% pure. Once purified, partially or to homogeneity as desired, if to be used therapeutically, the polypeptides should be substantially free of endotoxin.

Methods for expression of the antibodies, antigen binding fragments, and conjugates, and/or refolding to an appropriate active form, from mammalian cells, and bacteria such as *E. coli* have been described and are well-known and are applicable to the antibodies disclosed herein. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, 2<sup>nd</sup>, Cold Spring Harbor Laboratory, New York, 2013, Simpson ed., Basic methods in Protein Purification and Analysis: A laboratory Manual, Cold Harbor Press, 2008, and Ward et al., *Nature* 341:544, 1989.

Often, functional heterologous proteins from *E. coli* or other bacteria are isolated from inclusion bodies and require solubilization using strong denaturants, and subsequent refolding. During the solubilization step, as is well known in the art, a reducing agent must be present to separate disulfide bonds. An exemplary buffer with a reducing agent is: 0.1 M Tris pH 8, 6 M guanidine, 2 mM EDTA, 0.3 M DTE (dithioerythritol). Reoxidation of the disulfide bonds can occur in the presence of low molecular weight thiol reagents in reduced and oxidized form, as described in Saxena et al., *Biochemistry* 9: 5015-5021, 1970,

In addition to recombinant methods, the antibodies, antigen binding fragments, and/or conjugates can also be constructed in whole or in part using standard peptide synthesis. Solid phase synthesis of the polypeptides can be accomplished by attaching the C-terminal amino acid of the sequence to an insoluble support followed by sequential addition of the remaining amino acids in the sequence. Techniques for solid phase synthesis are described by Barany & Merrifield, The Peptides: Analysis, Synthesis, Biology. Vol. 2: Special Methods in Peptide Synthesis, Part A. pp. 3-284; Merrifield et al., J. Am. Chem. Soc. 85:2149-2156, 1963, and Stewart et al., Solid Phase Peptide Synthesis, 2nd ed., Pierce Chem. Co., Rockford, Ill., 1984. Proteins of greater length may be synthesized by condensation of the amino and carboxyl termini of shorter fragments. Methods of forming peptide bonds by activation of a carboxyl terminal end (such as by the use of the coupling reagent N, N'-dicylohexylcarbodimide) are well known in the art.

D. Methods and Composition

1. Methods of Inhibiting, Treating, and Preventing EBOV Infection and Disease

Methods are disclosed herein for the prevention or treatment of an EBOV infection or EVD, such as a ZEBOV infection, in a subject. Prevention can include inhibition of 25 infection with EBOV. The method can include administering to a subject a therapeutically effective amount of a disclosed antibody, antigen binding fragment, or conjugate that specifically binds EBOV GP, or a nucleic acid encoding such an antibody, antigen binding fragment, conjugate. In some examples, the antibody, antigen binding fragment, conjugate, or nucleic acid molecule, can be used pre-exposure (for example, to prevent or inhibit EBOV infection). In some examples, the antibody, antigen binding fragment, conju-35 gate, or nucleic acid molecule, can be used in post-exposure prophylaxis. In some examples, the antibody, antigen binding fragment, conjugate, or nucleic acid molecule, can be used to eliminate or reduce the viral load of EBOV in a subject infected with EBOV. For example a therapeutically 40 effective amount of an antibody, antigen binding fragment, conjugate, or nucleic acid molecule, can be administered to a subject with an EBOV infection. In some examples the antibody, antigen binding fragment, conjugate, or nucleic acid molecule is modified such that it is directly cytotoxic to 45 infected cells (e.g., by conjugation to a toxin), or uses natural defenses such as complement, antibody dependent cellular cytotoxicity (ADCC), or phagocytosis by macrophages, or can be modified to increase the natural defenses.

The EVD or EBOV infection in the subject does not need 50 to be completely eliminated for the method to be effective. For example, the method can reduce or ameliorate EVD or EBOV infection by a desired amount, for example by at least 10%, at least 20%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 98%, or even 55 at least 100% (elimination of detectable EBOV infection or EVD), as compared to EBOV infection or EVD in the absence of the treatment.

In one non-limiting example, the method reduces viral titer in a subject with an EBOV infection. For example, 60 administration of a therapeutically effective amount of a disclosed EBOV GP-specific antibody or antigen binding fragment or conjugate can reduce viral titer by at least 10%, at least 20%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 98%, or even at least 65 100% (elimination of detectable EBOV) in the subject. Methods of determining the EBOV viral titer in the subject

are known, and include, for example, obtaining a blood sample from the subject and assaying the sample for EBOV activity.

In several embodiments, administration of a therapeutically effective amount of a disclosed antibody, antigen binding fragment, conjugate, or nucleic acid molecule, results in a reduction in the establishment of EBOV infection and/or reducing subsequent EVD progression in a subject. A reduction in the establishment of EBOV infection and/or a reduction in subsequent EVD progression encompass any statistically significant reduction in EBOV activity.

In several embodiments, the subject can be selected for treatment, for example, a subject at risk of EBOV infection, or known to have an EBOV infection. In some embodiments, a subject can be selected that is at risk of or known to have an infection with a particular strain of EBOV, such as BDBV, RESTV, SUDV, TAFV, or ZEBOV.

In several embodiments, a method of preventing or inhibiting EBOV infection (e.g., ZEBOV infection) of a cell is 20 provided. The method includes contacting the cell with an effective amount of an antibody or antigen binding fragment as disclosed herein. For example the cell can be incubated with the effective amount of the antibody or antigen binding fragment prior to or contemporaneous with incubation with the EBOV. EBOV infection of the cell does not need to be completely eliminated for the method to be effective. For example, a method can reduce EBOV infection by a desired amount, for example by at least 10%, at least 20%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 98%, or even at least 100% (elimination of detectable EBOV infected cells), as compared to EBOV infection in the absence of the treatment. In some embodiments, the cell is also contacted with an effective amount of an additional agent, such as anti-viral agent. The cell can be in vivo or in vitro.

Studies in have shown that cocktails of EBOV neutralizing antibodies that target different epitopes of EBOV GP can treat macaques infected with ZEBOV (Qiu et al., *Sci. Transl. Med.*, 4, 138ra81, 2012). Accordingly, in some examples, a subject is further administered one or more additional antibodies that bind EBOV GP and that can neutralize EBOV infection. For example, the subject can be administered a therapeutically effective amount of a set of antibodies including two or more of the EVB100, EVB114 and EVB166 antibodies disclosed herein. The antibodies can be administered as a cocktail (that is, as a single composition including the two or more antibodies), or can be administered in sequence.

In some examples, a subject is administered the DNA encoding the antibody or antigen binding fragments thereof, to provide in vivo antibody production, for example using the cellular machinery of the subject. Immunization by nucleic acid constructs is well known in the art and taught, for example, in U.S. Pat. No. 5,643,578, and U.S. Pat. No. 5,593,972 and U.S. Pat. No. 5,817,637. U.S. Pat. No. 5,880,103 describes several methods of delivery of nucleic acids encoding to an organism. One approach to administration of nucleic acids is direct administration with plasmid DNA, such as with a mammalian expression plasmid. The nucleotide sequence encoding the disclosed antibody, or antigen binding fragments thereof, can be placed under the control of a promoter to increase expression. The methods include liposomal delivery of the nucleic acids. Such methods can be applied to the production of an antibody, or antigen binding fragments thereof. In some embodiments, a disclosed antibody or antigen binding fragment is expressed in a subject using the pVRC8400 vector (described in

Barouch et al., J. Virol, 79, 8828-8834, 2005, which is incorporated by reference herein).

The nucleic acid molecules encoding the disclosed antibodies or antigen binding fragments can be included in a viral vector, for example for expression of the antibody or 5 antigen binding fragment in a host cell, or a subject (such as a subject with or at risk of EBOV infection). A number of viral vectors have been constructed, that can be used to express the disclosed antibodies or antigen binding fragments, such as a retroviral vector, an adenoviral vector, or an 10 adeno-associated virus (AAV) vector. In several examples, the viral vector can be replication-competent. For example, the viral vector can have a mutation in the viral genome that does not inhibit viral replication in host cells. The viral vector also can be conditionally replication-competent. In 15 other examples, the viral vector is replication-deficient in host cells.

In several embodiments, a subject (such as a human subject with or at risk of HIV-1 infection) can be administered a therapeutically effective amount of an adeno-asso- 20 ciated virus (AAV) viral vector that includes one or more nucleic acid molecules encoding a disclosed antibody or antigen binding fragment. The AAV viral vector is designed for expression of the nucleic acid molecules encoding a disclosed antibody or antigen binding fragment, and admin- 25 istration of the therapeutically effective amount of the AAV viral vector to the subject leads to expression of a therapeutically effective amount of the antibody or antigen binding fragment in the subject. Non-limiting examples of AAV viral vectors that can be used to express a disclosed antibody or 30 antigen binding fragment in a subject include those provided in Johnson et al ("Vector-mediated gene transfer engenders long-lived neutralizing activity and protection against SIV infection in monkeys," Nat. Med., 15(8):901-906, 2009) and Gardner et al. ("AAV-expressed eCD4-Ig provides durable 35 protection from multiple SHIV challenges," Nature, 519 (7541): 87-91, 2015), each of which is incorporated by reference herein in its entirety.

In one embodiment, a nucleic acid encoding a disclosed antibody, or antigen binding fragments thereof, is introduced 40 directly into cells. For example, the nucleic acid can be loaded onto gold microspheres by standard methods and introduced into the skin by a device such as Bio-Rad's HELIOS<sup>TM</sup> Gene Gun. The nucleic acids can be "naked," consisting of plasmids under control of a strong promoter. 45

Typically, the DNA is injected into muscle, although it can also be injected directly into other sites. Dosages for injection are usually around 0.5  $\mu$ g/kg to about 50 mg/kg, and typically are about 0.005 mg/kg to about 5 mg/kg (see, e.g., U.S. Pat. No. 5,589,466).

2. Dosages

A therapeutically effective amount of an EBOV GPspecific antibody, antigen binding fragment, conjugate, or nucleic acid molecule encoding such molecules, will depend upon the severity of the disease and/or infection and the 55 general state of the patient's health. A therapeutically effective amount is that which provides either subjective relief of a symptom(s) or an objectively identifiable improvement as noted by the clinician or other qualified observer. The EBOV GP-specific antibody, antigen binding fragment, conjugate, 60 or nucleic acid molecule encoding such molecules, can be administered in conjunction with another therapeutic agent, either simultaneously or sequentially.

Single or multiple administrations of a composition including a disclosed EBOV GP-specific antibody, antigen 65 binding fragment, conjugate, or nucleic acid molecule encoding such molecules, can be administered depending on

the dosage and frequency as required and tolerated by the patient. Compositions including the EBOV GP-specific antibody, antigen binding fragment, conjugate, or nucleic acid molecule encoding such molecules, should provide a sufficient quantity of at least one of the EBOV GP-specific antibodies, antigen binding fragments, conjugates, or nucleic acid molecules to effectively treat the patient. The dosage can be administered once, but may be applied periodically until either a therapeutic result is achieved or until side effects warrant discontinuation of therapy. In one example, a dose of the antibody or antigen binding fragment is infused for thirty minutes every other day. In this example, about one to about ten doses can be administered, such as three or six doses can be administered every other day. In a further example, a continuous infusion is administered for about five to about ten days. The subject can be treated at regular intervals, such as daily, weekly, or monthly, until a desired therapeutic result is achieved. Generally, the dose is sufficient to treat or ameliorate symptoms or signs of disease without producing unacceptable toxicity to the patient.

Data obtained from cell culture assays and animal studies can be used to formulate a range of dosage for use in humans. The dosage normally lies within a range of circulating concentrations that include the  $ED_{50}$ , with little or minimal toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. The therapeutically effective dose can be determined from cell culture assays and animal studies.

In certain embodiments, the antibody or antigen binding fragment that specifically binds EBOV GP, or conjugate thereof, or a nucleic acid molecule or vector encoding such a molecule, can be administered at a dose in the range of from about 1 to about 100 mg/kg, such as about 5-50 mg/kg, about 25-75 mg/kg, or about 40-60 mg/kg. In some embodiments, the dosage can be administered at about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, or 300 mg/kg, or other dose deemed appropriate by the treating physician. Further, the doses described herein can be administered according to the dosing frequency or frequency of administration described herein, including without limitation daily, every other day, 2 or 3 times per week, weekly, every 2 weeks, every 3 weeks, monthly, etc. In some embodiments, the dosage is administered daily beginning at the time of diagnosis with EBOV and until EBOV symptoms are alleviated. Additional treatments, including additional courses of therapy with a disclosed agent can be performed as needed.

## 50 3. Modes of Administration

The EBOV GP-specific antibody, antigen binding fragment, conjugate, nucleic acid molecule, or composition, as well as additional agents, can be administered to subjects in various ways, including local and systemic administration, such as, e.g., by injection subcutaneously, intravenously, intra-arterially, intraperitoneally, intramuscularly, intradermally, or intrathecally. In an embodiment, a therapeutic agent is administered by a single subcutaneous, intravenous, intra-arterial, intraperitoneal, intramuscular, intradermal or intrathecal injection once a day. The therapeutic agent can also be administered by direct injection at or near the site of disease.

The therapeutic agent may also be administered orally in the form of microspheres, microcapsules, liposomes (uncharged or charged (e.g., cationic)), polymeric microparticles (e.g., polyamides, polylactide, polyglycolide, poly (lactide-glycolide)), microemulsions, and the like. A further method of administration is by osmotic pump (e.g., an Alzet pump) or mini-pump (e.g., an Alzet miniosmotic pump), which allows for controlled, continuous and/or slow-release delivery of the therapeutic agent or pharmaceutical composition over a pre-determined period. 5 The osmotic pump or mini-pump can be implanted subcutaneously, or near a target site.

It will be apparent to one skilled in the art that the therapeutic agent or compositions thereof can also be administered by other modes. The therapeutic agent can be admin- 10 istered as pharmaceutical formulations suitable for, e.g., oral (including buccal and sub-lingual), rectal, nasal, topical, pulmonary, vaginal or parenteral (including intramuscular, intraarterial, intrathecal, subcutaneous and intravenous) administration, or in a form suitable for administration by 15 inhalation or insufflation. Depending on the intended mode of administration, the pharmaceutical formulations can be in the form of solid, semi-solid or liquid dosage forms, such as tablets, suppositories, pills, capsules, powders, liquids, suspensions, emulsions, creams, ointments, lotions, and the 20 like. The formulations can be provided in unit dosage form suitable for single administration of a precise dosage. The formulations comprise an effective amount of a therapeutic agent, and one or more pharmaceutically acceptable excipients, carriers and/or diluents, and optionally one or more 25 other biologically active agents.

4. Composition

Compositions are provided that include one or more of the disclosed EBOV GP-specific antibodies, antigen binding fragments, conjugates, or nucleic acid molecules, in a car- 30 rier. The compositions are useful, for example, for the treatment or detection of an EBOV infection. The compositions can be prepared in unit dosage forms for administration to a subject. The amount and timing of administration are at the discretion of the treating physician to achieve the 35 desired purposes. The EBOV GP-specific antibody, antigen binding fragment, conjugate, or nucleic acid molecule encoding such molecules can be formulated for systemic or local administration. In one example, the EBOV GP-specific antibody, antigen binding fragment, conjugate, or nucleic 40 acid molecule encoding such molecules, is formulated for parenteral administration, such as intravenous administration.

In some embodiments, the compositions comprise an antibody, antigen binding fragment, or conjugate thereof, in 45 at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% purity. In certain embodiments, the compositions contain less than 10%, less than 5%, less than 4%, less than 3%, less than 2%, less than 1% or less than 0.5% 50 of macromolecular contaminants, such as other mammalian (e.g., human) proteins.

The compositions for administration can include a solution of the EBOV GP-specific antibody, antigen binding fragment, conjugate, or nucleic acid molecule encoding such 55 molecules, dissolved in a pharmaceutically acceptable carrier, such as an aqueous carrier. A variety of aqueous carriers can be used, for example, buffered saline and the like. These solutions are sterile and generally free of undesirable matter. These compositions may be sterilized by conventional, well 60 known sterilization techniques. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents and the like, for example, sodium acetate, sodium chloride, 65 potassium chloride, calcium chloride, sodium lactate and the like. The concentration of antibody in these formulations can

vary widely, and will be selected primarily based on fluid volumes, viscosities, body weight and the like in accordance with the particular mode of administration selected and the subject's needs.

A typical composition for intravenous administration includes about 0.01 to about 30 mg/kg of antibody or antigen binding fragment or conjugate per subject per day (or the corresponding dose of a conjugate including the antibody or antigen binding fragment). Actual methods for preparing administrable compositions will be known or apparent to those skilled in the art and are described in more detail in such publications as Remington's Pharmaceutical Science, 22th ed., Pharmaceutical Press, London, UK (2012). In some embodiments, the composition can be a liquid formulation including one or more antibodies, antigen binding fragments (such as an antibody or antigen binding fragment that specifically binds to EBOV GP), in a concentration range from about 0.1 mg/ml to about 20 mg/ml, or from about 0.5 mg/ml to about 20 mg/ml, or from about 1 mg/ml to about 20 mg/ml, or from about 0.1 mg/ml to about 10 mg/ml, or from about 0.5 mg/ml to about 10 mg/ml, or from about 1 mg/ml to about 10 mg/ml.

The disclosed antibodies, antigen binding fragments, conjugates, and nucleic acid encoding such molecules, can be provided in lyophilized form and rehydrated with sterile water before administration, although they are also provided in sterile solutions of known concentration. The antibody solution, or an antigen binding fragment or a nucleic acid encoding such antibodies or antigen binding fragments, can then be added to an infusion bag containing 0.9% sodium chloride, USP, and administered according to standard protocols. Considerable experience is available in the art in the administration of antibody drugs, which have been marketed in the U.S. since the approval of RITUXAN® 1997. Antibodies, antigen binding fragments, conjugates, or a nucleic acid encoding such molecules, can be administered by slow infusion, rather than in an intravenous push or bolus. In one example, a higher loading dose is administered, with subsequent, maintenance doses being administered at a lower level. For example, an initial loading dose of 4 mg/kg may be infused over a period of some 90 minutes, followed by weekly maintenance doses for 4-8 weeks of 2 mg/kg infused over a 30 minute period if the previous dose was well tolerated.

Controlled-release parenteral formulations can be made as implants, oily injections, or as particulate systems. For a broad overview of protein delivery systems see, Banga, A. J., Therapeutic Peptides and Proteins: Formulation, Processing, and Delivery Systems, Technomic Publishing Company, Inc., Lancaster, Pa., (1995). Particulate systems include microspheres, microparticles, microcapsules, nanocapsules, nanospheres, and nanoparticles. Microcapsules contain the therapeutic protein, such as a cytotoxin or a drug, as a central core. In microspheres the therapeutic is dispersed throughout the particle. Particles, microspheres, and microcapsules smaller than about 1 µm are generally referred to as nanoparticles, nanospheres, and nanocapsules, respectively. Capillaries have a diameter of approximately 5 µm so that only nanoparticles are administered intravenously. Microparticles are typically around 100 µm in diameter and are administered subcutaneously or intramuscularly. See, for example, Kreuter, J., Colloidal Drug Delivery Systems, J. Kreuter, ed., Marcel Dekker, Inc., New York, N.Y., pp. 219-342 (1994); and Tice & Tabibi, Treatise on Controlled Drug Delivery, A. Kydonieus, ed., Marcel Dekker, Inc. New York, N.Y., pp. 315-339, (1992).

1

Polymers can be used for ion-controlled release of the antibody compositions disclosed herein. Various degradable and nondegradable polymeric matrices for use in controlled drug delivery are known in the art (Langer, Accounts Chem. Res. 26:537-542, 1993). For example, the block copolymer, 5 polaxamer 407, exists as a viscous yet mobile liquid at low temperatures but forms a semisolid gel at body temperature. It has been shown to be an effective vehicle for formulation and sustained delivery of recombinant interleukin-2 and urease (Johnston et al., Pharm. Res. 9:425-434, 1992; and 10 Pec et al., J. Parent. Sci. Tech. 44(2):58-65, 1990). Alternatively, hydroxyapatite has been used as a microcarrier for controlled release of proteins (Ijntema et al., Int. J. Pharm. 112:215-224, 1994). In yet another aspect, liposomes are used for controlled release as well as drug targeting of the 15 lipid-capsulated drug (Betageri et al., Liposome Drug Delivery Systems, Technomic Publishing Co., Inc., Lancaster, Pa. (1993)). Numerous additional systems for controlled delivery of therapeutic proteins are known (see U.S. Pat. No. 5,055,303; U.S. Pat. No. 5,188,837; U.S. Pat. No. 4,235,871; 20 U.S. Pat. No. 4,501,728; U.S. Pat. No. 4,837,028; U.S. Pat. No. 4,957,735; U.S. Pat. No. 5,019,369; U.S. Pat. No. 5,055,303; U.S. Pat. No. 5,514,670; U.S. Pat. No. 5,413,797; U.S. Pat. No. 5,268,164; U.S. Pat. No. 5,004,697; U.S. Pat. No. 4,902,505; U.S. Pat. No. 5,506,206; U.S. Pat. No. 25 or secondary antibody are described above, and include 5,271,961; U.S. Pat. No. 5,254,342 and U.S. Pat. No. 5,534,496).

## 5. Methods of Detection and Diagnosis

Methods are also provided for the detection of the expression of EBOV GP in vitro or in vivo. In one example, 30 expression of EBOV GP is detected in a biological sample, and can be used to detect EBOV infection as the presence of EBOV in a sample. The sample can be any sample, including, but not limited to, tissue from biopsies, autopsies and pathology specimens. Biological samples also include sec- 35 tions of tissues, for example, frozen sections taken for histological purposes. Biological samples further include body fluids, such as blood, serum, plasma, sputum, spinal fluid or urine. The method of detection can include contacting a cell or sample, or administering to a subject, an 40 E. Kits antibody or antigen binding fragment that specifically binds to EBOV GP, or conjugate there of (e.g. a conjugate including a detectable marker) under conditions sufficient to form an immune complex, and detecting the immune complex (e.g., by detecting a detectable marker conjugated to the 45 antibody or antigen binding fragment.

In several embodiments, a method is provided for detecting EBOV disease and/or an EBOV infection in a subject. The disclosure provides a method for detecting EBOV in a biological sample, wherein the method includes contacting 50 a biological sample from a subject with a disclosed antibody or antigen binding fragment under conditions sufficient for formation of an immune complex, and detecting the immune complex, to detect the EBOV GP in the biological sample. In one example, the detection of EBOV GP in the sample 55 indicates that the subject has an EBOV infection. In another example, the detection of EBOV GP in the sample indicates that the subject has EVD. In another example, detection of EBOV GP in the sample confirms a diagnosis of EVD and/or an EBOV infection in the subject. 60

In some embodiments, the disclosed antibodies or antigen binding fragments are used to test vaccines. For example to test if a vaccine composition including EBOV GP assumes a conformation including the EBOV GP epitope to which EVB100, EVB114, or EVB166 antibody binds. Thus pro- 65 vided herein is a method for testing a vaccine, wherein the method includes contacting a sample containing the vaccine,

such as an EBOV GP immunogen, with a disclosed antibody or antigen binding fragment under conditions sufficient for formation of an immune complex, and detecting the immune complex. Detection of the immune complex confirms that the EBOV GP vaccine includes the epitope to which EVB100, EVB114, or EVB166 antibody, respectively binds. In one example, the detection of the immune complex in the sample indicates that a vaccine component, such as an EBOV GP immunogen assumes a conformation capable of binding the antibody or antigen binding fragment.

In one embodiment, the antibody or antigen binding fragment is directly labeled with a detectable marker. In another embodiment, the antibody that binds EBOV GP (the first antibody) is unlabeled and a second antibody or other molecule that can bind the antibody that binds the first antibody is utilized for detection. As is well known to one of skill in the art, a second antibody is chosen that is able to specifically bind the specific species and class of the first antibody. For example, if the first antibody is a human IgG, then the secondary antibody may be an anti-human-IgG. Other molecules that can bind to antibodies include, without limitation, Protein A and Protein G, both of which are available commercially.

Suitable labels for the antibody, antigen binding fragment various enzymes, prosthetic groups, fluorescent materials, luminescent materials, magnetic agents and radioactive materials. Non-limiting examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, beta-Non-limiting galactosidase, or acetylcholinesterase. examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin. Non-limiting examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin. A non-limiting exemplary luminescent material is luminol; a non-limiting exemplary a magnetic agent is gadolinium, and non-limiting exemplary radioactive labels include <sup>125</sup>I, <sup>131</sup>I, <sup>35</sup>S or <sup>3</sup>H.

Kits are also provided. For example, kits for treating a subject with an EBOV infection, or for detecting EBOV GP in a sample or in a subject. The kits will typically include a disclosed EBOV GP-specific antibody, antigen binding fragment, or nucleic acid molecule encoding such molecules, or compositions including such molecules. More than one of the disclosed EBOV GP-specific antibody, antigen binding fragment, conjugate, or nucleic acid molecule encoding such molecules, or compositions including such molecules can be included in the kit.

In one embodiment, the kit is a diagnostic kit and includes an immunoassay. Although the details of the immunoassays may vary with the particular format employed, the method of detecting EBOV GP in a biological sample generally includes the steps of contacting the biological sample with an antibody which specifically reacts, under conditions sufficient to form an immune complex, to EBOV GP. The antibody is allowed to specifically bind under immunologically reactive conditions to form an immune complex, and the presence of the immune complex (bound antibody) is detected directly or indirectly.

The kit can include a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic. The container typically holds a composition including one or more of the disclosed antibodies,

antigen binding fragments, conjugates, nucleic acid molecules, or compositions. In several embodiments the container may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection <sup>5</sup> needle). A label or package insert indicates that the composition is used for treating the particular condition.

The label or package insert typically will further include instructions for use of the antibodies, antigen binding fragments, conjugates, nucleic acid molecules, or compositions <sup>10</sup> included in the kit. The package insert typically includes instructions customarily included in commercial packages of therapeutic products that contain information about the indications, usage, dosage, administration, contraindications and/or warnings concerning the use of such therapeutic products. The instructional materials may be written, in an electronic form or may be visual. The kits may also include additional components to facilitate the particular application for which the kit is designed. Thus, for example, the kit may additionally contain means of detecting a label (such as 20 enzyme substrates for enzymatic labels, filter sets to detect fluorescent labels, appropriate secondary labels such as a secondary antibody, or the like). The kits may additionally include buffers and other reagents routinely used for the practice of a particular method. Such kits and appropriate <sup>25</sup> contents are well known to those of skill in the art.

### III. Examples

The following examples are provided to illustrate particu-<sup>30</sup> lar features of certain embodiments, but the scope of the claims should not be limited to those features exemplified.

## Example 1

## EBOV GP-Specific Monoclonal Antibodies that Neutralize EBOV Infection

This examples illustrates the isolation and characterization of the EVB100, EVB114, and EVB166 antibodies, 40 which specifically bind to EBOV GP and can neutralize EBOV.

Ebola virus disease (EVD) causes severe illness characterized by rapid onset of fever, vomiting, diarrhea and bleeding diathesis, and was first described in the Democratic 45 Republic of Congo in 1976. The 2014 outbreak in West Africa has affected over 27,000 and claimed at least 11,000 lives. The challenges of a large outbreak and the failure of traditional quarantine and contact tracing measures to control this outbreak highlights the urgency for therapies. The 50 success in nonhuman primates (NHP) of ZMapp, a cocktail of three mouse-human chimeric mAbs derived from immunized mice (Qiu et al., Clin. Immunol. 141, 218-27, 2011; Wilson et al., Science. 287, 1664-6, 2000), illustrated the potential impact of monoclonal antibody therapies against 55 EVD, and it is currently being evaluated in human trials. To date, efforts to simplify the ZMapp regimen to contain fewer mAbs have not been successful in the macaque EVD model (Qiu et al., Nature. 514, 47-53, 2014). Accordingly, mAbs were isolated from human survivors of Ebola virus infec- 60 tion, with the goal of identifying antibodies that confer clinical protection either as single or dual-combination agents.

Blood was obtained from two survivors of the 1995 Kikwit EVD outbreak (Muyembe-Tamfum et al., *J. Infect.* 65 *Dis.* 179 Suppl, S259-S262, 1999) eleven years after infection. These subjects were the sole survivors of a family of 15 68

people who were infected during the outbreak. At the time of infection, subject 1 (51) was a male 28-year-old who had severe laboratory-confirmed illness and, following recovery, worked for several months in the EVD ward caring for other patients. His sister (S2) was 20-years old and had moderate disease severity that was clinically diagnosed based on contact history and symptoms. To determine if the subjects retained circulating antibodies against Ebola virus (EBOV) glycoprotein (GP), GP-specific antibodies were assessed by ELISA (FIG. 1A). Reciprocal EC90 titers of 2,326 and 275 in the sera of 51 and S2 were observed, respectively. Moreover, the serum from 51, the more severely ill subject, displayed potent virus neutralizing activity (FIG. 1B). The results indicated that these survivors maintained serologic memory against EBOV GP more than a decade following infection, and suggested the potential to clone immunoglobulins with potent neutralizing activity from their memory B cells.

IgG memory B-cells were sorted from S1's peripheral blood mononuclear cells (PBMC), and immortalized individual clones with Epstein-Barr virus (using techniques described in Traggiai et al., *Nat. Med.* 10, 871-875, 2004). Forty immortalized clones whose supernatants displayed a range of GP-binding activity by ELISA were identified (FIG. 1C). Two mAb clones, mAb100 and mAb114 (termed EVB100 and EVB114), expressed antibodies with markedly higher neutralizing activity than all others (FIG. 1D).

A second immortalization yielded 21 clones, from which two additional GP-specific mAb clones, 165 and 166 (termed EVB165 and EVB166), were rescued (see the following table). In the second screening of immortalized memory B-cells from 51, 14 million PBMCs were used to isolate 59,500 IgG memory B cells which were immortalized as in FIGS. 1A-1D. After removal of non-specific binding, 21 culture supernatants were found to specifically bind Ebola GP as measured by ELISA. Shown are ELISA A450 values for undiluted and 1:27 dilutions of the supernatants. Amongst the 21 supernatants, only 2 B cell clones (EVB165, EVB166) were rescued for further analysis.

Cell	Abs 450		
line ID	Und.	1/27	
mAb151	1.119	0.162	
mAb152	0.672	0.106	
mAb153	0.854	0.131	
mAb154	2.361	0.95	
mAb155	0.115	0.08	
mAb156	1.111	0.161	
mAb157	2.256	0.554	
mAb158	0.074	0.075	
mAb159	3.298	2.529	
mAb160	1.493	0.489	
mAb161	0.227	0.086	
mAb162	0.083	0.074	
mAb163	3.099	1.805	
mAb164	0.076	0.069	
EVB165	3.171	1.722	
EVB166	2.894	2.4	
EVB167	3.368	2.974	
mAb168	0.507	0.114	
mAb169	0.081	0.072	
mAb170	0.95	0.165	
mAb171	1.998	0.895	

Immunoglobulin sequences were PCR-amplified from the four clones and used to produce EVB100, EVB114, EVB165 and EVB166 by transient transfection. ELISA binding to EBOV GP was assessed and it was observed that one antibody, EVB114, stood apart from the others, displaying nearly 100% higher maximal binding (FIG. 2A). The remaining three antibodies, EVB100, EVB165 and EVB166, exhibited reduced levels of maximal binding compared to EVB114, but were comparable to each other and to 5 KZ52, a prototype human EBOV GP-specific mAb (Maruyama et al., *J. Virol.* 73, 6024-6030, 1999). EVB114 achieved half maximal binding (EC50) at a concentration of 0.07 µg/mL, which was up to two orders of magnitude lower EC50 than the other mAbs. EVB100 and EVB166 had 10 similar binding profile (0.26 µg/mL, 0.40 µg/mL) while EVB165 bound less well with an EC50>1 Kg/mL.

To test potential functional properties of the mAbs the inhibition of GP mediated entry into HEK293T cells was evaluated in the absence of complement (FIG. **2**B) A sum- 15 mary of in vitro neutralization IC data (obtained as described for FIG. **2**B) is provided in the following table:

binding. It is intriguing that these mutations (A96V and Y108S) are located at the base of the HCDR3 loop which are most likely not in direct contact with GP but may have a stabilizing effect on the whole HCDR3. Indeed, restoration to the binding equivalent of the mature antibody required a fully matured light chain in addition to the two HCDR3 mutations. Inherent uncertainty in determining the germline configuration of the HCDR3 does not appear to apply to this case since the two mutations are located in the V and J regions of the junction and no polymorphisms have been described at those positions. Importantly, the fully mutated light chain gene, as shown in the case of the EVB114 UCA/sK variant, can partially compensate for the lack of somatic mutation in the heavy chain (FIG. 2I). The presence of additional mutations on either VH or VK is required to achieve the level of the fully matured EVB114 binding. These results suggest a rapid pathway of EVB114 affinity

mAb	$IC50~(\mu g/mL)$	95% CI	$IC90~(\mu g/mL)$	95% CI	$IC99~(\mu g/mL)$	95% CI	n
KZ52	0.06	0.02 to 0.14	17.21	8.47 to 35.00	>>1000	54,868	6
EVB100	0.06	0.05 to 0.08	0.61	0.39 to 0.93	7.58	2.999 to 19.16	6
EVB114	0.09	0.07 to 0.11	0.71	0.44 to 1.16	7.19	2.588 to 19.96	6
EVB166	0.86	0.72 to 1.02	6.84	4.78 to 9.80	97.25	31.32 to 138.2	4
EVB165	1.77	1.43 to 2.18	19.46	13.23 to 28.61	267.00	124.9 to 570.8	4

EVB165 and EVB166 both neutralized well and exhibited similar potencies for half maximal inhibition (IC50) concentration of 1.77 and 0.86  $\mu$ g/ml, respectively. EVB100 and EVB114 resided in the strongest neutralizing group, with IC50 about one-log greater (0.06 and 0.09  $\mu$ g/ml, respectively) than EVB165 and EVB166. Notably, all four of the neutralizing antibodies inhibited 100% of the input virus unlike KZ52, which consistently displayed only 80-90% at 10  $\mu$ g/mL. Importantly, neutralization of the 2014 West African Makona variant was achieved within similar concentration ranges seen for the Mayinga variant (FIG. **5**).

Sequence analysis revealed EVB114 and EVB165 to be 40  $IgG_1$  isotypes, and EVB100 and EVB166 to be  $IgG_3$  isotypes. Immunoglobulins displayed between 85-95% and 89-97% germline identity for heavy and light chains, respectively (FIG. **2**C). Analyses of germline gene usage and V(D)J recombination indicate that they originate from dif-45 ferent B-cell lineages. Interestingly, EVB114 utilizes IGHV3-13\*01, a rarely used VH gene, and IGKV1-27\*01.

The role of somatic hypermutations for the two most potent antibodies, EVB100 and EVB114, was analyzed using variants that were partially or completely reverted to 50 the unmutated common ancestors (UCAs) (FIGS. 2D to 2G). The fully reverted version of EVB100 (UCA/UCA), as well as a variant with germline VH and a VL with a single change from germline (A89T), recognized cells expressing GP with only a 2- to 4-fold weaker binding compared with the fully 55 matured antibody (FIG. 2H). GP binding comparable to the fully matured EVB100 heavy and light chains (sH/sL) was observed when three HCDR3 mutations (A96V/V103Y/ Y114S) were introduced in the reverted germline antibody (gH/UCA), illustrating that those mutations were sufficient 60 to mediate binding observed with fully matured EVB100. The addition of all the other mutations did not contribute further to EVB100 binding to GP. In the case of the EVB114, the fully reverted version of EVB114 (UCA/UCA) demonstrated negligible binding to EBOV GP (FIG. 2I). Introduc- 65 tion of two mutations (A96V and Y108S) in the HCDR3 of EVB114 germline was sufficient to confer an increase in GP

maturation through one or two somatic mutations, which became redundant as further mutations accumulated, a finding that is reminiscent of what was recently observed for the generation of broadly neutralizing influenza antibodies (Pappas et al., *Nature.* 516, 418-22, 2014).

Since EVB100 and EVB114 were the most potently neutralizing antibodies, they were considered optimal candidates for further evaluation. The potential for synergy between these antibodies in the context of combination therapy was assessed. First, to possibility of cross-competition for antigen binding or targeting of a single immunodominant region of GP was assessed. It was found that each antibody bound to GP in the presence of the other, suggesting that they recognize distinct regions on GP (FIG. 3A) and therefore could be used together in combination immunotherapy to improve efficacy and diminish the likelihood of emergence viral escape mutants (Barouch et al., Nature. 503, 224-8, 2013; Shingai et al., Nature. 503, 277-80, 2013). To define the regions targeted by EVB100 and EVB114, biolayer interferometry was employed to assess GP binding in competition with mAbs KZ52 and 13C6, which have known epitopes in the GP base and glycan cap, respectively (Lee et al., Nature. 454, 177-82, 2008; Murin et al., Proc. Natl. Acad. Sci. U.S.A 111, 17182-17187, 2014). It was found that EVB100 competes with KZ52 for binding at the base of GP, while EVB114 recognizes at least in part the glycan cap region, as demonstrated by the partial competition observed with 13C6 (FIGS. 3B and 3C). Since some EBOV GP antibodies have been suggested to mediate antibody dependent cell-mediated cytotoxicity (ADCC) (Olinger et al., Proc. Natl. Acad. Sci. U.S.A 109, 18030-5, 2012) the ADCC activity of EVB100 and EVB114 were determined in a flow cytometry-based assay using GPexpressing target cells (FIG. 3D). It was found that both EVB100 and EVB114 mediated ADCC, and maximum activity was observed at a mAb concentration of 0.03 µg/ml, which is similar to the IC50 values for neutralization Killing of target cells was demonstrated to be mediated through Fc receptors since LALA mutations in the mAb Fc regions (Hezareh et al., J. Virol. 75, 12161-12168, 2001) of the antibodies abrogated ADCC activity. Therefore, in addition to neutralization, these mAbs have the potential to induce direct killing of infected cells in vivo, a key viral clearance mechanism.

The presence of potent neutralizing and ADCC activity. and the absence of cross competition, supported testing EVB100 and EVB114 in vivo for protective efficacy in macaques. Four rhesus macaques (Macaca mulatta) were challenged with a lethal dose of Ebola virus, Kikwit 1995 variant. One day post-challenge, the treatment group (n=3) was given an intravenous injection with a mixture of EVB100 and EVB114 at a total combined dose of 50 mg/kg, and the treatment was repeated twice more at 24-hour intervals (FIG. 4A). Circulating Ebola GP-specific antibody titers in the mAb recipients peaked after the second mAb injection and reciprocal ELISA titers remained above 10<sup>5</sup> throughout the study, suggesting minimal clearance of the mAbs during the observation period (FIG. 4B). The naive untreated macaque succumbed to EVD on day 10 with a 20 circulating viral load exceeding 10<sup>8</sup> ge/ml (FIGS. 4C and 4D). In contrast, all three mAb treated macaques survived challenge without detectable systemic viremia. Consistent with historic controls, the untreated animal displayed hallmark indicators of Ebola infection including hematologic, 25 liver and renal dysfunction as indicated by thrombocytopenia and striking elevations in alanine transaminase (ALT) and creatinine from day 6 through the time of death (FIGS. 4E, and 6-8). In contrast, macaques in the treatment group remained within normal ranges for these parameters, and 30 remained free of all EVD symptoms.

Assays were also performed to determine if mAb monotherapy is sufficient for protection of NHP. Initial testing focused on EVB114, since it showed higher maximal binding than EVB100. As in the first experiment four macaques 35 were exposed to a lethal dose of EBOV and administered 50 mg/kg of EVB114 (n=3) to the treatment group after a one-day delay, followed by two more doses at 24-hour intervals. All treated macaques survived, whereas the control animal succumbed to EVD on day 6 with a peak viral load 40 of  $10^{10}$  ge/ml (FIGS. 4F to 4H). In contrast to the previous experiment, transient viremia was observed in the treated animals, but it remained at levels less than 0.1% of the untreated control animal, and returned to undetectable levels. Despite transient viremia, treated animals remained free 45 of clinical and laboratory abnormalities (FIGS. 41, and 9-11).

The neutralization activity of the EVB100, EVB114, EVB165, and EVB166 antibodies was further assayed in the presence of soluble GP (sGP), which is believed to interfere with the natural immune response to EBOV in human subjects (FIG. **12**). The neutralization assay was performed as above and using the IC50 concentration of each antibody in the presence or absence of sGP. As shown in FIG. **12**, the neutralization potency of EVB100 was not affected by the presence of sGP; however, neutralization by the EVB114, EVB165, and EVB166 antibodies was diminished in this in vitro assay.

EBOV GP is initially synthesized as precursor protein  $(GP_0)$  that is cleaved by cellular furin to form the surface 60 glycoprotein GP<sub>1</sub> and the membrane anchoring protein GP<sub>2</sub>. GP<sub>1</sub> and GP<sub>2</sub> are linked in the mature virion trimer by disulfide bonds. There are other GP forms created during cellular trafficking and processing that may also bind to antibodies in the infected host. Since binding specificity to 65 the various GP forms may influence in vivo efficacy of the antibodies, the EVB100, EVB114, EVB165, and

EVB166 antibodies were assayed to determine if they could immunoprecipitate GP (FIG. 13). KZ52 was used as a positive control. As shown in FIG. 13, each of the EVB100, EVB114, EVB165, and EVB166 antibodies can immunoprecipitate GP. EVB114 immunoprecipitated GP<sub>1</sub> and GP<sub>2</sub>, suggesting that it binds to the mature, disulfide-linked trimer. EVB114 also immunoprecipitated the lower molecular weight cathepsin-cleaved form of GP (GPCatL). The EVB100, EVB165, and EVB166 antibodies bind to precursor forms of GP present prior to furin or cathepsin cleavage.

To further assess the binding properties of the EVB100, EVB114, EVB165, and EVB166 antibodies for GP, several GP deletion mutants were assayed for binding to these antibodies (FIGS. **14** and **15**). GP deletion mutants were constructed that lack the N-terminus (containing the putative receptor binding domain, (d47-277, "GP&RBS"), the highly glycosylated mucin-like PdMUC"), or the C terminus of EBOV GP (d494-635, "GP dGP2") (see FIG. **14**). Additionally, binding to sGP was also assessed (FIGS. **17**A and **17**B).

Gross epitope mapping was performed by expressing the GP mutants in mammalian cells, followed by immunoprecipitation using the EVB100, EVB114, EVB165, or EVB166 antibodies and detection of precipitated protein by Western blot with EBOV GP specific antibody (FIGS. 15-17). Deletion of the GP mucin-like domain reduced recognition by EVB100 in immunoprecipitation assays, but the other antibodies recognized GP in the absence of the mucin-like domain (FIG. 15). This finding indicates that deletion of the mucin-like domain alters (or deletes) at least part of the EVB100 epitope on EBOV GP. Further, deletion of GP<sub>2</sub> reduced recognition by EVB100 in immunoprecipitation assays, but the other antibodies recognized GP in the absence of the GP<sub>2</sub> polypeptide (FIG. 16). This finding indicates that deletion of the GP<sub>2</sub> polypeptide alters (or deletes) at least part of the EVB100 epitope on EBOV GP.

The EVB100, EVB114, EVB165, and EVB166 antibodies were also assessed for binding to soluble GP (sGP) by immunoprecipitation (FIG. 17A). sGP contains the GP receptor binding domain; therefore, the N-terminal portions of sGP and GP as found on the viral envelope are homologous in that they each contain putative receptor binding domain. EVB114, EVB165, and EVB166 all immunoprecipitated sGP indicating that each of these antibodies binds to an epitope within the N-terminal portion of the GP protein, which includes the receptor binding domain. Further, recognition by direct WB suggests that binding is to a linear, not conformation-dependent epitope (FIG. 17B). In contrast, EVB100 does not bind to N-terminal GP sequences by either direct WB or IP/WB that preserves conformation. This is consistent with the previous finding that EVB100 binding requires elements in the mucin and GP<sub>2</sub> regions of GP.

The EVB100, EVB114, EVB165, and EVB166 monoclonal antibodies were further assayed for neutralization of the Bundibugyo (BDBV) and Sudan (SUDV) Ebola virus strains, which have previously causes highly lethal outbreaks in humans (FIG. 18). Cross-species neutralization was assayed using a single-round replication assay with virus pseudotyped with the envelope glycoprotein (GP) from each species. As shown in FIG. 18, the EVB100 and EVB166 antibodies potently neutralized BDBV, while EVB114 and EVB165 displayed modest cross recognition and neutralization. EVB166 was additionally neutralized the SUDV pseudovirus.

EVB114 has several characteristics that may contribute to protection as a monotherapy compared to KZ52 and 13C6,

which were non-protective in NHPs (Qiu et al., Nature. 514, 47-53, 2014; Oswald et al., PLoS Pathog. 3, e9, 2007). Firstly, both KZ52 and EVB114 neutralize with potent IC50s, however

EVB114 neutralizes 100% of input virus whereas KZ52 5 plateaus at 80-90%. Secondly, EVB114 does not require complement for neutralizing activity in contrast to 13C6 (FIG. 2B) (Wilson et al., Science. 287, 1664-6, 2000). Based on these observations, one non-limiting conclusion is that protective monotherapy requires both potent binding and 10 complete neutralization in the absence of complement. In addition, ADCC activity may contribute to the unique ability of EVB114 to protect as a monotherapy against lethal Ebola infection of macaques.

Data presented in this example shows that circulating 15 functional antibodies as well as memory B cells specific to Ebola virus are maintained in survivors for more than a decade following infection. mAbs isolated from a survivor of the 1995 Kikwit EVD outbreak exhibited ADCC activity and showed potent neutralizing activity against two other 20 Ebola variants, including one from the recent West Africa outbreak. Macaques who received EVB114 and EVB100 as combination therapy remained healthy with no signs of viremia after EBOV challenge. Strikingly, when a single antibody, EVB114, was therapeutically administered after 25 lethal EBOV challenge of macaques, all treated animals were fully protected and asymptomatic, despite a low transient level of circulating virus being detected.

Materials and Methods

Isolation of Monoclonal Antibodies from EBOV Survi- 30 vors.

Two subjects who survived the 1995 EBOV Kikwit variant outbreak in the Democratic Republic of Congo were identified and enrolled in VRC200 clinical trial #NCT00067054 after giving signed informed consent. 35 Peripheral blood mononuclear cells (PBMCs) were obtained, stained with directly labeled antibodies to CD22 (Pharmingen) and to immunoglobulin IgM, IgD, and IgA. CD22+IgM-IgD-IgA-B cells were isolated using FACS Aria, pulsed with Epstein-Barr Virus (50% B958 superna- 40 of EVD survivor's polyclonal sera, monoclonal antibodies tant) and seeded at 30 cells/well (for a total of 2.7×105 purified cells) in replicate cultures in medium supplemented with CpG 2006 and irradiated allogeneic PBMCs, as previously described (Traggiai et al., Nat. Med. 10, 871-875, 2004). Culture supernatants were collected after 2 weeks and 45 tested for binding to ELISA plates coated with EBOV GP (Mavinga variant), their specificity was confirmed using an unrelated antigen (tetanus toxoid) and positive cultures were further tested for their ability to neutralize EBOV pseudoviruses. Cultures that scored positive in the EBOV neutral- 50 ization assay were subcloned by limiting dilution.

Antibody Purification, Labeling, Genetic Analysis, and Reversion to Germline.

The usage of VH and VL gene segments was determined by sequencing, and analysis for homology to known human 55 V, D, and J genes was performed using the IMGT database (http://www.imgt.org/). Human antibodies were affinity purified by protein A chromatography (GE Healthcare) and dialyzed against PBS. Selected antibodies were biotinylated using the EZ-Link NHS-PEO Solid Phase Biotinylation Kit 60 (Pierce).

Antibodies were also produced recombinantly by cloning VH and VL genes via PCR into human Igy1, Igk (EVB114, 165, 166), and Ig $\lambda$  (EVB100) expression vectors using gene specific primers (Tiller et al., J. Immunol. Methods. 329, 65 112-124, 2008). Antibodies used for animal studies were produced by transient transfection of suspension cultured

293FreeStyle cells (Invitrogen) with PEI or Expi cells with Expifectamine293 (Invitrogen). Supernatants from transfected cells were collected after 6-10 days of culture and IgGs were affinity purified by Protein A chromatography (GE Healthcare) and dialyzed against PBS. Purified mAbs were then concentrated with Amicon Ultra centrifugal filters and sterilized by 0.22 µm filtration. The purity was assessed by SEC-HPLC and SDS-PAGE. Endotoxin content was measured with the Endpoint Chromogenic LAL assay (QCL-1000 TM assay, Lonza) according to manufacturing instructions and shown to be below 0.25 EU/ml. Antibody concentrations were determined using the BCA Protein Assay Kit (Thermo Scientific) using Rituximab (Roche) as internal standard or A280 using an Nanodrop (Thermo Scientific). Germlined VH and VL nucleotide sequences were synthesized by Genscript, and their accuracy was confirmed by sequencing.

Antibodies.

KZ52 monoclonal antibody used in ELISA assay a kind gift from Dennis Burton. KZ52 used elsewhere and 13C6 was purchased from IBT Bioservices. Unless otherwise noted isotype control antibody was an anti-HIV gp120 IgG1. Antibody Neutralization Assay.

Supernatants or purified mAbs from immortalized B cell clones isolated from EVD survivor donors were assessed for neutralization potency using a single-round infection assay with EBOV GP-pseudotyped lentiviruses particles which express a luciferase reporter gene following entry (Sullivan et al., PLoS Med. 3, e177, 2006). Unless indicated, all experiments utilized particles bearing GP from the EBOV Mayinga variant. In brief, HEK293T cells were used as infection targets and incubated in a 96-well plate 1 day before infection with pseudovirus in the presence of serially diluted supernatant or purified mAbs. Infected target cells were lysed 72 hours after infection and assayed with the Luciferase Assay System or Bright Glo (Promega), using a Victor X3 Plate Reader (PerkinElmer) to detect luciferase

activity. ELISA for serum antibody titer and GP-binding. Binding and antibody in non-human primates to EBOV GP was evaluated by enzyme-linked immunosorbent assay (ELISA) as described previously (Sullivan et al., PLoS Med. 3, e177, 2006). Titers for survivor and non-human primates were calculated as reciprocal EC90 values (Sullivan et al., PLoS Med. 3, e177, 2006).

Ebola Virus GP Vectors.

Plasmid vector pVR1012 WT GP (Z) has been described previously (21). A vector expressing a soluble mucin deleted (ΔMuc) GP, GPΔMucΔTM-GCN4 HisSA (4309-505, Δ657-676), was made using codon optimization and then synthesized and directly cloned in frame to a GCN4 trimerization domain-His-Strep Tactin domains (MKQIEDKIEEILSKI-YHIENEIARIKKLIGEVASSSIEGRGSHHHHHHSAW-

SHPQFEK, SEQ ID NO: 66) and sequence verified by Genscript. EBOV GP variant Makona-005 (Acc#KJ660348) was codon optimized, synthesized and sequence verified by Genscript.

Antibody-Dependent Cell-Mediated Cytotoxicity (ADCC).

rAd5 EBOV GP-transduced and non-transduced HEK293T cells were double labeled with membrane-bound and intracellular stains in order to detect ADCC activity. Cells were incubated with 8 µM Plum stain (Plum cell labeling kit M.T.T.I. CellVue) followed by FBS. The cells were then washed with RPMI 1640, incubated with 5 µM Carboxyflourescein Succinimidyl ester (CFSE) (Vybrant

55

65

CFDA SE cell tracer kit, Invitrogen), incubated with FBS and washed again with RPMI 1640. Doubly labeled EBOV GP expressing cells were plated in a V-bottomed 96-well plate at 5,000 cells/well. Antibodies were added to duplicate samples at 31.6 ng/ml to the target cells for 20 minutes at 5 room temperature. RSV antibody (palivizumab) was used as a control antibody. Effector cells resuspended in RPMI were then added to the target cells at the effector-to-target cell (ET) ratio 1:50 which was found to give the best signal to noise ratio. Each plate was incubated for 4 hr at 37° C./5% CO2. After 4 hr, plates were centrifuged at 250×g and cells were fixed with 1% Paraformaldehyde (PFA) and analyzed via flow cytometry. As a control, labeled non-transduced HEK293T cells were also used as targets for ADCC activity. Thirty thousand non-gated events were acquired within 6 hr 15 after the ADCC assay using an LSRII cytometer (Becton Dickinson). The CFSE emission channel was read in B515 using a neutral density filter and Plum emission was read in R660. Following acquisition, analysis was performed using FlowJo software (Tree Star). Percent killing was obtained by 20 quantifying dead cells (Plum+,CFSE-) out of the total Plum positive population. For mAbs, ADCC killing was measured by subtracting percent killing of nontransduced cells from percent killing of transduced cells.

Antibody Variants.

UCA sequences of the isolated antibodies were determined with reference to the IMGT database (imgt.org). Antibody variants in which single or multiple mutations were reverted to the germline sequence were produced by gene synthesis (Genscript) and used to produce a large set of 30 EVB114 and EVB100 antibody variants.

Binding of antibody variants to transfected cells.

EVB114 and EVB100 antibody variants were used to stain MDCK-SIAT1 cell lines transduced to express EBOV GP as a stable membrane protein (Makona variant). Binding 35 of antibodies was analysed using a Becton Dickinson FACS Canto2 (BD Biosciences) with FlowJo software (TreeStar). The relative affinities of antibody binding to surface GP were determined by interpolating the concentration of antibody required to achieve 50% maximal binding (EC50) 40 from the plotted binding curves using the mean-fluorescence intensity (MFI) fitted with a 4-parameter nonlinear regression with a variable slope.

Inhibition of Binding Assay on GP-Expressing Cells.

EVB100 and EVB114 were biotinylated using the EZ- 45 Link NHS-PEO solid phase biotinylation kit (Pierce). Labeled antibodies were tested for binding to GP-expressing MDCK-SIAT-1 cells to determine the optimal concentration of each antibody to achieve 70-80% maximal binding. The biotin-labelled antibodies were then used as probes to 50 assess, by flow cytometry, whether their binding (measured using fluorophore-conjugated streptavidin) was inhibited by preincubation of GP cells with homologous or heterologous unlabelled antibodies.

Production of Purified GP.

Expi (Invitrogen) cells were transfected with GP  $\Delta$ Muc $\Delta$ TM-GCN4 HisSA and pCMV-Sport Furin (7:3 ratio) using 293Fection (Invitrogen) at a ratio of 2 mL 293Fectin: Img total DNA. 18-24 hours following transfection, <sup>1</sup>/10th volume of AbBooster (ABI Scientific) was added and cul- 60 ture media collected 5 days later. Supernatant was filtered and protein purified as described previously (Cote et al., *Nature*. 477, 344-348, 2011).

Biolayer Interferometry Antibody Cross-Competition Assay.

Antibody cross competition was determined based on biolayer interferometry using a fortéBio Octet HTX instru-

76

ment. EBOV GP AMuc protein was loaded onto HIS biosensors (AR2G, fortéBio) through amine coupling for 600 s. Biosensors were equilibrated for 120 s in 1% BSA in PBS (BSA-PBS) prior to capturing competitor mAbs. GP proteins were diluted to 10 µg/mL; mAbs KZ52, EVB100, EVB114, 13C6, and IgG1 isotype control Ab were diluted to 35 µg/mL in BSA-PBS. Binding of competitor mAbs was assessed for 300 s followed by a brief equilibration for 60 s prior to binding assessment of probing mAbs. Binding of probing mAbs was assessed for 300 s. Percent inhibition (PI) of probing mAbs binding to GP by competitor mAbs was carried out by an equation: PI=100-[(probing mAb binding in the presence competitor mAb)/(probing mAb binding in the absence of competitor mAb)]×100. All the assays were performed in duplicate and with agitation set to 1,000 rpm at 30° C.

Animal Study and Safety.

Research was conducted under an IACUC-approved protocol in compliance with the Animal Welfare Act, PHS Policy, and other Federal statutes and regulations relating to animals and experiments involving animals. The facilities where this research was conducted are accredited by the Association for Assessment and Accreditation of Laboratory Animal Care, International and adhere to principles stated in the Guide for the Care and Use of Laboratory Animals, National Research Council, 2011. Animal study protocols were approved by both the Vaccine Research Center and United States Army Medical Research Institute of Infectious Diseases IACUCs. All animals were Vietnamese-origin rhesus macaques (Macaca mulatta), female, approximately 2-5 years of age and were obtained from Covance. Animals were randomly assigned to treatment groups based on sequential selection from a population inventory. Sample sizes of three animals per BSL4 EBOV challenge group provide 80% power to detect a difference in survival rates assuming 100% survival (3/3 treated survive) vs. 0% survival in negative controls at the 95% confidence level (1-tailed Fisher exact test). Prior to blood sampling or treatment, animals were anesthetized with ketamine or telazol.

Antibody Administration.

In the EVB100/EVB114 cocktail challenge, antibodies were mixed in PBS at 4 mg/mL of EVB100 and 46 mg/mL of EVB114 for a total antibody concentration of 50 mg/mL. In the second challenge, animals received 50 mg/mL of EVB114 in PBS. Antibodies were administered via intravenous injection in peripheral veins using  $\leq$ 20 gauge butterfly needles over a period  $\geq$ 15 minutes in a single bolus via syringe pump.

EBOV Challenge.

Animal studies conducted at USAMRIID were approved by the IACUC. Animals were transferred one week prior to challenge to the Bio-Safety Level-4 (BSL-4) facility for exposure to a lethal (1000 PFU) i.m. EBOV Kikwit variant challenge. Challenge studies included a single unvaccinated animal (control); the use of historical control (n>50) allows for one untreated control to be used in each challenge experiment. While at USAMRIID the monkeys were fed and checked daily. During the EBOV challenge study, blood was collected from the NHP for hematological, biochemical and virological analyses. Following the development of clinical signs, animals were checked multiple times daily. Institute scoring criteria were used to determine timing of humane euthanasia under anesthesia.

Detection of EBOV.

RNA was isolated from plasma of EBOV-exposed NHP by real time qPCR as described previously (Malhotra et al., *PLoS* Negl. Trop. Dis. 7, e2171, 2013). EDTA plasma was
added to TriReagent LS (Sigma), 1 part to 3 parts, in preparation for qRT-PCR. Inactivated samples were Extracted and eluted with AVE Buffer (QIAGEN, Valencia, Calif.) using a QIAamp Viral RNA Mini Kit (Qiagen, Valencia, Calif.). All samples were run on an Applied Biosystems 7500 Fast Dx Real-Time PCR instrument (Life Technologies, Grand Island, N.Y.). Reactions were performed with SuperScript II One-Step RT-PCR System (Life Technologies, Grand Island, N.Y.) with additional MgSO4 added to a final concentration of 3.0 mM. All samples were 10 run in triplicate 5 µL each. The average of the triplicates was multiplied by 200 to obtain genomes equivalents per mL, then multiplied by a dilution factor of 4 for the final reported value. The sequence of the primer and probes for the EBOV glycoprotein are described below. The genomic equivalents <sup>15</sup> were determined using a synthetic RNA standard curve of known concentration. Forward primer: 5'-TTTTCAATCCT-CAACCGTAAGGC (SEQ ID NO: 63)-3'; REVERSE PRIMER: 5'-CAGTCCGGTCCCAGAATGTG (SEQ ID NO: 64)-3'; PROBE: CCCATCGCTGC (SEQ ID NO: 65)-TAMRA

#### Example 2

#### Antibodies Specific to EBOV GP for Detecting EBOV in a Sample or a Subject

This example describes an exemplary use of EBOV monoclonal neutralizing antibodies specific to EBOV GP for the detection of EBOV in a sample or a subject. This 30 example further describes the use of these antibodies to confirm the diagnosis of EBOV infection in a subject.

A biological sample, such as a blood sample, is obtained from the patient diagnosed with, undergoing screening for, or suspected of having, an EBOV infection. A blood sample 35 can be taken from a patient who is not infected is used as a control, alternateively, a standard result can also be used as a control. An ELISA is performed to detect the presence of EBOV in the blood sample. Proteins present in the blood samples (the patient sample and control sample) are immo- 40 bilized on a solid support, such as a 96-well plate, according to methods well known in the art (see, for example, Robinson et al., Lancet 362:1612-1616, 2003, incorporated herein by reference). Following immobilization, EBOV monoclonal neutralizing antibodies specific to EBOV GP 45 that are directly labeled with a fluorescent marker are applied to the protein-immobilized plate. The plate is washed in an appropriate buffer, such as PBS, to remove any unbound antibody and to minimize non-specific binding of antibody. Fluorescence can be detected using a fluorometric 50 plate reader according to standard methods. An increase in fluorescence intensity of the patient sample, relative to the control sample, indicates the EBOV GP antibody specifically bound proteins from the blood sample, thus detecting the presence of EBOV protein in the sample. Detection of 55 EBOV protein in the patient sample indicates the patient has EBOV infection, or confirms diagnosis of EBOV in the subject.

#### Example 3

#### EBOV Monoclonal Neutralizing Antibodies Specific for EBOV GP for the Treatment of EBOV

This example describes a particular method that can be 65 used to treat EBOV infection in a human subject by administration of one or more EBOV GP-specific neutralizing

antibodies or antigen binding fragments. Although particular methods, dosages, and modes of administrations are provided, one skilled in the art will appreciate that variations can be made without substantially affecting the treatment. Screening Subjects

In particular examples, the subject is first screened to determine if they have an EBOV infection. Examples of methods that can be used to screen for EBOV infection include evaluating the patient for EBOV symptoms (e.g., hemorrhagic fever), determining prior exposure to EBOV infected subjects or EBOV materials (e.g., bodily fluids from an EBOV infected patient), and/or measuring the levels of one or more EBOV proteins or nucleic acid in a biological sample from the subject (e.g., assaying for EBOV sGP in a blood sample from the subject).

In some examples, EBOV testing consists of initial screening with an enzyme-linked immunosorbent assay (ELISA) to detect antibodies to an EBOV protein, such as to EBOV GP. Specimens with a reactive ELISA result are 6FAM-CATGTGCCGC- 20 retested in duplicate. If the result of the duplicate test is reactive, the specimen is reported as repeatedly reactive and undergoes confirmatory testing with a more specific supplemental test (e.g., Western blot or an immunofluorescence assay (IFA)). Specimens that are repeatedly reactive by ELISA and positive by IFA or reactive by Western blot are considered EBOV-positive and indicative of EBOV infection. In additional examples, nucleic acid testing (e.g., viral RNA or proviral DNA amplification method) can also help diagnosis in certain situations.

> The detection of EBOV protein in a subject's blood is indicative that the subject is infected with EBOV and is a candidate for receiving the therapeutic compositions disclosed herein. However, pre-screening is not required prior to administration of the therapeutic compositions disclosed herein.

Administration of Therapeutic Compositions

Following subject selection, a therapeutically effective amount of an EBOV GP-specific neutralizing mAb described herein (e.g., EVB114 or EVB100) or a combination of such mAbs is administered to the subject (such as an adult human either at risk for contracting EBOV or known to be infected with EBOV). Additional agents, such as anti-viral agents, can also be administered to the subject simultaneously or prior to or following administration of the disclosed mAb. Typically the antibody is administered intravenously.

The amount of the antibody administered to prevent, reduce, inhibit, and/or treat EBOV or a condition associated with it depends on the subject being treated, the severity of the disorder, and the manner of administration of the therapeutic composition. Ideally, a therapeutically effective amount of an agent is the amount sufficient to prevent, reduce, and/or inhibit, and/or treat the condition (e.g., EBOV or EVD) in a subject without causing a substantial cytotoxic effect in the subject. An effective amount can be readily determined by one skilled in the art, for example using routine trials establishing dose response curves. As such, these compositions may be formulated with an inert diluent or with a pharmaceutically acceptable carrier.

In one specific example, a subject known to have an EBOV infection is administered 50 mg/kg of a disclosed antibody (or combination thereof) every day for 3 days following initial diagnosis of EBOV infection. In another example, the antibodies are administered continuously. Assessment

60

Following the administration of one or more therapies, subjects with EBOV can be monitored for a reduction in EBOV levels (such as viral titer or the EBOV GP level in serum), or reductions in one or more clinical symptoms associated with EBOV infection. Subjects can be monitored using any method known in the art. For example, biological samples from the subject, including blood, can be obtained 5 and alterations in EBOV levels evaluated. Additional Treatments

In particular examples, if subjects are stable or have a minor, mixed or partial response to treatment, they can be re-treated after re-evaluation with the same schedule and 10 preparation of agents that they previously received for the

SEQUENCE LISTING

80

desired amount of time, including the duration of a subject's lifetime. A partial response is a reduction, such as at least a 10%, at least 20%, at least 30%, at least 40%, at least 50%, or at least 70% in EBOV infection (e.g., as measured by EBOV GP level or viral titer in serum), EBOV replication, or combination thereof.

It will be apparent that the precise details of the methods or compositions described may be varied or modified without departing from the spirit of the described embodiments. We claim all such modifications and variations that fall within the scope and spirit of the claims below.

<160> NUMBER OF SEQ ID NOS: 66 <210> SEO TD NO 1 <211> LENGTH: 119 <212> TYPE · PRT <213> ORGANISM: homo sapiens <400> SEOUENCE: 1 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Ile Gln Pro Gly Gly 5 10 1 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ala Leu Arg Met Tyr 20 25 Asp Met His Trp Val Arg Gln Thr Ile Asp Lys Arg Leu Glu Trp Val 35 40 Ser Ala Val Gly Pro Ser Gly Asp Thr Tyr Tyr Ala Asp Ser Val Lys 55 Gly Arg Phe Ala Val Ser Arg Glu Asn Ala Lys Asn Ser Leu Ser Leu 65 70 75 80 Gln Met Asn Ser Leu Thr Ala Gly Asp Thr Ala Ile Tyr Tyr Cys Val 90 85 Arg Ser Asp Arg Gly Val Ala Gly Leu Phe Asp Ser Trp Gly Gln Gly 100 105 Ile Leu Val Thr Val Ser Ser 115 <210> SEQ ID NO 2 <211> LENGTH: 107 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 2 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly 10 1 5 15 Asp Arg Ile Thr Ile Thr Cys Arg Ala Ser Gln Ala Phe Asp Asn Tyr 25 20 Val Ala Trp Tyr Gln Gln Arg Pro Gly Lys Val Pro Lys Leu Leu Ile 40 45 Ser Ala Ala Ser Ala Leu His Ala Gly Val Pro Ser Arg Phe Ser Gly 50 55 60 Ser Gly Ser Gly Thr His Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro 65 70 75 80 Glu Asp Val Ala Thr Tyr Tyr Cys Gln Asn Tyr Asn Ser Ala Pro Leu 85 90 95 Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys 100 105

81

< Z I .	1> LH	ENGTH	H: 12	25											
<21: <21:	2> TY 3> OF	(PE : RGANI	PRT [SM:	homo	s sa]	piens	3								
<40	0> SI	EQUEN	ICE :	3											
Gln 1	Val	Gln	Leu	Gln 5	Glu	Ser	Gly	Pro	Gly 10	Leu	Val	Lys	Pro	Ser 15	Asp
Thr	Leu	Ser	Leu 20	Thr	Сүз	Thr	Val	Ser 25	Gly	Gly	Ser	Leu	Ser 30	Ser	Phe
Tyr	Trp	Ser 35	Trp	Ile	Arg	Gln	Pro 40	Pro	Gly	Lys	Gly	Leu 45	Glu	Trp	Ile
Gly	Tyr 50	Ile	Tyr	Tyr	Ser	Gly 55	Ser	Pro	Asn	Tyr	Ser 60	Pro	Ser	Leu	Glu
Ser 65	Arg	Val	Thr	Met	Ser 70	Val	Asp	Thr	Thr	Arg 75	Asn	Gln	Ile	Ser	Leu 80
ГЛа	Leu	Aap	Ser	Val 85	Thr	Ala	Ala	Asp	Thr 90	Ala	Val	Tyr	Tyr	Сув 95	Val
Arg	Ala	Ser	Arg 100	Ser	Tyr	Tyr	Trp	Gly 105	Ser	Tyr	Arg	Pro	Thr 110	Ala	Phe
Aap	Ser	Trp 115	Gly	Gln	Gly	Thr	Leu 120	Val	Thr	Val	Ser	Ser 125			
<21) <21; <21; <21;	0> SH 1> LH 2> TY 3> OH	EQ II ENGTH ZPE : RGANJ	) NO H: 10 PRT [SM:	4 05 homa	s saj	piens	3								
<40	)> SH	EQUEN	ICE :	4											
Ser 1	Tyr	Glu	Leu	Thr 5	Gln	Pro	Leu	Ser	Val 10	Ser	Val	Ser	Pro	Gly 15	Gln
Thr	Ala	Ile	Phe 20	Thr	Сүз	Ser	Gly	Asp 25	Asn	Leu	Gly	Asp	Lys 30	Tyr	Val
СЛа	Trp	Phe 35	Gln	Gln	Arg	Pro	Gly 40	Gln	Ser	Pro	Met	Leu 45	Leu	Ile	Tyr
Cys Gln	Trp Asp 50	Phe 35 Asn	Gln Lys	Gln Arg	Arg Pro	Pro Ser 55	Gly 40 Gly	Gln Ile	Ser Pro	Pro Glu	Met Arg 60	Leu 45 Phe	Leu Ser	Ile Gly	Ser
Cys Gln Asn 65	Trp Asp 50 Ser	Phe 35 Asn Gly	Gln Lys Asn	Gln Arg Thr	Arg Pro Ala 70	Pro Ser 55 Thr	Gly 40 Gly Leu	Gln Ile Thr	Ser Pro Ile	Pro Glu Ser 75	Met Arg 60 Gly	Leu 45 Phe Thr	Leu Ser Gln	Ile Gly Ser	Ser Thr 80
Cys Gln Asn 65 Asp	Trp Asp 50 Ser Glu	Phe 35 Asn Gly Ala	Gln Lys Asn Asp	Gln Arg Thr Tyr 85	Arg Pro Ala 70 Tyr	Pro Ser 55 Thr Cys	Gly 40 Gly Leu Gln	Gln Ile Thr Thr	Ser Pro Ile Trp 90	Pro Glu Ser 75 Asp	Met Arg 60 Gly Ser	Leu 45 Phe Thr Thr	Leu Ser Gln Val	Ile Gly Ser Val 95	Ser Thr 80 Phe
Cys Gln Asn 65 Asp Gly	Trp Asp 50 Ser Glu Gly	Phe 35 Asn Gly Ala Gly	Gln Lys Asn Asp Thr 100	Gln Arg Thr Tyr 85 Lys	Arg Pro Ala 70 Tyr Leu	Pro Ser 55 Thr Cys Thr	Gly 40 Gly Leu Gln Val	Gln Ile Thr Thr Leu 105	Ser Pro Ile Trp 90	Pro Glu Ser 75 Asp	Met Arg 60 Gly Ser	Leu 45 Phe Thr Thr	Leu Ser Gln Val	Ile Gly Ser Val 95	Tyr Ser Thr 80 Phe
Cys Gln Asn 65 Asp Gly <211 <211 <211 <211	Trp 50 Ser Glu Gly 52 SF 52 SP 52 SF 53 53 SF 53 53 50 55 55 55 55 55 55 55 55 55 55 55 55	Phe 35 Asn Gly Ala Gly Cly EQ II EQ II	Gln Lys Asn Asp Thr 100 C NO C NO C NO C SM:	Gln Arg Thr Tyr 85 Lys 523 homo	Arg Pro Ala 70 Tyr Leu Leu	Pro Ser 55 Thr Cys Thr Thr	Gly 40 Gly Leu Gln Val	Gln Ile Thr Thr Leu 105	Ser Pro Ile Trp 90	Pro Glu Ser 75 Asp	Met Arg 60 Gly Ser	Leu 45 Phe Thr Thr	Leu Ser Gln Val	Ile Gly Ser Val 95	Ser Thr 80 Phe
Cys Gln Asn 65 Gly <211 <211 <211 <211 <211	Trp 50 Ser Glu Gly 2> SE 2> TY 3> OF	Phe 35 Asn Gly Ala Gly Gly EQ II ENGTH (PE: CGANJ	Gln Lys Asn Asp Thr 100 C NO 1: 12 PRT (SM: UCE:	Gln Arg Thr Tyr 85 Lys 523 homo 5	Arg Pro Ala 70 Tyr Leu Sap	Pro Ser 55 Thr Cys Thr	Gly 40 Gly Leu Gln Val	Gln Ile Thr Thr Leu 105	Ser Pro Ile Trp 90	Pro Glu Ser 75 Asp	Met Arg 60 Gly Ser	Leu 45 Phe Thr Thr	Leu Ser Gln Val	Ile Gly Ser Val 95	Ser Thr 80 Phe
Cys Gln Asn 65 Asp Gly <211 <211 <211 <211 <400 Gln 1	Trp 50 Ser Glu Gly 0> SI 1> LH 2> TY 3> OF 0> SI 0> SI Val	Phe 35 Asn Gly Ala Gly EQ III (PE: SQAN) GQUEN Gln	Gln Lys Asn Asp Thr 100 C NO C NO C NO C SM: ISM: ISM: Leu	Gln Arg Thr Tyr 85 Lys 5 23 homo 5 Val 5	Arg Pro Ala 70 Tyr Leu Gln	Pro Ser 55 Thr Cys Thr Thr Ser	Gly 40 Gly Leu Gln Val	Gln Ile Thr Thr Leu 105	Ser Pro Ile Trp 90 Glu 10	Pro Glu Ser 75 Asp Val	Met Arg 60 Gly Ser Lys	Leu 45 Phe Thr Thr Lys	Leu Ser Gln Val	Ile Gly Ser Val 95	Ser Thr 80 Phe Ser
Cys Gln Asn 65 Asp Gly <211 <211 <211 <211 <211 <10 0 Gln 1 Ser	Trp 50 Ser Glu Gly 5) SF 5) SF 5) SF 5) SF 5) SF 50 5) SF 50 50 50 50 50 50 50 50 50 50 50 50 50	Phe 35 Asn Gly Ala Gly EQ II CQ II C	Gln Lys Asn Asp Thr 100 O NO O NO O NO O S S S S S S S S S S S S S S S S S S	Gln Arg Thr Tyr 85 Lys 5 23 homo 5 Val 5 Ser	Arg Pro Ala 70 Tyr Leu Gln Cys	Pro Ser 55 Thr Cys Thr Thr Ser Lys	Gly 40 Gly Leu Gln Val Gly Gly Thr	Gln Ile Thr Thr Leu 105 Ala Ser 25	Ser Pro Ile Trp 90 Glu 10 Gly	Pro Glu Ser 75 Asp Val Gly	Met Arg 60 Gly Ser Lys Thr	Leu 45 Phe Thr Thr Lys Leu	Leu Ser Gln Val Pro Ser 30	Ile Gly Ser Val 95 Gly 15 Asn	Tyr Ser Thr 80 Phe Ser Tyr
Cys Gln Asn 65 Gly <211 <211 <211 <211 <211 Ser Ala	Trp Asp 50 Ser Glu Gly D> SE 1> LH 2> TY 3> OF D> SE Val Val Ile	Phe 35 Asn Gly Ala Gly Gly Gly Gly CQUEN Gln Lys Ser 35	Gln Lys Asn Asp Thr 100 O NO H: 12 PRT (SM: Leu Val 20 Trp	Gln Arg Thr Tyr 85 Lys bomo 5 Val 5 Ser Val	Arg Pro Ala 70 Tyr Leu Gln Cys Arg	Pro Ser 55 Thr Cys Thr Thr Ser Lys Gln	Gly 40 Gly Leu Gln Val S Gly Thr Ala 40	Gln Ile Thr Leu 105 Ala Ser 25 Pro	Ser Pro Ile Trp 90 Glu 10 Gly Gly	Pro Glu Ser 75 Asp Val Gly Gln	Met Arg 60 Gly Ser Lys Thr Gly	Leu 45 Phe Thr Thr Lys Leu 45	Leu Ser Gln Val Pro Ser 30 Glu	Ile Gly Ser Val 95 Gly 15 Asn Trp	Tyr Ser Thr 80 Phe Ser Tyr Met

#### -continued

Gln Gly Arg Val Ala Ile Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr 70 75 65 80 Met Glu Leu Ser Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys 85 90 Ala Thr Met Gly Ser Ala Asp Thr Ser Phe Tyr Phe Tyr Met Asp Val 100 105 110 Trp Gly Lys Gly Thr Thr Val Thr Val Ser Ser 115 120 <210> SEQ ID NO 6 <211> LENGTH: 108 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 6 Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly 10 1 5 15 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser 20 25 30 Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu 35 40 45 Ile Tyr Gly Thr Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser 50 55 60 Gly Ser Ala Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu 65 70 75 80 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Ala Tyr Ser Pro 85 90 95 Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys 100 105 <210> SEQ ID NO 7 <211> LENGTH: 358 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEQUENCE: 7 gaggtgcagc tggtggagtc tggggggggt ttaattcagc cggggggggtc cctgagactc teetgtgeag eetetggatt egeceteaga atgtaegaea tgeaetgggt eegteagaea atagataaac gtctcgagtg ggtctcagct gtgggtcctt ctggtgacac ctactatgca gactccgtga agggccgatt cgccgtctcc agagagaatg ccaagaactc cttgtctctt cagatgaaca gcctgacagc cggggacacg gctatatact attgtgtaag gtctgaccga ggagtggetg geetttttga cagetgggge cagggaatee tggteaeegt etetteag <210> SEQ ID NO 8 <211> LENGTH: 322 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEQUENCE: 8 gacatecaga tgacceagte tecateatee etgtetgeat etgtgggaga eagaateace atcacttgcc gggcgagtca ggcctttgac aattatgtag cctggtatca acagagacca qqqaaqqttc ctaaqctcct qatctctqct qcatccqctt tqcacqcaqq qqtcccatct cgcttcagcg gcagtggctc tgggacacat ttcactctca ccatcagcag cctgcagcct qaaqatqttq caacttatta ctqtcaaaac tataacaqtq ccccqctcac tttcqqcqqa

84

60

120

180

240

300

358

60

120

180

240

85

-continued

gggaccaagg tggagatcaa ac 322 <210> SEQ ID NO 9 <211> LENGTH: 376 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEQUENCE: 9 60 caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcggatac cctgtccctc 120 acctgtactg tetetggtgg etceetcagt agtttetact ggagetggat eeggeageee ccagggaagg gactggagtg gattgggtat atctattaca gtgggagccc caactacagc 180 ccctccctcg agagtcgagt caccatgtca gtagacacga ccaggaacca gatctccctg 240 aagttggact ctgtgaccgc ggcggacacg gccgtgtatt actgtgtgag agcctcccga 300 agttactatt ggggggagtta tcgcccaacg gcttttgact cctgggggcca gggaaccctg 360 376 gtcaccgtct cctcag <210> SEQ ID NO 10 <211> LENGTH: 316 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEOUENCE: 10 tectatgage tgactcagee acteteagtg teegtgteee caggeeagae ageeatette 60 acctgctctg gagataattt gggggataag tatgtttgct ggtttcaaca gaggccaggc 120 cagteeceta tgetgeteat etateaagae aataagegge eetegggggat eeetgagega 180 ttetetgget ceaactetgg gaacacagee actetgaeta teagegggae ceagtetaea 240 gatgaggetg actattactg teagaegtgg gaeageaeeg tggtgttegg eggagggaee 300 aaactgaccg tcctgg 316 <210> SEQ ID NO 11 <211> LENGTH: 370 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEQUENCE: 11 60 caggtccagc tggtgcagtc tggggctgag gtgaagaagc ctgggtcctc ggtgaaagtc teetgeaaga ettetggagg cacceteage aactatgeta teagetgggt gegaeaggee 120 cctggacaag ggcttgagtg gatgggaggc accatteeta ccettggtat gtecaectae 180 gcaccgaact tccagggcag agtcgcgatt accgcggaca aatccacgag cacagcctac 240 atggagttga gtagtetgag gtetgaegae aeggeegttt attattgtge gaetatggge 300 agtgcggaca ctagtttcta cttctacatg gacgtctggg gcaaagggac cacggtcacc 360 370 qtctcctcaq <210> SEQ ID NO 12 <211> LENGTH: 325 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEQUENCE: 12 gaaattgtgt tgacgcagtc tccaggcacc ctgtctttgt ctccagggga gagagccacc 60 120 ctctcctgca gggccagtca gagtgttagt agcagctact tagcctggta ccagcagaaa

88

# 87

### -continued

cctggccagg ctcccagact cctcatctat ggtacatcca gcagggccac tggcatccca	180
gacaggttca gtggcagtgc gtctgggaca gacttcactc tcaccatcag cagactggag	240
cctgaagatt ttgcagtgta ttactgtcag cagtatgctt actcaccatt cactttcggc	300
cctgggacca cagtggatat caaac	325
<210> SEQ ID NO 13 <211> LENGTH: 676 <212> TYPE: PRT <213> ORGANISM: Ebola virus	
<400> SEQUENCE: 13	
Met Val Thr Ser Gly Ile Leu Gln Leu Pro Arg Glu Arg Phe Arg Lys 1 5 10 15	
Thr Ser Phe Val Trp Val Ile Ile Leu Phe His Lys Val Phe Pro 20 25 30	
Ile Pro Leu Gly Val Val His Asn Asn Thr Leu Gln Val Ser Asp Ile 35 40 45	
Asp Lys Leu Val Cys Arg Asp Lys Leu Ser Ser Thr Ser Gln Leu Lys 50 55 60	
Ser Val Gly Leu Asn Leu Glu Gly Asn Gly Val Ala Thr Asp Val Pro 65 70 75 80	
Thr Ala Thr Lys Arg Trp Gly Phe Arg Ala Gly Val Pro Pro Lys Val 85 90 95	
Val Asn Tyr Glu Ala Gly Glu Trp Ala Glu Asn Cys Tyr Asn Leu Asp 100 105 110	
Ile Lys Lys Ala Asp Gly Ser Glu Cys Leu Pro Glu Ala Pro Glu Gly 115 120 125	
Val Arg Gly Phe Pro Arg Cys Arg Tyr Val His Lys Val Ser Gly Thr 130 135 140	
Gly Pro Cys Pro Glu Gly Tyr Ala Phe His Lys Glu Gly Ala Phe Phe 145 150 155 160	
Leu Tyr Asp Arg Leu Ala Ser Thr Ile Ile Tyr Arg Ser Thr Thr Phe 165 170 175	
Ser Glu Gly Val Val Ala Phe Leu Ile Leu Pro Glu Thr Lys Lys Asp 180 185 190	
Phe Phe Gln Ser Pro Pro Leu His Glu Pro Ala Asn Met Thr Thr Asp 195 200 205	
Pro Ser Ser Tyr Tyr His Thr Val Thr Leu Asn Tyr Val Ala Asp Asn 210 215 220	
Phe Gly Thr Asn Met Thr Asn Phe Leu Phe Gln Val Asp His Leu Thr 225 230 235 240	
Tyr Val Gln Leu Glu Pro Arg Phe Thr Pro Gln Phe Leu Val Gln Leu 245 250 255	
Asn Glu Thr Ile Tyr Thr Asn Gly Arg Arg Ser Asn Thr Thr Gly Thr 260 265 270	
Leu Ile Trp Lys Val Asn Pro Thr Val Asp Thr Gly Val Gly Glu Trp 275 280 285	
Ala Phe Trp Glu Asn Lys Lys Asn Phe Thr Lys Thr Leu Ser Ser Glu	
Glu Leu Ser Val Ile Phe Val Pro Arg Ala Gln Asp Pro Gly Ser Asn	
305     310     315     320       Gln Lys Thr Lys Val Thr Pro Thr Ser Phe Ala Asn Asn Gln Thr Ser	
325 330 335	

Lys Asn His Glu Asp Leu Val Pro Glu Asp Pro Ala Ser Val Val Gln 340 345 350

#### -continued

Va	1 A)	rg	Asp 355	Leu	Gln	Arg	Glu	Asn 360	Thr	Val	Pro	Thr	Pro 365	Pro	Pro	Азр
Th	r Va 3'	al 70	Pro	Thr	Thr	Leu	Ile 375	Pro	Asp	Thr	Met	Glu 380	Glu	Gln	Thr	Thr
Se 38	r H: 5	is	Tyr	Glu	Pro	Pro 390	Asn	Ile	Ser	Arg	Asn 395	His	Gln	Glu	Arg	Asn 400
As	n Tł	hr	Ala	His	Pro 405	Glu	Thr	Leu	Ala	Asn 410	Asn	Pro	Pro	Asp	Asn 415	Thr
Th	r Pi	ro	Ser	Thr 420	Pro	Pro	Gln	Asp	Gly 425	Glu	Arg	Thr	Ser	Ser 430	His	Thr
Th	r Pi	ro	Ser 435	Pro	Arg	Pro	Val	Pro 440	Thr	Ser	Thr	Ile	His 445	Pro	Thr	Thr
Ar	g G 49	lu 50	Thr	His	Ile	Pro	Thr 455	Thr	Met	Thr	Thr	Ser 460	His	Aab	Thr	Азр
Se 46	r A: 5	sn	Arg	Pro	Asn	Pro 470	Ile	Asp	Ile	Ser	Glu 475	Ser	Thr	Glu	Pro	Gly 480
Pr	o Le	eu	Thr	Asn	Thr 485	Thr	Arg	Gly	Ala	Ala 490	Asn	Leu	Leu	Thr	Gly 495	Ser
Ar	g Ai	rg	Thr	Arg 500	Arg	Glu	Ile	Thr	Leu 505	Arg	Thr	Gln	Ala	Lys 510	Cys	Asn
Pr	o Af	sn	Leu 515	His	Tyr	Trp	Thr	Thr 520	Gln	Asp	Glu	Gly	Ala 525	Ala	Ile	Gly
Le	u A1	la 30	Trp	Ile	Pro	Tyr	Phe	Gly	Pro	Ala	Ala	Glu 540	Gly	Ile	Tyr	Thr
Gl	u GI	ly	Ile	Met	His	Asn	Gln	Asn	Gly	Leu	Ile	Суз	Gly	Leu	Arg	Gln
Le	u Al	la	Asn	Glu	Thr	Thr	Gln	Ala	Leu	Gln	Leu	Phe	Leu	Arg	Ala	Thr
Th	r G	lu	Leu	Arg	565 Thr	Phe	Ser	Ile	Leu	Asn	Arg	Lys	Ala	Ile	Asp	Phe
Le	u Le	eu	Gln	580 Arg	Trp	Gly	Gly	Thr	585 Суз	His	Ile	Leu	Gly	590 Pro	Asp	Суа
су	s II	le	595 Glu	Pro	His	Asp	Trp	600 Thr	Lys	Asn	Ile	Thr	605 Asp	Lys	Ile	Азр
Gl	6: n I:	10 le	Ile	His	Asp	Phe	615 Ile	Asp	Lys	Pro	Leu	620 Pro	Asp	Gln	Thr	Азр
62 As	5 n A:	.sp	Asn	Trp	Trp	630 Thr	Gly	Trp	Arg	Gln	635 Trp	Val	Pro	Ala	Gly	640 Ile
Gl	y II	le	Thr	- Gly	645 Val	Ile	Ile	- Ala	Val	650 Ile	- Ala	Leu	Leu	Cys	655 Ile	Суз
Lo	 s Pł	he	Leu	660 Leu		-			665	-				670	-	-
y			675	4												
<2 <2 <2 <2	10> 11> 12> 13>	SE LE TY OR	Q II NGTH PE : GANJ	) NO H: 6' PRT SM:	14 76 Ebo:	la v:	irus									
< 4	00>	SE	QUEN	ICE :	14											
Me 1	t G	lu	Gly	Leu	Ser 5	Leu	Leu	Gln	Leu	Pro 10	Arg	Asp	ГÀа	Phe	Arg 15	ГЛа
Se	r Se	er	Phe	Phe 20	Val	Trp	Val	Ile	Ile 25	Leu	Phe	Gln	Lys	Ala 30	Phe	Ser

#### -continued

Met Pro Leu Gly Val Val Thr Asn Ser Thr Leu Glu Val Thr Glu Ile Asp Gln Leu Val Cys Lys Asp His Leu Ala Ser Thr Asp Gln Leu Lys Ser Val Gly Leu Asn Leu Glu Gly Ser Gly Val Ser Thr Asp Ile Pro Ser Ala Thr Lys Arg Trp Gly Phe Arg Ser Gly Val Pro Pro Lys Val Phe Ser Tyr Glu Ala Gly Glu Trp Ala Glu Asn Cys Tyr Asn Leu Glu Ile Lys Lys Pro Asp Gly Ser Glu Cys Leu Pro Pro Pro Pro Asp Gly Val Arg Gly Phe Pro Arg Cys Arg Tyr Val His Lys Ala Gln Gly Thr Gly Pro Cys Pro Gly Asp Tyr Ala Phe His Lys Asp Gly Ala Phe Phe Leu Tyr Asp Arg Leu Ala Ser Thr Val Ile Tyr Arg Gly Val Asn Phe Ala Glu Gly Val Ile Ala Phe Leu Ile Leu Ala Lys Pro Lys Glu Thr Phe Leu Gln Ser Pro Pro Ile Arg Glu Ala Val Asn Tyr Thr Glu Asn Thr Ser Ser Tyr Tyr Ala Thr Ser Tyr Leu Glu Tyr Glu Ile Glu Asn Phe Gly Ala Gln His Ser Thr Thr Leu Phe Lys Ile Asn Asn Asn Thr Phe Val Leu Leu Asp Arg Pro His Thr Pro Gln Phe Leu Phe Gln Leu Asn Asp Thr Ile His Leu His Gln Gln Leu Ser Asn Thr Thr Gly Lys Leu Ile Trp Thr Leu Asp Ala Asn Ile Asn Ala Asp Ile Gly Glu Trp - 285 Ala Phe Trp Glu Asn Lys Lys Asn Leu Ser Glu Gln Leu Arg Gly Glu Glu Leu Ser Phe Glu Thr Leu Ser Leu Asn Glu Thr Glu Asp Asp Asp Ala Thr Ser Ser Arg Thr Thr Lys Gly Arg Ile Ser Asp Arg Ala Thr Arg Lys Tyr Ser Asp Leu Val Pro Lys Asp Ser Pro Gly Met Val Ser Leu His Val Pro Glu Gly Glu Thr Thr Leu Pro Ser Gln Asn Ser Thr Glu Gly Arg Arg Val Asp Val Asn Thr Gln Glu Thr Ile Thr Glu Thr Thr Ala Thr Ile Ile Gly Thr Asn Gly Asn Asn Met Gln Ile Ser Thr Ile Gly Thr Gly Leu Ser Ser Ser Gln Ile Leu Ser Ser Ser Pro Thr Met Ala Pro Ser Pro Glu Thr Gln Thr Ser Thr Thr Tyr Thr Pro Lys Leu Pro Val Met Thr Thr Glu Glu Ser Thr Thr Pro Pro Arg Asn Ser

### -continued

Pro	Gl <sub>3</sub> 450	Sei	r Th	ır	Thr	Glu	Ala 455	Pro	Thr	Leu	Thr	Thr 460	Pro	Glu	Asn	Ile
Thi 465	Thi	Ala	a Va	1	Lys	Thr 470	Val	Leu	Pro	Gln	Glu 475	Ser	Thr	Ser	Asn	Gly 480
Leu	. Il€	Th:	s Se	er	Thr 485	Val	Thr	Gly	Ile	Leu 490	Gly	Ser	Leu	Gly	Leu 495	Arg
Lys	Arg	Sei	r Ar 50	g	Arg	Gln	Val	Asn	Thr 505	Arg	Ala	Thr	Gly	Lys 510	Cys	Asn
Pro	Asr	ι Leι 515	ı Hi 5	s	Tyr	Trp	Thr	Ala 520	Gln	Glu	Gln	His	Asn 525	Ala	Ala	Gly
Ile	e Ala 530	Tr <u>r</u>	, Il	.e	Pro	Tyr	Phe 535	Gly	Pro	Gly	Ala	Glu 540	Gly	Ile	Tyr	Thr
Glu 545	ι Glչ	' Lei	ı Me	t	His	Asn 550	Gln	Asn	Ala	Leu	Val 555	Суз	Gly	Leu	Arg	Gln 560
Leu	ı Ala	. Asr	n Gl	u	Thr 565	Thr	Gln	Ala	Leu	Gln 570	Leu	Phe	Leu	Arg	Ala 575	Thr
Thi	Glu	. Lei	1 Ar 58	g	Thr	Tyr	Thr	Ile	Leu 585	Asn	Arg	ГЛа	Ala	Ile 590	Asp	Phe
Leu	. Lei	1 Arg 595	g Ar 5	g	Trp	Gly	Gly	Thr 600	Суз	Arg	Ile	Leu	Gly 605	Pro	Asp	Суз
Суз	11e 610	e Glu	ı Pr	0	His	Asp	Trp 615	Thr	Гла	Asn	Ile	Thr 620	Asp	Гла	Ile	Asn
Glr 625	ı Ile	e Ile	∍ Hi	s	Asp	Phe 630	Ile	Asp	Asn	Pro	Leu 635	Pro	Asn	Gln	Asp	Asn 640
Asr	) Ast	) Asr	ı Tr	р	Trp 645	Thr	Gly	Trp	Arg	Gln 650	Trp	Ile	Pro	Ala	Gly 655	Ile
Glγ	' Ile	Th:	6 Gl	.у 0	Ile	Ile	Ile	Ala	Ile 665	Ile	Ala	Leu	Leu	Cys 670	Val	Сүз
Lys	Leu	Leu 675	ı Cy 5	s												
<21 <21 <21 <21	.0> 9 .1> 1 .2> 7 .3> 0	EQ I ENGI YPE : RGAI	IDN TH: PR NISM	10 67 2T I :	15 76 Ebo]	La v:	irus									
<40	0> 5	EQUI	ENCE	:	15											
Met 1	Glγ	Val	L Th	ır	Gly 5	Ile	Leu	Gln	Leu	Pro 10	Arg	Asp	Arg	Phe	Lys 15	Lys
Thi	Se1	Phe	e Ph 20	le I	Leu	Trp	Val	Ile	Ile 25	Leu	Phe	Gln	Arg	Thr 30	Phe	Ser
Ile	e Pro	) Leu 35	ı Gl	·У	Val	Ile	His	Asn 40	Ser	Thr	Leu	Gln	Val 45	Ser	Asp	Val
Ast	ь Lys 50	Lei	ı Va	1	Суз	Arg	Asp 55	Lys	Leu	Ser	Ser	Thr 60	Asn	Gln	Leu	Arg
Se1 65	Va]	Gl	/ Le	u	Asn	Leu 70	Glu	Gly	Asn	Gly	Val 75	Ala	Thr	Asp	Val	Pro 80
Sei	Ala	. Thi	r Ly	s	Arg 85	Trp	Gly	Phe	Arg	Ser 90	Gly	Val	Pro	Pro	Lys 95	Val
Val	. Asr	. Туз	c Gl 10	.u 0	Ala	Gly	Glu	Trp	Ala 105	Glu	Asn	Суа	Tyr	Asn 110	Leu	Glu
Ile	e Lys	Lys 119	s Pr 5	0	Asp	Gly	Ser	Glu 120	Сүз	Leu	Pro	Ala	Ala 125	Pro	Asp	Gly
Ile	Arg	∣ Gl∑	7 Ph	ıe	Pro	Arg	Cys 135	Arg	Tyr	Val	His	Lys 140	Val	Ser	Gly	Thr

### -continued

Gly 145	Pro	Суз	Ala	Gly	Asp 150	Phe	Ala	Phe	His	Lys 155	Glu	Gly	Ala	Phe	Phe 160
Leu	Tyr	Asp	Arg	Leu 165	Ala	Ser	Thr	Val	Ile 170	Tyr	Arg	Gly	Thr	Thr 175	Phe
Ala	Glu	Gly	Val 180	Val	Ala	Phe	Leu	Ile 185	Leu	Pro	Gln	Ala	Lys 190	Lys	Asp
Phe	Phe	Ser 195	Ser	His	Pro	Leu	Arg 200	Glu	Pro	Val	Asn	Ala 205	Thr	Glu	Asp
Pro	Ser 210	Ser	Gly	Tyr	Tyr	Ser 215	Thr	Thr	Ile	Arg	Tyr 220	Gln	Ala	Thr	Gly
Phe 225	Gly	Thr	Asn	Glu	Thr 230	Glu	Tyr	Leu	Phe	Glu 235	Val	Asp	Asn	Leu	Thr 240
Tyr	Val	Gln	Leu	Glu 245	Ser	Arg	Phe	Thr	Pro 250	Gln	Phe	Leu	Leu	Gln 255	Leu
Asn	Glu	Thr	Ile 260	Tyr	Thr	Ser	Gly	Lys 265	Arg	Ser	Asn	Thr	Thr 270	Gly	ГÀа
Leu	Ile	Trp 275	Lys	Val	Asn	Pro	Glu 280	Ile	Asp	Thr	Thr	Ile 285	Gly	Glu	Trp
Ala	Phe 290	Trp	Glu	Thr	Lys	Lys 295	Asn	Leu	Thr	Arg	Lys 300	Ile	Arg	Ser	Glu
Glu 305	Leu	Ser	Phe	Thr	Ala 310	Val	Ser	Asn	Arg	Ala 315	Lys	Asn	Ile	Ser	Gly 320
Gln	Ser	Pro	Ala	Arg 325	Thr	Ser	Ser	Asp	Pro 330	Gly	Thr	Asn	Thr	Thr 335	Thr
Glu	Asp	His	Lys 340	Ile	Met	Ala	Ser	Glu 345	Asn	Ser	Ser	Ala	Met 350	Val	Gln
Val	His	Ser 355	Gln	Gly	Arg	Glu	Ala 360	Ala	Val	Ser	His	Leu 365	Thr	Thr	Leu
Ala	Thr 370	Ile	Ser	Thr	Ser	Pro 375	Gln	Pro	Pro	Thr	Thr 380	Lys	Pro	Gly	Pro
Asp 385	Asn	Ser	Thr	His	Asn 390	Thr	Pro	Val	Tyr	Lys 395	Leu	Asp	Ile	Ser	Glu 400
Ala	Thr	Gln	Ala	Glu 405	Gln	His	His	Arg	Arg 410	Thr	Asp	Asn	Asp	Ser 415	Thr
Thr	Ser	Aab	Thr 420	Pro	Pro	Ala	Met	Thr 425	Ala	Ala	Gly	Pro	Pro 430	Lys	Ala
Glu	Asn	Thr 435	Asn	Thr	Ser	LÀa	Gly 440	Thr	Asp	Leu	Pro	Asp 445	Pro	Ala	Thr
Thr	Thr 450	Ser	Pro	Gln	Asn	His 455	Ser	Glu	Thr	Ala	Gly 460	Asn	Asn	Asn	Thr
His 465	His	Gln	Asp	Thr	Gly 470	Glu	Glu	Ser	Ala	Ser 475	Ser	Gly	Lys	Leu	Gly 480
Leu	Ile	Thr	Asn	Thr 485	Ile	Ala	Gly	Val	Ala 490	Gly	Leu	Ile	Thr	Gly 495	Gly
Arg	Arg	Thr	Arg 500	Arg	Glu	Ala	Ile	Val 505	Asn	Ala	Gln	Pro	Lys 510	Cya	Asn
Pro	Asn	Leu 515	His	Tyr	Trp	Thr	Thr 520	Gln	Asp	Glu	Gly	Ala 525	Ala	Ile	Gly
Leu	Ala 530	Trp	Ile	Pro	Tyr	Phe 535	Gly	Pro	Ala	Ala	Glu 540	Gly	Ile	Tyr	Thr
Glu 545	Gly	Leu	Met	His	Asn 550	Gln	Asp	Gly	Leu	Ile 555	Суз	Gly	Leu	Arg	Gln 560

### -continued

Leu	Ala	Asn	Glu	Thr 565	Thr	Gln	Ala	Leu	Gln 570	Leu	Phe	Leu	Arg	Ala 575	Thr
Thr	Glu	Leu	Arg 580	Thr	Phe	Ser	Ile	Leu 585	Asn	Arg	Lys	Ala	Ile 590	Asp	Phe
Leu	Leu	Gln 595	Arg	Trp	Gly	Gly	Thr 600	Суз	His	Ile	Leu	Gly 605	Pro	Asp	Суз
Суз	Ile 610	Glu	Pro	His	Asp	Trp 615	Thr	Lys	Asn	Ile	Thr 620	Asp	Lys	Ile	Asp
Gln 625	Ile	Ile	His	Asp	Phe 630	Val	Asp	Lys	Thr	Leu 635	Pro	Asp	Gln	Gly	Asp 640
Asn	Asp	Asn	Trp	Trp 645	Thr	Gly	Trp	Arg	Gln 650	Trp	Ile	Pro	Ala	Gly 655	Ile
Gly	Val	Thr	Gly 660	Val	Ile	Ile	Ala	Val 665	Ile	Ala	Leu	Phe	Cys 670	Ile	Суз
Lys	Phe	Val 675	Phe												
<210 <211 <212	)> SE .> LE :> TY	EQ II ENGTH PE :	) NO 1: 6' PRT	16 77											
<213	> OF )> SE	igani Iquen	ISM: ICE:	Ebo:	la v:	irus									
Met 1	Gly	Ser	Gly	Tyr	Gln	Leu	Leu	Gln	Leu 10	Pro	Arg	Glu	Arg	Phe	Arg
rÀa	Thr	Ser	Phe	Leu	Val	Trp	Val	Ile	Ile	Leu	Phe	Gln	Arg	Ala	Ile
Ser	Met	Pro	20 Leu	Gly	Ile	Val	Thr	25 Asn	Ser	Thr	Leu	Lys	30 Ala	Thr	Glu
Ile	Asp	35 Gln	Leu	Val	Cys	Arg	40 Asp	Lys	Leu	Ser	Ser	45 Thr	Ser	Gln	Leu
Lys	50 Ser	Val	Gly	Leu	Asn	55 Leu	Glu	Gly	Asn	Gly	60 Ile	Ala	Thr	Asp	Val
65 Pro	Ser	Ala	Thr	Lys	70 Arg	Trp	Gly	Phe	Arg	75 Ser	Gly	Val	Pro	Pro	80 Lys
Val	Val	Ser	Tvr	85 Glu	Ala	Glv	Glu	Trp	90 Ala	Glu	Asn	Cvs	Tvr	95 Asn	Leu
Glu	TIA	Larg	100 Lvs	Ser	Agn	Glv	Ser	105 Glu	Cva	I.e11	Pro	Leu	110 Pro	Pro	Agn
oru	C	115	сл <u>-</u> -	DF	b	U LY	120	J.	-у» Т	Jeu Wal	110 114	125	110	- I U	сл тЪ
GTÀ	vai 130	arg	GIÀ	rne	PTO	агд 135	- -	Arg	1yr	vai	нія 140	- гүа	vai	GIN	<u>-</u> ,
Thr 145	Gly	Pro	Суз	Pro	Gly 150	Asp	Leu	Ala	Phe	His 155	LÀa	Asn	Gly	Ala	Phe 160
Phe	Leu	Tyr	Asp	Arg 165	Leu	Ala	Ser	Thr	Val 170	Ile	Tyr	Arg	Gly	Thr 175	Thr
Phe	Thr	Glu	Gly 180	Val	Val	Ala	Phe	Leu 185	Ile	Leu	Ser	Glu	Pro 190	Lys	Lys
His	Phe	Trp 195	Lys	Ala	Thr	Pro	Ala 200	His	Glu	Pro	Val	Asn 205	Thr	Thr	Asp
Asp	Ser 210	Thr	Ser	Tyr	Tyr	Met 215	Thr	Leu	Thr	Leu	Ser 220	Tyr	Glu	Met	Ser
Asn 225	Phe	Gly	Gly	Lys	Glu 230	Ser	Asn	Thr	Leu	Phe 235	Lys	Val	Asp	Asn	His 240
Thr	Tyr	Val	Gln	Leu 245	Asp	Arg	Pro	His	Thr	Pro	Gln	Phe	Leu	Val 255	Gln

### -continued

Leu	Asn	Glu	Thr 260	Leu	Arg	Arg	Asn	Asn 265	Arg	Leu	Ser	Asn	Ser 270	Thr	Gly
Arg	Leu	Thr 275	Trp	Thr	Leu	Asp	Pro 280	Lys	Ile	Glu	Pro	Asp 285	Val	Gly	Glu
Trp	Ala 290	Phe	Trp	Glu	Thr	Lys 295	Lys	Asn	Phe	Ser	Gln 300	Gln	Leu	His	Gly
Glu 305	Asn	Leu	His	Phe	Gln 310	Ile	Leu	Ser	Thr	His 315	Thr	Asn	Asn	Ser	Ser 320
Asp	Gln	Ser	Pro	Ala 325	Gly	Thr	Val	Gln	Gly 330	Lys	Ile	Ser	Tyr	His 335	Pro
Pro	Thr	Asn	Asn 340	Ser	Glu	Leu	Val	Pro 345	Thr	Asp	Ser	Pro	Pro 350	Val	Val
Ser	Val	Leu 355	Thr	Ala	Gly	Arg	Thr 360	Glu	Glu	Met	Ser	Thr 365	Gln	Gly	Leu
Thr	Asn 370	Gly	Glu	Thr	Ile	Thr 375	Gly	Phe	Thr	Ala	Asn 380	Pro	Met	Thr	Thr
Thr 385	Ile	Ala	Pro	Ser	Pro 390	Thr	Met	Thr	Ser	Glu 395	Val	Asp	Asn	Asn	Val 400
Pro	Ser	Glu	Gln	Pro 405	Asn	Asn	Thr	Ala	Ser 410	Ile	Glu	Asp	Ser	Pro 415	Pro
Ser	Ala	Ser	Asn 420	Glu	Thr	Ile	Asp	His 425	Ser	Glu	Met	Asn	Pro 430	Ile	Gln
Gly	Ser	Asn 435	Asn	Ser	Ala	Gln	Ser 440	Pro	Gln	Thr	Lys	Thr 445	Thr	Pro	Ala
Pro	Thr 450	Ala	Ser	Pro	Met	Thr 455	Gln	Asp	Pro	Gln	Glu 460	Thr	Ala	Asn	Ser
Ser 465	Lys	Leu	Gly	Thr	Ser 470	Pro	Gly	Ser	Ala	Ala 475	Glu	Pro	Ser	Gln	Pro 480
Gly	Phe	Thr	Ile	Asn 485	Thr	Val	Ser	L'As	Val 490	Ala	Asp	Ser	Leu	Ser 495	Pro
Thr	Arg	Lys	Gln 500	Lys	Arg	Ser	Val	Arg 505	Gln	Asn	Thr	Ala	Asn 510	Lys	Cys
Asn	Pro	Asp 515	Leu	His	Tyr	Trp	Thr 520	Ala	Val	Asp	Glu	G1y 525	Ala	Ala	Val
GTÀ	Leu 530	Ala	Trp	11e	Pro	Tyr 535	Pne	GIY	Pro	Ala	A1a 540	GIU	GIY	11e	lyr
11e 545	Giu	GIY	Val	Met	H1S 550	Asn	GIN	Asn	GIY	Leu 555	11e	CYS	GIY	Leu	Arg 560
GIN	Leu	Ala	Asn	GIU 565	Thr	Thr	GIN	Ala	Leu 570	GIN	Leu	Phe	Leu	Arg 575	AIA
Thr	Thr	Glu	Leu 580	Arg	Thr	Tyr	Ser	Leu 585	Leu	Asn	Arg	гал	Ala 590	11e	Aap
Рhe	Leu	Leu 595	GIN	Arg	Trp	стλ	600 617	Thr	сув	Arg	⊥le	Leu 605	ЧЦΥ	Pro	ser
СЛа	Cys 610	Ile	Glu	Pro	His	Asp 615	Trp	Thr	Lys	Asn	Ile 620	Thr	Asp	Glu	Ile
Asn 625	Gln	Ile	Lys	His	Asp 630	Phe	Ile	Asp	Asn	Pro 635	Leu	Pro	Asp	His	Gly 640
Asp	Asp	Leu	Asn	Leu 645	Trp	Thr	Gly	Trp	Arg 650	Gln	Trp	Ile	Pro	Ala 655	Gly
Ile	Gly	Ile	Ile 660	Gly	Val	Ile	Ile	Ala 665	Ile	Ile	Ala	Leu	Leu 670	Cys	Ile

-continued

Cys Lys Ile Leu Cys 675

<210	)> SH L> LH	EQ II ENGTH	) NO 1: 67	17 76											
<212	2 > TY	ZPE :	PRT												
<213	8 > OF	RGANI	[SM:	Ebol	la v:	irus									
<400	)> SI	EQUEN	ICE :	17											
Met 1	Gly	Ala	Ser	Gly 5	Ile	Leu	Gln	Leu	Pro 10	Arg	Glu	Arg	Phe	Arg 15	ГÀа
Thr	Ser	Phe	Phe 20	Val	Trp	Val	Ile	Ile 25	Leu	Phe	His	ГЛа	Val 30	Phe	Ser
Ile	Pro	Leu 35	Gly	Val	Val	His	Asn 40	Asn	Thr	Leu	Gln	Val 45	Ser	Asp	Ile
Asp	Lys 50	Phe	Val	Cya	Arg	Asp 55	Lys	Leu	Ser	Ser	Thr 60	Ser	Gln	Leu	Lys
Ser 65	Val	Gly	Leu	Asn	Leu 70	Glu	Gly	Asn	Gly	Val 75	Ala	Thr	Asp	Val	Pro 80
Thr	Ala	Thr	Lys	Arg 85	Trp	Gly	Phe	Arg	Ala 90	Gly	Val	Pro	Pro	Lys 95	Val
Val	Asn	Сув	Glu 100	Ala	Gly	Glu	Trp	Ala 105	Glu	Asn	Сув	Tyr	Asn 110	Leu	Ala
Ile	Lys	Lys 115	Val	Asp	Gly	Ser	Glu 120	Cys	Leu	Pro	Glu	Ala 125	Pro	Glu	Gly
Val	Arg 130	Asp	Phe	Pro	Arg	Cys 135	Arg	Tyr	Val	His	Lys 140	Val	Ser	Gly	Thr
Gly 145	Pro	Сув	Pro	Gly	Gly 150	Leu	Ala	Phe	His	Lys 155	Glu	Gly	Ala	Phe	Phe 160
Leu	Tyr	Asp	Arg	Leu 165	Ala	Ser	Thr	Ile	Ile 170	Tyr	Arg	Gly	Thr	Thr 175	Phe
Ala	Glu	Gly	Val 180	Ile	Ala	Phe	Leu	Ile 185	Leu	Pro	Lys	Ala	Arg 190	Lys	Asp
Phe	Phe	Gln 195	Ser	Pro	Pro	Leu	His 200	Glu	Pro	Ala	Asn	Met 205	Thr	Thr	Asp
Pro	Ser 210	Ser	Tyr	Tyr	His	Thr 215	Thr	Thr	Ile	Asn	Tyr 220	Val	Val	Asp	Asn
Phe 225	Gly	Thr	Asn	Thr	Thr 230	Glu	Phe	Leu	Phe	Gln 235	Val	Asp	His	Leu	Thr 240
Tyr	Val	Gln	Leu	Glu 245	Ala	Arg	Phe	Thr	Pro 250	Gln	Phe	Leu	Val	Leu 255	Leu
Asn	Glu	Thr	Ile 260	Tyr	Ser	Asp	Asn	Arg 265	Arg	Ser	Asn	Thr	Thr 270	Gly	ГÀа
Leu	Ile	Trp 275	Lys	Ile	Asn	Pro	Thr 280	Val	Asp	Thr	Ser	Met 285	Gly	Glu	Trp
Ala	Phe 290	Trp	Glu	Asn	Lys	Lys 295	Asn	Phe	Thr	Lys	Thr 300	Leu	Ser	Ser	Glu
Glu 305	Leu	Ser	Phe	Val	Pro 310	Val	Pro	Glu	Thr	Gln 315	Asn	Gln	Val	Leu	Asp 320
Thr	Thr	Ala	Thr	Val 325	Ser	Pro	Pro	Ile	Ser 330	Ala	His	Asn	His	Ala 335	Ala
Glu	Asp	His	Lys 340	Glu	Leu	Val	Ser	Glu 345	Aap	Ser	Thr	Pro	Val 350	Val	Gln
Met	Gln	Asn 355	Ile	Lys	Gly	Lys	Asp 360	Thr	Met	Pro	Thr	Thr 365	Val	Thr	Gly

Val Pro Thr Thr Thr Pro Ser Pro Phe Pro Ile Asn Ala Arg Asn Thr

Asp His Thr Lys Ser Phe Ile Gly Leu Glu Gly Pro Gln Glu Asp His

#### -continued

Ser Thr Thr Gln Pro Ala Lys Thr Thr Ser Gln Pro Thr Asn Ser Thr Glu Ser Thr Thr Leu Asn Pro Thr Ser Glu Pro Ser Ser Arg Gly Thr Gly Pro Ser Ser Pro Thr Val Pro Asn Thr Thr Glu Ser His Ala Glu Leu Gly Lys Thr Thr Pro Thr Thr Leu Pro Glu Gln His Thr Ala Ala Ser Ala Ile Pro Arg Ala Val His Pro Asp Glu Leu Ser Gly Pro Gly Phe Leu Thr Asn Thr Ile Arg Gly Val Thr Asn Leu Leu Thr Gly Ser Arg Arg Lys Arg Arg Asp Val Thr Pro Asn Thr Gln Pro Lys Cys Asn Pro Asn Leu His Tyr Trp Thr Ala Leu Asp Glu Gly Ala Ala Ile Gly Leu Ala Trp Ile Pro Tyr Phe Gly Pro Ala Ala Glu Gly Ile Tyr Thr Glu Gly Ile Met Glu Asn Gln Asn Gly Leu Ile Cys Gly Leu Arg Gln Leu Ala Asn Glu Thr Thr Gln Ala Leu Gln Leu Phe Leu Arg Ala Thr Thr Glu Leu Arg Thr Phe Ser Ile Leu Asn Arg Lys Ala Ile Asp Phe Leu Leu Gln Arg Trp Gly Gly Thr Cys His Ile Leu Gly Pro Asp Cys Cys Ile Glu Pro Gln Asp Trp Thr Lys Asn Ile Thr Asp Lys Ile Asp Gln Ile Ile His Asp Phe Val Asp Asn Asn Leu Pro Asn Gln Asn Asp Gly Ser Asn Trp Trp Thr Gly Trp Lys Gln Trp Val Pro Ala Gly Ile Gly Ile Thr Gly Val Ile Ile Ala Ile Ile Ala Leu Leu Cys Ile Cys Lys Phe Met Leu <210> SEQ ID NO 18 <211> LENGTH: 364 <212> TYPE: PRT <213> ORGANISM: Ebola virus <400> SEQUENCE: 18 Met Gly Val Thr Gly Ile Leu Gln Leu Pro Arg Asp Arg Phe Lys Arg Thr Ser Phe Phe Leu Trp Val Ile Ile Leu Phe Gln Arg Thr Phe Ser Ile Pro Leu Gly Val Ile His Asn Ser Thr Leu Gln Val Ser Asp Val Asp Lys Leu Val Cys Arg Asp Lys Leu Ser Ser Thr Asn Gln Leu Arg

Ser Val Gly Leu Asn Leu Glu Gly Asn Gly Val Ala Thr Asp Val Pro

Ser Ala Thr Lys Arg Trp Gly Phe Arg Ser Gly Val Pro Pro Lys Val

Val Asn Tyr Glu Ala Gly Glu Trp Ala Glu Asn Cys Tyr Asn Leu Glu

Ile Lys Lys Pro Asp Gly Ser Glu Cys Leu Pro Ala Ala Pro Asp Gly

-continued

Ile Arg Gly Phe Pro Arg Cys Arg Tyr Val His Lys Val Ser Gly Thr Gly Pro Cys Ala Gly Asp Phe Ala Phe His Lys Glu Gly Ala Phe Phe Leu Tyr Asp Arg Leu Ala Ser Thr Val Ile Tyr Arg Gly Thr Thr Phe Ala Glu Gly Val Val Ala Phe Leu Ile Leu Pro Gln Ala Lys Lys Asp Phe Phe Ser Ser His Pro Leu Arg Glu Pro Val Asn Ala Thr Glu Asp Pro Ser Ser Gly Tyr Tyr Ser Thr Thr Ile Arg Tyr Gln Ala Thr Gly Phe Gly Thr Asn Glu Thr Glu Tyr Leu Phe Glu Val Asp Asn Leu Thr Tyr Val Gln Leu Glu Ser Arg Phe Thr Pro Gln Phe Leu Leu Gln Leu Asn Glu Thr Ile Tyr Thr Ser Gly Lys Arg Ser Asn Thr Thr Gly Lys Leu Ile Trp Lys Val Asn Pro Glu Ile Asp Thr Thr Ile Gly Glu Trp Ala Phe Trp Glu Thr Lys Lys Thr Ser Leu Glu Lys Phe Ala Val Lys Ser Cys Leu Ser Gln Leu Tyr Gln Thr Glu Pro Lys Thr Ser Val Val Arg Val Arg Arg Glu Leu Leu Pro Thr Gln Gly Pro Thr Gln Gln Leu Lys Thr Thr Lys Ser Trp Leu Gln Lys Ile Pro Leu Gln Trp Phe Lys Cys Thr Val Lys Glu Gly Lys Leu Gln Cys Arg Ile <210> SEQ ID NO 19 <211> LENGTH: 128 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 19 Asp Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Gly Ser Leu Lys Leu Ala Cys Val Val Ser Gly Phe Arg Phe Ser Asp Tyr Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Ala Asn Ile Lys Gln Asp Gly Ser Gly Lys Tyr Tyr Val Asp Ser Val 

#### -continued

Lys Gly Arg Phe Thr Val Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr 65 70 75 80 Leu His Met Thr Ser Leu Gly Ala Glu Asp Thr Ala Val Tyr Phe Cys 85 90 95 Ala Arg Ala Ala Pro Thr Gly Ser Tyr Thr Asn Ile Leu Val Asp Asn 105 100 110 Val His Phe Asp Tyr Trp Gly Gln Gly Ile Leu Val Ala Val Ser Ser 115 120 125 <210> SEQ ID NO 20 <211> LENGTH: 107 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 20 Gly Ile Gln Leu Thr Gln Ser Pro Gly Ser Leu Ser Ala Ser Val Gly 10 5 15 1 Asp Ser Val Thr Ile Thr Cys Arg Pro Asn Gln Asn Ile Ala Thr Tyr 25 30 20 Ile Asn Trp Tyr Gln Gln Thr Pro Gly Lys Ala Pro Lys Leu Leu Ile 40 35 45 Tyr Ala Ala Ser Ile Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly 50 55 60 Ala Gly Ser Gly Thr His Phe Thr Leu Ile Ile Ser Thr Leu Gln Pro 65 70 75 80 Glu Asp Ser Ala Thr Tyr Tyr Cys Gln Gln Ser Tyr Ser Thr Pro Trp 85 90 95 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys 100 105 <210> SEO ID NO 21 <211> LENGTH: 385 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEQUENCE: 21 gatgtgcagt tggtggagtc tggggggggc gtggtccagc cgggggggtc cctgaaactc 60 gcctgtgtag tctctggatt caggtttagt gactactgga tgagttgggt ccgccaggcc 120 ccagggaagg ggctggaatg ggtggccaac ataaaacaag atggaagtgg gaagtactat 180 gtggactccg tgaagggccg attcaccgtc tccagagaca acgccaagaa ctcactgtat 240 ctacacatga ccagcctggg agccgaggac acggccgtat acttctgcgc gagagcagcc 300 cccaccggct cctacactaa tatcctagtc gacaacgtcc acttcgacta ctggggccag 360 385 qqaatcctqq tcqccqtctc ctcaq <210> SEQ ID NO 22 <211> LENGTH: 322 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEQUENCE: 22 ggcatccagc tgacccagtc tccaggctcc ctgtctgcat ctgtaggaga cagtgtcacc 60 atcacttgcc ggccaaatca gaacatcgcc acctatataa attggtatca gcagacacca 120 gggaaagccc ctaagctcct gatctatgcc gcatccattt tgcagagtgg ggtcccatca 180 aggttcagtg gcgctggatc tgggacacat ttcactctca tcatcagtac cctacaacct 240

110

## 109

### -continued

-continued	
gaggattetg caaettaeta etgecaaeag agttaeagta eeeegggae atteggeeaa	300
gggaccaaag tggaaatcaa ac	322
<210> SEQ ID NO 23 <211> LENGTH: 113 <212> TYPE: PRT <213> ORGANISM: homo sapiens	
<400> SEQUENCE: 23	
Ala Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Thr 1 5 10 15	
Thr Val Lys Ile Ser Cys Lys Val Ser Gly Tyr Thr Phe Ile Gln Glu 20 25 30	
Tyr Ile His Trp Val Gln Gln Ala Pro Gly Lys Gly Leu Val Trp Met 35 40 45	
Gly Leu Gly Asp Pro Glu Asn Asn Glu Thr Leu Tyr Ser Glu Asp Phe 50 55 60	
Gln Gly Arg Val Thr Met Thr Ala Asp Thr Ser Ser Asp Thr Ala Tyr 65 70 75 80	
Leu Glu Leu Arg Ser Leu Thr Phe Ala Asp Thr Ala Val Tyr Phe Cys 85 90 95	
Thr Ser Arg Lys Ser Trp Trp Gly Gln Gly Thr Leu Val Thr Val Ala 100 105 110	
Ser	
<210> SEQ ID NO 24 <211> LENGTH: 111 <212> TYPE: PRT <213> ORGANISM: homo sapiens	
<400> SEQUENCE: 24	
Glu Leu Val Leu Thr Gln Ser Pro GlyThr Leu Ser Leu Ser Pro Gly151015	
Glu Ser Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Leu Ser Ser Asp 20 25 30	
Ser Val Ser Trp Phe Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Val 35 40 45	
Ile His Gly Thr Ser Lys Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser 50 55 60	
Gly Gly Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ala Arg Leu Glu 65 70 75 80	
Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Arg Ser Gly Tyr Gly Met 85 90 95	
Ser Val Thr Trp Thr Phe Gly Gln Gly Thr Thr Val Glu Ile Lys 100 105 110	
<210> SEQ ID NO 25 <211> LENGTH: 340 <212> TYPE: DNA <213> ORGANISM: homo sapiens	
<400> SEQUENCE: 25	
gcggtccagt tggtacaatc tggggctgag gtgaagaagc ctgggaccac cgtcaaaatc	60
teetgeaaag tttetggata caeetteatt caagaataca taeaetgggt geaacaggee	120
cctggaaaag ggcttgtgtg gatgggactt ggtgaccctg aaaataatga gactctatat	180
tcagaggatt tccaaggcag agtcaccatg accgcggaca catcctcaga cacagcctat	240

1	1	1	
	н		

-continued	
------------	--

collected logication acceleration acceleration acceleration	300
teetggtggg geeagggaae eetggteaee gtegeeteag	340
<210> SEQ ID NO 26 <211> LENGTH: 334 <212> TYPE: DNA <213> ORGANISM: homo sapiens	
<400> SEQUENCE: 26	
gaacttgtgt tgacgcagtc tccaggcacc ctgtctttgt ctccagggga aagegccace	60
ctctcctgta gggccagtca gagtcttagc agcgactctg tatcttggtt ccagcagaaa	120
cctggccagg ctcccaggct cgtcatccat ggtacatcaa agagggccac tggcatccca	180
gacaggttca gtggcggtgg gtctgggaca gacttcactc tcaccatcgc cagactggag	240
cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg	300
acgttcggcc aagggaccac ggtggagatc aaac	334
<210> SEQ ID NO 27 <211> LENGTH: 334 <212> TYPE: DNA <213> ORGANISM: homo sapiens	
<400> SEQUENCE: 27	
gaacttgtgt tgacgcagtc tccaggcacc ctgtctttgt ctccagggga aagcgccacc	60
ctctcctgta gggccagtca gagtcttagc agcgactctg tatcttggtt ccagcagaaa	120
cctggccagg ctcccaggct cgtcatccat ggtacatcaa agagggccac tggcatccca	180
gacaggttca gtggcggtgg gtctgggaca gacttcactc tcaccatcgc cagactggag	240
cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg	300
cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg acgtttggcc aagggaccac ggtggagatc aaac	300 334
<pre>cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg acgtttggcc aagggaccac ggtggagatc aaac &lt;210&gt; SEQ ID NO 28 &lt;211&gt; LENGTH: 119 &lt;212&gt; TYPE: PRT &lt;213&gt; ORGANISM: homo sapiens &lt;400&gt; SEQUENCE: 28</pre>	300 334
<pre>cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg acgtttggcc aagggaccac ggtggagatc aaac &lt;210&gt; SEQ ID NO 28 &lt;211&gt; LENGTH: 119 &lt;212&gt; TYPE: PRT &lt;213&gt; ORGANISM: homo sapiens &lt;400&gt; SEQUENCE: 28 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Ile Gln Pro Gly Gly 1 5 10 15</pre>	300 334
cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg acgtttggcc aagggaccac ggtggagatc aaac <210> SEQ ID NO 28 <211> LENGTH: 119 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 28 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Ile Gln Pro Gly Gly 10 15 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ala Leu Arg Ser Tyr 20 25 30	300 334
<pre>cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg acgtttggcc aagggaccac ggtggagatc aaac &lt;210&gt; SEQ ID NO 28 &lt;211&gt; LENGTH: 119 &lt;212&gt; TYPE: PRT &lt;213&gt; ORGANISM: homo sapiens &lt;400&gt; SEQUENCE: 28 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Ile Gln Pro Gly Gly 15 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ala Leu Arg Ser Tyr 20 Asp Met His Trp Val Arg Gln Thr Ile Asp Lys Arg Leu Glu Trp Val 40 35</pre>	300 334
<pre>cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg acgtttggcc aagggaccac ggtggagatc aaac </pre>	300 334
cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtagtc agtcacgtgg acgtttggcc aagggaccac ggtggagatc aaac 2210> SEQ ID NO 28 2212> TYPE: PRT 212> ORGANISM: homo sapiens 2400> SEQUENCE: 28 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Ile Gln Pro Gly Gly Gly 10 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ala Leu Arg Ser Tyr 20 Asp Met His Trp Val Arg Gln Thr Ile Asp Lys Arg Leu Glu Trp Val 35 Ser Ala Val Gly Pro Ser Gly Asp Thr Tyr Tyr Ala Asp Ser Val Lys 60 Cly Arg Phe Ala Val Ser Arg Glu Asn Ala Lys Asn Ser Leu Ser Leu 80	300 334
<pre>cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg acgtttggcc aagggaccac ggtggagatc aaac &lt;210&gt; SEQ ID NO 28 &lt;211&gt; LENGTH: 119 &lt;212&gt; TYPE: PRT &lt;213&gt; ORGANISM: homo sapiens &lt;400&gt; SEQUENCE: 28 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Ile Gln Pro Gly Gly Gly 10 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ala Leu Arg Ser Tyr 20 Asp Met His Trp Val Arg Gln Thr Ile Asp Lys Arg Leu Glu Trp Val 35 Ser Ala Val Gly Pro Ser Gly Asp Thr Tyr Tyr Ala Asp Ser Val Lys 50 Gly Arg Phe Ala Val Ser Arg Glu Asn Ala Lys Asn Ser Leu Ser Leu 80 Gln Met Asn Ser Leu Thr Ala Gly Asp Thr Ala Ile Tyr Tyr Cys Val 90</pre>	300 334
cctgaggatt ttgcagtcta ttattgtcag cggtctgggt atggtatgtc agtcacgtgg acgtttggcc aagggaccac ggtggagatc aaac <210> SEQ ID NO 28 211> LENGTH: 119 212> TYPE: PRT 213> ORGANISM: homo sapiens <400> SEQUENCE: 28 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Ile Gln Pro Gly Gly Gly 10 10 10 10 10 15 15 170 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ala Leu Arg Ser Tyr 20 20 20 20 10 10 10 10 10 10 10 10 10 10 10 10 Asp Met His Trp Val Arg Gln Thr Ile Asp Lys Arg Leu Glu Trp Val 45 Ser Ala Val Gly Pro Ser Gly Asp Thr Tyr Tyr Ala Asp Ser Val Lys 50 Gly Arg Phe Ala Val Ser Arg Glu Asn Ala Lys Asn Ser Leu Ser Leu 80 Gln Met Asn Ser Leu Thr Ala Gly Asp Thr Ala Ile Tyr Tyr Gys Val 95 Arg Ser Asp Arg Gly Val Ala Gly Leu Phe Asp Ser Trp Gly Gln Gly 100	300 334

114

113

-continued

<211> LENGTH: 106 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 29 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly 10 1 5 15 Asp Arg Ile Thr Ile Thr Cys Arg Ala Ser Gln Ala Phe Ser Asn Tyr 25 20 30 Val Ala Trp Tyr Gln Gln Arg Pro Gly Lys Val Pro Lys Leu Leu Ile 40 35 Ser Ala Ala Ser Ala Leu His Ala Gly Val Pro Ser Arg Phe Ser Gly 55 Ser Gly Ser Gly Thr His Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro 65 70 75 80 Glu Asp Val Ala Thr Tyr Tyr Cys Gln Asn Tyr Asn Ser Ala Pro Leu 85 90 95 Thr Phe Gly Gly Gly Thr Lys Val Glu Ile 100 105 <210> SEQ ID NO 30 <211> LENGTH: 357 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEQUENCE: 30 gaagtgcagc tggtggagtc tggaggaggt ctgattcagc ccgggggttc cctgcgtctg 60 agttgtgccg catctggatt tgctctgcga agctacgaca tgcactgggt gagacagact 120 atcgataagc gcctggagtg ggtgtctgct gtcggcccca gtggagacac ctactatgca 180 gattcagtga aggggaggtt cgcagtctcc cgggaaaacg ccaaaaattc cctgagcctg 240 cagatgaact ctctgaccgc cggcgacaca gctatctact attgcgtcag gagcgataga 300 ggggtcgcag gactgtttga ttcatggggt cagggtattc tggtcaccgt gtcttca 357 <210> SEQ ID NO 31 <211> LENGTH: 318 <212> TYPE: DNA <213> ORGANISM: homo sapiens <400> SEQUENCE: 31 gatattcaga tgactcagag cccttcctca ctgtccgcat ccgtgggaga ccgtattact 60 attacttgta gagettetea ggetttttet aactaegtgg ettggtatea geagaggeee 120 ggcaaggtcc ctaaactgct gatctccgcc gcttctgcac tgcatgctgg agtgccaagc 180 240 cqqttctctq qaaqtqqatc aqqqactcac ttcaccctqa caatttccaq cctqcaqccc gaggatgtcg caacctacta ttgccagaac tacaacagtg ctcccctgac attcggtggt 300 318 ggaacaaagg tcgagatc <210> SEQ ID NO 32 <211> LENGTH: 8 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 32 Gly Phe Ala Leu Arg Met Tyr Asp 1 5

115

-continued

116

<210> SEQ ID NO 33 <211> LENGTH: 7 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 33 Val Gly Pro Ser Gly Asp Thr 1 5 <210> SEQ ID NO 34 <211> LENGTH: 13 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 34 Val Arg Ser Asp Arg Gly Val Ala Gly Leu Phe Asp Ser - 5 10 1 <210> SEQ ID NO 35 <211> LENGTH: 6 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 35 Gln Ala Phe Asp Asn Tyr 1 5 <210> SEQ ID NO 36 <211> LENGTH: 3 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 36 Ala Ala Ser 1 <210> SEQ ID NO 37 <211> LENGTH: 9 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 37 Gln Asn Tyr Asn Ser Ala Pro Leu Thr 1 5 <210> SEQ ID NO 38 <211> LENGTH: 8 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 38 Gly Phe Ala Leu Arg Ser Tyr Asp 1 5 <210> SEQ ID NO 39 <211> LENGTH: 6 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 39 Gln Ala Phe Ser Asn Tyr 1 5 <210> SEQ ID NO 40 <211> LENGTH: 8

1	1	7
Т	T	1

118

	-continued
<212> TYPE: PRT <213> ORGANISM:	homo sapiens
<400> SEQUENCE:	40
Gly Gly Ser Leu 1	Ser Ser Phe Tyr 5
<210> SEQ ID NO <211> LENGTH: 7 <212> TYPE: PRT <213> ORGANISM:	41 homo sapiens
<400> SEQUENCE:	41
Ile Tyr Tyr Ser 1	Gly Ser Pro 5
<210> SEQ ID NO <211> LENGTH: 1 <212> TYPE: PRT <213> ORGANISM:	42 9 homo sapiens
<400> SEQUENCE:	42
Val Arg Ala Ser 1	Arg Ser Tyr Tyr Trp Gly Ser Tyr Arg Pro Thr Ala 5 10 15
Phe Asp Ser	
<210> SEQ ID NO <211> LENGTH: 6 <212> TYPE: PRT <213> ORGANISM:	43 homo sapiens
<400> SEQUENCE:	43
Asn Leu Gly Asp 1	Lys Tyr 5
<210> SEQ ID NO <211> LENGTH: 3 <212> TYPE: PRT <213> ORGANISM:	44 homo sapiens
<400> SEQUENCE:	44
Gln Asp Asn 1	
<210> SEQ ID NO <211> LENGTH: 8 <212> TYPE: PRT <213> ORGANISM:	45 homo sapiens
<400> SEQUENCE:	45
Gln Thr Trp Asp 1	Ser Thr Val Val 5
<210> SEQ ID NO <211> LENGTH: 8 <212> TYPE: PRT <213> ORGANISM:	46 homo sapiens
<400> SEQUENCE:	46
Gly Phe Arg Phe 1	Ser Asp Tyr Trp 5
<210> SEQ ID NO <211> LENGTH: 8	47

-continued

1	1	Ω.
		u.
л	л.	1

-continued	

<212> TYPE: PRT <213> ORGANISM:	homo sapiens
<400> SEQUENCE:	47
Ile Lys Gln Asp 1	Gly Ser Gly Lys 5
<210> SEQ ID NO <211> LENGTH: 2: <212> TYPE: PRT <213> ORGANISM:	48 1 homo sapiens
<400> SEOUENCE:	48
~ Ala Arg Ala Ala 1	Pro Thr Gly Ser Tyr Thr Asn Ile Leu Val Asp Asn 5 10 15
Val His Phe Asp 20	Tyr
<210> SEQ ID NO <211> LENGTH: 6 <212> TYPE: PRT <213> ORGANISM:	49 homo sapiens
<400> SEQUENCE:	49
Gln Asn Ile Ala 1	Thr Tyr 5
<210> SEQ ID NO <211> LENGTH: 9 <212> TYPE: PRT <213> ORGANISM:	50 homo sapiens
<400> SEQUENCE:	50
Gln Gln Ser Tyr 1	Ser Thr Pro Trp Thr 5
<210> SEQ ID NO <211> LENGTH: 8 <212> TYPE: PRT	51
<213> ORGANISM:	homo sapiens
<400> SEQUENCE:	51
Gly Gly Thr Leu 1	Ser Asn Tyr Ala 5
<210> SEQ ID NO <211> LENGTH: 8 <212> TYPE: PRT <213> ORGANISM:	52 homo sapiens
<400> SEQUENCE:	52
Thr Ile Pro Thr 1	Leu Gly Met Ser 5
<210> SEQ ID NO <211> LENGTH: 1 <212> TYPE: PRT <213> ORGANISM:	53 5 homo sapiens
<400> SEQUENCE:	53
Ala Thr Met Gly 1	Ser Ala Asp Thr Ser Phe Tyr Phe Tyr Met Asp Val 5 10 15

121

-continued	

	-continued
<pre>&lt;211&gt; LENGTH: 7 &lt;212&gt; TYPE: PRT &lt;213&gt; ORGANISM:</pre>	homo sapiens
<400> SEQUENCE:	54
Gln Ser Val Ser 1	Ser Ser Tyr 5
<210> SEQ ID NO <211> LENGTH: 3 <212> TYPE: PRT <213> ORGANISM:	55 homo sapiens
<400> SEQUENCE:	55
Gly Thr Ser 1	
<210> SEQ ID NO <211> LENGTH: 9 <212> TYPE: PRT <213> ORGANISM:	56 homo sapiens
<400> SEQUENCE:	56
Gln Gln Tyr Ala 1	Tyr Ser Pro Phe Thr 5
<210> SEQ ID NO <211> LENGTH: 8 <212> TYPE: PRT <213> ORGANISM:	57 homo sapiens
<400> SEQUENCE:	57
Gly Tyr Thr Phe 1	Ile Gln Glu Tyr 5
<210> SEQ ID NO <211> LENGTH: 8 <212> TYPE: PRT <213> ORGANISM:	58 homo sapiens
<400> SEQUENCE:	58
Gly Asp Pro Glu 1	Asn Asn Glu Thr 5
<210> SEQ ID NO <211> LENGTH: 6 <212> TYPE: PRT <213> ORGANISM:	59 homo sapiens
<400> SEQUENCE:	59
Thr Ser Arg Lys 1	Ser Trp 5
<210> SEQ ID NO <211> LENGTH: 7 <212> TYPE: PRT <213> ORGANISM:	60 homo sapiens
<400> SEQUENCE:	60
Gln Ser Leu Ser 1	Ser Asp Ser 5
<210> SEQ ID NO <211> LENGTH: 12 <212> TYPE: PRT	61 2

-continued

<213> ORGANISM: homo sapiens <400> SEQUENCE: 61 Gln Arg Ser Gly Tyr Gly Met Ser Val Thr Trp Thr 5 10 1 <210> SEQ ID NO 62 <211> LENGTH: 108 <212> TYPE: PRT <213> ORGANISM: homo sapiens <400> SEQUENCE: 62 Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Leu Ser Pro Gly 1 5 10 15 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Ser Ser Ser 25 20 30 Tyr Leu Ala Trp Tyr Gl<br/>n Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu 35 40 45 Ile Tyr Gly Thr Ser Ser Arg Ala Thr Gly Ile Pro Asp Arg Phe Ser505560 Gly Ser Ala Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu 65 75 70 80 Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr Ala Tyr Ser Pro 85 90 95 Phe Thr Phe Gly Pro Gly Thr Thr Val Asp Ile Lys 100 105 <210> SEQ ID NO 63 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: DNA primer <400> SEQUENCE: 63 ttttcaatcc tcaaccgtaa ggc 23 <210> SEQ ID NO 64 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: DNA primer <400> SEQUENCE: 64 cagtccggtc ccagaatgtg 20 <210> SEQ ID NO 65 <211> LENGTH: 21 <212> TYPE: DNA <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: DNA primer <400> SEQUENCE: 65 catgtgccgc cccatcgctg c 21 <210> SEQ ID NO 66 <211> LENGTH: 58 <212> TYPE: PRT <213> ORGANISM: Artificial sequence <220> FEATURE: <223> OTHER INFORMATION: Modified Ebola virus GP

We claim:

1. An isolated monoclonal antibody, comprising:

- a heavy chain variable region  $(V_H)$  comprising a heavy chain complementarity determining region (HCDR)1, a 20 HCDR2, and a HCDR3 of the  $V_H$  set forth as SEQ ID NO: 1 (EVB114  $V_H$ ), and a light chain variable region  $(V_L)$  comprising a light chain complementarity determining region (LCDR)1, a LCDR2, and a LCDR3 of the  $V_L$  set forth as SEQ ID NO: 2 (EVB114  $V_L$ ); or 25
- a  $V_H$  comprising a HCDR1, a HCDR2, and a HCDR3 of the  $V_H$  set forth as SEQ ID NO: 3 (EVB100  $V_H$ ), and a  $V_L$  comprising a LCDR1, a LCDR2, and a LCDR3 of the  $V_L$  set forth as SEQ ID NO: 4 (EVB100  $V_L$ ); and
- wherein the monoclonal antibody or antigen binding 30 fragment specifically binds to Ebola virus glycoprotein (GP).
- 2. The antibody of claim 1, wherein:
- the HCDR1, the HCDR2, and the HCDR3 comprise amino acids 26-33, 51-57, and 96-108 of SEQ ID NO: 35 1, respectively, and the LCDR1, the LCDR2, and the LCDR3 comprise amino acids 27-32, 50-52, and 89-97 of SEQ ID NO: 2, respectively; or
- the HCDR1, the HCDR2, and the HCDR3 comprise amino acids 26-33, 51-57, and 96-114 of SEQ ID NO: 40 3, respectively, and the LCDR1, the LCDR2, and the LCDR3 comprise amino acids 26-31, 49-51, and 88-95 of SEQ ID NO: 4, respectively.

3. The antibody of claim 1, wherein:

- the  $V_H$  comprises an amino acid sequence at least 80% 45 identical to the sequence set forth as SEQ ID NO: 1 and/or the  $V_L$  comprises an amino acid sequence at least 80% identical to the sequence set forth as SEQ ID NO: 2; or
- the  $V_H$  comprises an amino acid sequence at least 80% 50 identical to the sequence set forth as SEQ ID NO: 3, and/or the  $V_L$  comprises an amino acid sequence at least 80% identical to the sequence set forth as SEQ ID NO: 4.

4. The antibody of claim 1, wherein:

- the  $V_H$  comprises the amino acid sequence set forth as SEQ ID NO: 1, and/or the  $V_L$  comprises the amino acid sequence set forth as SEQ ID NO: 2; or
- the  $V_H$  comprises the amino acid sequence set forth as SEQ ID NO: 3, and/or the  $V_L$  comprises the amino acid 60 sequence set forth as SEQ ID NO: 4.

5. The antibody of claim 1, wherein the  $V_H$  and  $V_L$  comprise or consist of the amino acid sequences set forth as SEQ ID NOs: 1 and 2, respectively, or SEQ ID NOs: 3 and 4, respectively.

**6**. The antibody of claim **1**, comprising the  $V_H$  comprising the HCDR1, the HCDR2, and the HCDR3 of the  $V_H$  set forth

as SEQ ID NO: 1, and the  $V_L$  comprising the LCDR1, the LCDR2, and the LCDR3 of the  $V_L$  set forth as SEQ ID NO: 2, and further comprising

- a M31S substitution according to kabat positioning in the  $V_{H}$ ;
- a D30S substitution according to kabat positioning in the  $\mathrm{V_{\it L}};$  or
- a M31S substitution according to kabat positioning in the  $V_H$  and a D30S substitution according to kabat positioning in the  $V_L$ .

7. The antibody of claim **6**, wherein the HCDR1, the HCDR2, the HCDR3, the LCDR1, the LCDR2, and the LCDR3 comprise SEQ ID NOs: 38, 33, 34, 35, 36, and 37, respectively, SEQ ID NOs: 32, 33, 34, 39, 36, and 37, respectively, or SEQ ID NOs: 38, 33, 34, 39, 36, and 37, respectively.

**8**. The antibody of claim **6**, wherein the  $V_H$  and the  $V_L$  comprises the amino acid sequences set forth as SEQ ID NOs: 28 and 2, respectfully, SEQ ID NOs: 1 and 29, respectfully, or SEQ ID NOs: 28 and 29, respectfully.

9. The antibody of claim 1, comprising a human frame-work region.

**10**. The antibody of claim **1**, comprising a human constant region.

**11**. The antibody of claim **1**, wherein the antibody is an IgG, IgM or IgA.

**12**. The antibody of claim **1**, wherein the antibody is an IgG1 and comprises a human constant region.

**13**. The antibody of claim **1**, comprising a recombinant constant region comprising one or more modifications that increase binding to the neonatal Fc receptor and/or increase antibody-dependent cell cytotoxicity (ADCC).

14. The antibody of claim 13, wherein the antibody is a human IgG1 and comprises a recombinant constant region comprising M428L and N434S mutations to increase binding to the neonatal Fc receptor.

15. An antigen binding fragment that specifically binds to 55 the extracellular domain Ebola virus GP, comprising the  $V_H$ and the  $V_L$  of the antibody of claim 1.

16. The antigen binding fragment of claim 15, wherein the antigen binding fragment is a Fv, Fab,  $F(ab')_2$ , scFV or a scFV<sub>2</sub> fragment.

17. A bispecific antibody that specifically binds to the extracellular domain Ebola virus GP, comprising the  $V_H$  and the  $V_I$  of the antibody of claim 1.

**18**. The antibody or antigen binding fragment of claim **1**, wherein the monoclonal antibody or antigen binding fragment neutralizes Ebola virus.

**19**. The antibody or antigen binding fragment of claim **14**, wherein the Ebola virus is Zaire Ebola virus.

10

**20**. The antibody or antigen binding fragment of claim **1**, wherein the Ebola virus GP is a Zaire Ebola virus GP.

**21**. The antibody or antigen binding fragment of claim **16**, wherein the Zaire Ebola virus GP comprises the amino acid sequence set forth as SEQ ID NO: 15.

**22**. The antibody or antigen binding fragment of claim **1**, linked to an effector molecule or a detectable marker.

23. The antibody or antigen binding fragment of claim 22, wherein the detectable marker is a fluorescent, enzymatic, or radioactive marker.

24. An antibody or an antigen binding fragment thereof that binds to the same epitope as the antibody of claim 1, wherein the antibody or antigen binding fragment thereof neutralizes Ebola virus.

**25**. An isolated nucleic acid molecule encoding the  $V_H$  <sup>15</sup> and/or the  $V_L$  of the antibody or antigen binding fragment of claim **1**.

**26**. The nucleic acid molecule of claim **25**, wherein the nucleic acid molecule is a recombinant nucleic acid molecule.

27. The nucleic acid molecule of claim 25, comprising a cDNA molecule encoding the antibody or antigen binding fragment.

**28**. The nucleic acid molecule of claim **25**, wherein the  $V_H$  and/or the  $V_L$  of the antibody or antigen binding frag-<sup>25</sup> ment comprise the nucleic acid sequences set forth as

SEQ ID NOs: 7 and 8, respectively, or degenerate variants thereof; or

SEQ ID NOs: 9 and 10, respectively, or degenerate variants thereof. 30

**29**. The nucleic acid molecule of claim **25**, operably linked to a promoter.

**30**. An expression vector comprising the nucleic acid molecule of claim **25**.

**31**. A pharmaceutical composition for use in treating or <sup>35</sup> inhibiting an Ebola virus infection, comprising:

a therapeutically effective amount of the antibody of claim 1, an antigen binding fragment thereof, a nucleic acid molecule encoding the antibody or antigen binding fragment, or an expression vector comprising the nucleic acid molecule; and

a pharmaceutically acceptable carrier.

**32**. The pharmaceutical composition of claim **31**, wherein the composition is sterile and/or is in unit dosage form or a multiple thereof.

**33**. A method of detecting an Ebola virus infection in a subject, comprising:

contacting a biological sample from the subject with the antibody of claim 1 or an antigen binding fragment thereof under conditions sufficient to form an immune <sup>50</sup> complex; and

detecting the presence of the immune complex on the sample, wherein the presence of the immune complex on the sample indicates that the subject has the Ebola virus infection.

**34**. A method of preventing or treating an Ebola virus infection in a subject, comprising administering to the subject a therapeutically effective amount of the antibody of claim 1, an antigen binding fragment thereof, a nucleic acid molecule encoding the antibody or antigen binding fragment, or an expression vector comprising the nucleic acid molecule, thereby preventing or treating the Ebola virus infection.

**35**. The method of claim **34**, further comprising administering to the subject one or more additional antibodies or antigen binding fragments that specifically bind to Ebola virus GP and neutralize Ebola virus, or one or more nucleic acid molecules encoding the additional antibodies or antigen binding fragments.

**36.** The method of **35**, comprising administering to the  $_{20}$  subject a pharmaceutical composition comprising a first antibody and a second antibody, wherein

- the first antibody comprises the  $V_H$  comprising the HCDR1, the HCDR2, and the HCDR3 of the  $V_H$  set forth as SEQ ID NO: 1, and the  $V_L$  comprising the LCDR1, the LCDR2, and the LCDR3 of the  $V_L$  set forth as SEQ ID NO: 2; and
- the second antibody comprises the  $V_H$  comprising the HCDR1, the HCDR2, and the HCDR3 of the  $V_H$  set forth as SEQ ID NO: 3, and the  $V_L$  comprising the LCDR1, the LCDR2, and the LCDR3 of the  $V_L$  set forth as SEQ ID NO: 4.

**37**. The method of claim **36**, wherein the first antibody comprises the  $V_H$  and  $V_L$  set forth as SEQ ID NOs: 1 and 2, respectively, and the second antibody comprises the  $V_H$  and  $V_L$  set forth as SEQ ID NOs: 3 and 4, respectively.

**38**. The method of claim **34**, wherein the Ebola virus is Ebola virus Zaire.

**39**. A method of producing an antibody or antigen binding fragment that specifically binds to Ebola virus GP, comprising:

expressing in a host cell:

- one or more nucleic acid molecules encoding antibody heavy and light chains comprising the  $V_H$  and the  $V_L$ of the antibody of claim 1; or
- one or more nucleic acid molecules encoding an antigen binding fragment comprising the  $V_H$  and the  $V_L$ ; and

purifying the antibody or antigen binding fragment;

thereby producing the antibody or antigen binding fragment.

\* \* \* \* \*